



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XX International Grassland Congress

The Effect of Fertiliser Treatment on the Development of Rangelands in Argentina

Ezequiel M. Latorre

Universidad Nacional del Centro de la Provincia de Buenos Aires, Argentina

Mónica B. Sacido

Universidad Nacional del Centro de la Provincia de Buenos Aires, Argentina

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium3/112>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

The effect of fertiliser treatment on the development of rangelands in Argentina

E.F. Latorre and M.B. Sacido

The Faculty of Agronomy, Azul National University of the Centre of the Buenos Aires Province, Argentina. CP: 7300. Commission of Investigation Scientific, Buenos Aires Province, Argentina. Email: msacido@faa.unicen.edu.ar

Keywords: rangelands, fertiliser application, productivity

Introduction In Argentina grazing of rangelands may result in a decrease in winter gramineous species with an increase in summer weeds such as *Cynodon dactylon*. *Lolium multiflorum* is an important forage resource for grazing in the autumn, winter and spring. A delay in its emergence may occur because of summer weeds, which reduces the germination rate. The proportion of the seed bank as ryegrass allows the recovery of natural grassland and facilitates an increase in the productivity of livestock. The objective of this study was the evaluation of the impact of application of fertiliser in the short term on the relationship with botanical composition at different herbage availabilities.

Materials and methods The experimental site was a typical natural pasture in the Flooding Pampas, with three level of availability of herbage mass (high, >5000 kg dry matter (DM)/ha; medium, between 5000 and 3000 kg DM/ha; and low <3000 kg DM/ha). The treatments were with and without fertilization. On the fertilized treatment the technique of promotion of rangeland (“promoción de pasturas”) involved the annual application of ammonium phosphate (80 kg/ha) and glyphosate (5 l/ha). Measurements were made at 45 and 75 days after fertiliser application of floristic composition, herbage availability and species abundance. Correlation (software STADISTIC 5.0) and similarity coefficients by the Czekanowski, Dice, Sørensen Index were determined.

Results The floristic composition, at the three different availabilities of herbage, in both treatments (-F without fertilized and F fertilized) is shown in Figure 1. It shows the variation in floristic composition, especially for *Lolium multiflorum* Lam, between the fertiliser and no fertiliser treatments ($P<0.01$). Within a fertiliser treatment there were no significant differences in floristic composition between the three levels of availability. In relation to differences between fertiliser treatments in floristic composition, these were significant ($P<0.05$) using a Dependent Samples test.




Figure 1 The floristic composition of the pastures as affected by three levels of availability of herbage and fertilizer treatment (-F, without fertilizer; and F, fertilised)

Conclusion The composition in the forage offered differed according to the fertiliser application regime. The high fertilizer treatment increased the composition of forage species, especially *Lolium multiflorum*, and decreased the proportion of weeds.