



Molecular Breeding of White Clover for Transgenic Resistance to Alfalfa Mosaic Virus and Natural Resistance to Clover Yellow Vein Virus

P. Chu
CSIRO, Australia

G. Zhao
Gansu Agricultural University, China

G. C. Spangenberg
La Trobe University, Australia

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium5/93>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

Molecular breeding of white clover for transgenic resistance to *Alfalfa mosaic virus* and natural resistance to *Clover yellow vein virus*

P. Chu¹, G. Zhao^{2,3} and G.C. Spangenberg²

¹CSIRO Plant Industry, GPO Box 1600, Canberra, ACT 2601, Australia; ²Primary Industries Research Victoria, Plant Biotechnology Centre, La Trobe University, Bundoora, Victoria 3086, Australia; ³Department of Grassland Science, Gansu Agricultural University, Lanzhou 730070, China. Email: paul.chu@csiro.au

Keywords: virus resistance, white clover, alfalfa mosaic virus, clover yellow vein virus

Introduction *Trifolium repens* L. (white clover) is one of the most important pastoral plants in temperate Australia. Its productivity and persistence is being reduced significantly by *Alfalfa mosaic virus* (AMV), *Clover yellow vein virus* (CIYVV) and *White clover mosaic virus* (WCIMV). These viruses are also widespread in other legumes and are inflicting large economic losses to farmers throughout the world (Campbell, 1984). To reduce the economic impact of these viruses, white clover plants resistant to both CIYVV and AMV are being developed for future commercial release. Since introducing viral transgenes from two or more viruses into a transgenic plant has the potential threat of viral recombination, we have decided to develop white clover with transgenic resistance to AMV and natural resistance to CIYVV.

Materials and methods Transgenic white clover plants expressing the AMV coat protein (CP) gene and showing immunity to that virus have been produced. White clover plants with natural resistance to CIYVV were identified by screening various cultivars of the white clover against different isolates of the virus. To combine the transgenic AMV immunity and natural CIYVV resistance into a single transgenic white clover cultivar, crosses between singly AMV immune and CIYVV resistant white clover plants were performed. The offspring plants were subjected to molecular analysis and infectivity screens to identify plants carrying AMV CP and showing immunity to both viruses. The best of three AMV + CIYVV resistant genotypes was selected for crossing with 12 Sustain-type elite plants in order to produce a new white clover cultivar with double virus resistance.

Results Crosses between singly AMV immune and CIYVV resistant white clover plants produced offspring plants that were resistant to both viruses. After crossing with the Sustain-type parents, almost 100% of the offspring plants carrying the AMV CP showed immunity to that virus, irrespective of the Sustain-type parent used. In contrast, there were significant variations in the percentage of CIYVV resistant progenies produced by the different Sustain-type parents (Figure 1). Diallele crossing between these first generation double virus resistant progenies (T1 plants) increased the proportion of CIYVV resistant plants among the T2 population from 34% to 49% (Figure 2). Furthermore, T2 population showed a higher proportion of plants with milder symptoms than plants in the T1 population.

Figure 1





Figure 2



Conclusions The results showed that it is possible to produce white clover plants with transgenic resistance to AMV and natural resistance to CIYVV. The differential inheritance of CIYVV resistance from different Sustain genotypes suggests that multiple resistance genes may be involved in the natural CIYVV resistance.

References

Campbell C.L. and J.W. Moyer (1984). Yield responses of 6 white clover clones to virus infection under field condition. *Plant Disease*, 68, 1033-1035.