



Population Genetics of Perennial Ryegrass (*Lolium Perenne* L.): Differentiation of Pasture and Turf Cultivars

M. P. Dobrowolski
Hamilton Centre, Australia

N. R. Bannan
Hamilton Centre, Australia

R. C. Ponting
La Trobe University, Australia

J. W. Forster
La Trobe University, Australia

K. F. Smith
Molecular Plant Breeding Cooperative Research Centre, Australia

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium5/144>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

Population genetics of perennial ryegrass (*Lolium perenne* L.): differentiation of pasture and turf cultivars

M.P. Dobrowolski^{1,3}, N.R. Bannan^{1,3}, R.C. Ponting^{2,3}, J.W. Forster^{2,3} and K.F. Smith^{1,3}

¹Primary Industries Research Victoria, Hamilton Centre, Hamilton, Victoria 3300, Australia ²Primary Industries Research Victoria, Plant Biotechnology Centre, La Trobe University, Bundoora, Victoria 3086, Australia

³Molecular Plant Breeding Cooperative Research Centre, Australia. Email: mark.dobrowolski@dpi.vic.gov.au

Keywords: perennial ryegrass, population genetics, simple sequence repeat markers, cultivar differentiation

Introduction Cultivar differentiation using molecular markers to assess genetic variation may be of value in obtaining or protecting plant breeders rights. A knowledge of genetic variation and how it is structured within perennial ryegrass (*Lolium perenne* L.) populations will also help us understand the consequences to fitness and adaptation when implementing molecular breeding strategies. In a study of the population genetic structure of a number of perennial ryegrass varieties we examined the cultivar differentiation potential of marker technology.

Materials and methods We used 26 genomic DNA-derived simple sequence repeat (SSR) marker loci. Based on genetic mapping in the p150/112 reference population, these loci are distributed across each of the 7 linkage groups (chromosomes) of the perennial ryegrass genome. Two groups of populations were compared to assess the capacity to distinguish them using the SSR marker loci: pasture varieties derived from the Kangaroo Valley ecotype from New South Wales, Australia, and turf varieties from a range of sources.

Results An analysis of molecular variance (AMOVA) showed that 90% of the variation is located within the cultivars, 8% among cultivars within the pasture and turf groups and 2% between the pasture and turf groups. Individuals in this analysis could not reliably be allocated to their source cultivar or group (pasture or turf) on the basis of their 26 loci SSR genotype (Figure 1). Due to the obligate outbreeding nature of perennial ryegrass, relatively high levels of diversity are maintained within cultivars that have undergone selection in their development.

Definition and identification of cultivars based on molecular markers may depend on careful selection of markers, informed by the population structure and breeding history of the cultivars to be distinguished. This data has important implications for the design and selection of metapopulations for association genetics analyses in outbreeding grass species.




Figure 1 Log-log plot of the likelihood of a perennial ryegrass individual belonging to the population defined by the Kangaroo Valley-derived cultivars or the population defined by the turf-type cultivars. The probability that an individual is derived from the defined populations is calculated from the observed allele frequencies in the population excluding that individual if necessary, and assuming Hardy-Weinberg equilibrium and panmixia.