

Physical and Quality Seed Traits Observed in New Pigeon Pea (*Cajanus Cajan*) Hybrids

R. Usberti

Plant Protection Agency, Brazil

J. A. Usberti Jr.

Campinas Agronomic Institute, Brazil

R. H. Aguiar

Campinas Agronomic Institute, Brazil

F. C. Simoes


Campinas Agronomic Institute, Brazil

F. N. Cavalcante

Campinas Agronomic Institute, Brazil

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/themeA/57>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

Presenter Information

R. Usberti, J. A. Usberti Jr., R. H. Aguiar, F. C. Simoes, F. N. Cavalcante, M. Harris, Maria T. Colozza, and J. C. T. Freitas

Physical and quality seed traits observed in new pigeon pea (*Cajanus cajan*) hybrids

R. Usberti¹, J.A. Usberti Jr.², R.H. Aguiar³, F.C. Simoes³, F.N. Cavalcante, Jr.³, M. Harris³, M.T. Colozza⁴ and J.C.T. Freitas⁴

¹Plant Protection Agency, P.O. Box 960, CEP 13073-001, Campinas, Brazil, Email: usberti@cati.sp.gov.br,

²Agronomic Institute, Campinas, Brazil, ³Faculty of Agricultural Engineering, Campinas, Brazil, ⁴Zootechnic Institute, Nova Odessa, Brazil

Keywords: hardseededness, germination, purity

Introduction Pigeon pea seed production may be affected by factors such as % of pure seeds, mean seed weight, incidence of pests and diseases and environmental stresses. Harvested seeds from different cultivars may also vary in germination %, hardseededness and germination speed. Hardseededness (seed coat impermeability to water) commonly occurs in forage legume species (Hopkinson, 1993). There is considerable variation among different entries for seed characters but this is not considered within genetic materials. This research analysed harvested seeds of selected individuals of two segregating F₂ pigeon pea populations for the above cited traits and assessed the range of variation for them resulting from the hybridisation process.

Materials and methods Selected individuals of two segregating F₂ pigeon pea populations, arisen from previous artificial crossings, had their seeds carefully harvested in special cloth bags and transferred to the laboratory. After a manual cleaning for separating seeds damaged by insects and fungi (Kashyap & Punia, 1996), the following physical seed traits were scored: physical purity (percentage of pure seeds/sample) and number of seeds/10g (ISTA, 1985). Seed standard germination tests were then performed using 4x50 replicates and the number of hard seeds scored at the first count (4th day), when their coats were partially removed by forceps; mean germination time test (T₅₀) was also carried out (Alvarado & Bradford, 1988).

Results In both segregating F₂ populations, most of the genotypes analysed showed low incidences of damaged seeds by insects and fungi (0-10%). For physical purity, there was a marked concentration of individuals in the ranges of 61-90 and >90% (population 1 and 2, respectively). There was a clear difference in the number of seeds/10g between the populations, the first showing high concentration of individuals in the range 91-100 and >100 while the second had a more uniform distribution (Table 1). Most of the selected individuals of both populations (75 and 60%, respectively) showed hardseededness, varying from <20 to 40%, while population 1 showed a higher frequency of individuals in the germination percentage interval of 61-100%. Both segregating populations presented similar distributions of their individuals as to mean germination times.

Table 1 Frequency distribution of physical and physiological seed traits in two pigeon pea (*Cajanus cajan*) segregating populations

| SP | Damaged seeds (%) | | | | | Physical purity (%) | | | Number of seeds/10g | | | |
|----|--------------------|-------|-------|-------|-----------------|---------------------|-------|-----|--|-------|--------|------|
| | Insects | | | Fungi | | | | | | | | |
| | 0-10 | 11-20 | >20 | 0-10 | 11-20 | 30-60 | 61-90 | >90 | 71-80 | 81-90 | 91-100 | >100 |
| 1 | 16 | 5 | 3 | 18 | 6 | 4 | 13 | 7 | - | 2 | 9 | 13 |
| 2 | 15 | 3 | 2 | 17 | 3 | 1 | 6 | 13 | 2 | 8 | 7 | 3 |
| | Hardseededness (%) | | | | Germination (%) | | | | Mean germination time (T ₅₀ , days) | | | |
| | <20 | 21-40 | 41-60 | >60 | <40 | 41-60 | 61-80 | >81 | 4-5 | 5-6.3 | >6.3 | |
| | 1 | 4 | 11 | 6 | 3 | 1 | 16 | 6 | 9 | 7 | 8 | |
| 2 | 6 | 6 | 4 | 4 | 2 | 9 | 7 | 2 | 6 | 6 | 8 | |

SP, segregating population with SP1 and SP2 having 24 and 20 individuals respectively; physical purity is pure seed weight/working seed sample weight

Conclusions The hybridisation process used was effective as to significantly increasing the available genetic variability for several seed quality traits, making feasible effective selection for them in the near future.

References

- Alvarado, N.A. & K.J. Bradford, (1988). Priming and storage of tomato (*Lycopersicon lycopersicum*) seeds. I. Effects of storage temperature on germination rate and viability. *Seed Science & Technology*, 16, 601-612.
- Hopkinson, J.M. (1993). Tropical pasture establishment .2. Seed characteristics and field establishment. *Tropical Grasslands*, 27, 276-290.
- ISTA, International Seed Testing Association. (1985). International rules for seed testing. RULES 1985. *Seed Science and Technology*, 13, 299-355; 356-513.
- Kashyap, R.K. & R.C. Punia, (1996). Seed quality as influence by pod infesting insect pests of pigeon pea *Cajanus cajan* L. Millsp. *Seed Science and Technology*, 23, 873-876.