



Yield and Quality Parameters of an Interspecific Hybrid *Pennisetum Purpureum* Schum. (Elephant-Grass) X *Pennisetum Glaucum* (L.) R. Br. Stuntz (Pearl Millet)

J. A. Usberti Jr.
Campinas Agronomic Institute, Brazil


R. Usberti
Plant Protection Agency, Brazil

P. B. Alcantara
Zootecnia Institute, Brazil

V. B. G. Alcantara
Zootecnia Institute, Brazil

R. A. Possenti
Zootecnia Institute, Brazil

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 [next page for additional authors](#)

Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/themeA/62>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005. The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

Presenter Information

J. A. Usberti Jr., R. Usberti, P. B. Alcantara, V. B. G. Alcantara, R. A. Possenti, and M. A. C. Lucena

Yield and quality parameters of an interspecific hybrid *Pennisetum purpureum* Schum. (elephant-grass) x *Pennisetum glaucum* (L.) R. Br. Stuntz (pearl millet)

J.A. Usberti, Jr.¹, R. Usberti², P.B. Alcantara³, V.B.G. Alcantara³, R.A. Possenti³ and M.A.C. Lucena³

¹Agronomic Institute, Campinas, Brazil. P.O. Box 28, CEP 13073-001, Email: usberti@iac.sp.gov.br, ²Plant Protection Agency, Campinas, Brazil, ³Zootechnic Institute, Nova Odessa, Brazil

Keywords: recurrent selection, interspecific hybrid, forage yielding and quality traits

Introduction Elephant-grass is a tropical forage grass used either as a supplement fodder or for direct grazing. It usually shows regular nutritive value (6-13% crude protein, CP, and 55-60% forage digestibility) (Alcantara *et al.*, 1981). Most of the available cultivars produce no viable seeds. On the other hand, pearl millet has high seed yielding potential along with high quality forage (>15% CP and 70% forage digestibility). However, it shows poor forage production, low field persistence under grazing and low regrowth potential after cutting or grazing. During the 90's, an interspecific hybrid between the two species was developed, trying to combine the elephant-grass adaptability and forage yielding potential with the pearl millet forage quality and seed yielding potential (Schank *et al.*, 1993; Schank, 1996). The new genetic material was able to produce viable seeds in variable amounts (Diz *et al.*, 1995). The main aim of this research was to produce selected populations with high phenotypic uniformities, showing high average forage production and quality.

Materials and methods Cutting and grazing types populations (CT and GT) were established, through the selection of 250 individual plants within the original F₂ population. The CT was made up of tall plants with low-tillering potential, long and broad leaves, thick stems with long internodes and erect growth habit while GT presented small and high-tillered plants, short and narrow leaves, thin stems with short internodes and prostrated growth habit. During two consecutive selection cycles, both were allowed free intercrossing in isolated fields. Plant height, tiller number, leaf length and width, stem diameter, internode length, inflorescence axis length, inflorescence length and leaf/stem ratios were scored, as well as dry matter (DM) production; leaf and stem CP contents; leaf+stem CP production and leaf and stem digestibility.

Results The CT and GT populations showed marked gains in phenotypic uniformities, while maintaining their original differential traits. Similar results were showed for internode length, inflorescence axis length and inflorescence length (data not shown here). However, there were differences among them for leaf/stem ratios (Table 1). Both selected populations showed DM production higher than the control with GT outstanding. The highest leaf and stem CP contents were also in GT, but there were no significant differences in leaf and stem digestibility. Therefore, GT and CT gave higher overall CP production than the original F₂ population.

Table 1 Mean results for selected morpho-agronomic, yield and quality traits in two selected populations compared with original F₂ population for 2004 selection cycle

| Population | Plant height | Tiller number | Stem diameter | Leaf stem ratio | DM yield (t/ha) | Leaf CP (%) | Stem CP (%) | Leaf digestibility (%) | Stem digestibility (%) | CP yield (kg/ha) |
|-------------------------|--------------|---------------|---------------|-----------------|-----------------|-------------|-------------|------------------------|------------------------|------------------|
| Cutting type | 3.41a | 18.0b | 1.80a | 32.3b | 15.2a | 8.1b | 4.3b | 71.9a | 56.3a | 779b |
| Grazing type | 3.01b | 34.4a | 1.21b | 38.2a | 14.1a | 9.4a | 5.8a | 70.9a | 59.0a | 965a |
| Original F ₂ | 3.32a | 19.5b | 1.73a | 26.5a | 11.8b | 7.8b | 4.6b | 69.0a | 57.4a | 620c |
| Mean | 3.25 | 24.0 | 1.58 | 32.3 | 13.6 | 8.4 | 4.9 | 70.6 | 57.6 | 788 |
| C.V (%) | 7.09 | 10.4 | 14.1 | 15.2 | 17.2 | 12.0 | 12.8 | 10.1 | 10.3 | 15.4 |

C.V. (%) = coefficient of variation; b) Means followed by different small letters, in the same column, are statistically different according to the Duncan test at $p < 0.05$

Conclusions The phenotypic uniformities of both selected populations were improved; forage yielding and quality parameters were enhanced, making feasible the use of those populations in cultivated pastures.

References

- Alcantara, P.B., V.B.G. Alcantara & J.E. Almeida (1981). *Proceedings of the Fourteenth International Grassland Congress*, 91.
- Diz, D.A., S. C. Schank & D. S. Wofford (1995). Defoliation effects and seed yield components in pearl millet X elephantgrass hybrids. *Agronomy Journal*, 87, 56-62.
- Schank, S. C. & D. P. Chynoweth (1993). The value of triploid, tetraploid, and hexaploid Napier grass derivatives as biomass and/or forage. *Tropical Agriculture*, 70, 83-87.
- Schank, S. C., D. A. Diz, P. J. Hogue, C. V. Vann. (1996). Evaluation of pearl millet X elephantgrass hybrids for use as high quality forage for livestock. *Soil and Crop Science Society of Florida Proceedings*, 55, 120-121.