

## FACILITATING AND COMPLICATING FACTORS IN COPING WITH FEAR OF COVID-19

Mehmet Yalçın<sup>1</sup>, Eyüp Sabır Erbiçer<sup>2</sup> & Ercan Akın<sup>2</sup>

<sup>1</sup>Department of Psychology, Hasan Kalyoncu University, Gaziantep, Turkey

<sup>2</sup>Department of Psychological Counseling and Guidance, Hacettepe University, Ankara, Turkey

received: 2.3.2022;

revised: 11.5.2022;

accepted: 8.6.2022

### SUMMARY

**Background:** The Covid-19 pandemic has affected almost all countries' people and affected them in many ways. Turkey has been one of the countries most affected by the Covid-19 pandemic. In this study, it is aimed to examine facilitating and complicating factors in coping with the fear of COVID-19.

**Subjects and methods:** This study was conducted with 530 people (305 females, 225 males) living in different regions of Turkey and reached online via google e-forms. Sociodemographic information of the participants was collected, and the Fear of Covid-19 Scale (FCV-19S), Short Version of the Intolerance of Uncertainty Scale (IUS-12), The State-Trait Anxiety Inventory (STAI), and The World Health Organization Quality of Life (WHOQOL-BREF) scales were administered. All statistical analyzes were performed using IBM SPSS Statistics. T-test, One Way Anova and Linear Regression analysis were used to analyze the collected data.

**Results:** In this study, the participants reported that watching movies, reading books, and listening to music were the most important facilitating factors in coping with Fear of COVID-19. Covid-19 fear scores were analyzed in terms of some demographic variables. According to the results, there was no difference between COVID-19 fear scores in terms of gender, having a chronic disease or not, knowing someone diagnosed with Covid-19 in their immediate vicinity, and the region they lived in. We found that Prospective anxiety, inhibitory anxiety, state anxiety, and psychological health were significant predictors of Covid-19 fear. However, trait anxiety, physical health, social relationships and environment were not significant predictors of Covid-19 fear. All these variables together explained 25% of the variance in the model.

**Conclusion:** Although this study has some limitations, it has an important place in the literature in revealing the facilitating and complicating factors in dealing with the fear of Covid -19. It mediates the provision of recommendations to policy makers and mental health professionals for providing psychological support services to individuals adversely affected by the COVID-19 pandemic process.

**Key words:** COVID-19 – fear - anxiety - intolerance to uncertainty

\* \* \* \* \*

### INTRODUCTION

Coronaviruses are a family of viruses that can spread from animals to humans and cause many diseases (Weiss & Leibowitz 2011). It is estimated that coronaviruses, whose origin is still unknown, have circulated the world for centuries (Woo et al. 2012, Huynh et al. 2012). SARS-CoV and MERS-CoV, which are from this family of viruses, caused epidemics at the beginning of this century and infected more than 10,000 people (Hui & Zumla 2019, WHO 2003). A series of unexplained cases of pneumonia occurred in Wuhan, China's Hubei province on 31 December 2019, and these cases were associated with the seafood and wet animal wholesale market (Bogoch et al. 2020, Lu et al. 2020). To date, the COVID-19 pandemic caused by the SARS-CoV-2 virus has affected more than 165 million (22 May 2021) people worldwide (WHO 2020).

This pandemic has affected both physical health and mental health significantly (Anjum et al. 2020, Brooks et al. 2020, Shigemura et al. 2020, Abuhmaidan & Al-Majali 2020, Ren & Guo 2020). Limitations on leaving home and other issues, having to leave certain routines, reduced social and physical contact with others have

caused stress and frustration that cause psychological and physical problems in people (Blendon et al. 2004, Braunack-Mayer et al. 2013, Šljivo et al. 2020, Karasar & Canli 2020). For this reason, with the epidemic, mental health professionals all over the world have started to serve individuals who have mental problems, as well as those who are mentally affected by the pandemic. The COVID-19 outbreak caused both death risk and unbearable psychological pressure in humans worldwide (Xiao 2020, Duan & Zhu 2020). When the threat of epidemic increased, the cancellation of travel plans, social isolation, media news, and people flocking to markets to meet their needs created an atmosphere of anxiety and depression all over the world (Ho et al. 2020). Wang et al. (2020) reported that there was no improvement in individuals' stress, anxiety, and depression levels at the beginning of the COVID-19 epidemic and 4 weeks later. It has been reported that especially isolation and loneliness cause an increase in individuals' depression, anxiety and anger levels and affect physical health negatively (Huremović 2019, Leigh-Hunt et al. 2017). In such crisis times, an increase in mental health problems is expected. It has been reported that previous pandemics and epidemics significantly

affect the mental health of individuals and society. It has been reported that especially individuals who had previously experienced mental problems face higher rates of psychological problems and problems such as fear of disease (Chua et al. 2004, Chong et al. 2004, Mak et al. 2009).

Fear and anxiety are physiologically basic emotions that involve activating the nervous system's "fight or flight" response and enable us to react quickly when faced with imminent threat. In this pandemic period, the most important causes of fear and anxiety in the society could be sorted as contagiousness of the infection, being invisible, uncertainties regarding vaccination and treatment and relatively high mortality rate (Pappas et al. 2009). Individuals' psychological responses to these emerging stressful situations vary in a wide range from extreme fear to emotionlessness, with the effect of personality traits such as trait anxiety, harm avoidance and intolerance to uncertainty (Taylor 2019).

Trait anxiety refers to the individuals' tendency to continue and experience negative emotions such as fear and anxiety. People with high level trait anxiety tend to see the world as dangerous and threatening. Therefore, these people are very scared and extremely anxious, especially in a sense of danger (Spielberger 1979). Studies on the SARS outbreak have reported that trait anxiety was a significant predictor of epidemic anxiety (Cheng & Cheung 2005).

Another dimension that affects this tendency to experience anxiety and fear is intolerance to uncertainty (McEvoy & Mahoney 2013). The pandemic process contains uncertainties on a wide variety of issues. The unpredictability of infection was also a source of great fear in the Spanish flu epidemic (Graves 1969). In times of pandemics, it is important for people to be able to tolerate or accept uncertainty. It has been suggested that intolerance to uncertainty will make a particularly significant contribution to the anxiety and fear associated with the epidemic (Taylor 2019). Individuals with a high level of intolerance to uncertainty tend to be anxious because they feel they have limited control over a threatening situation such as a pandemic (Taha et al. 2014).

The number of people who need psychiatric help since the beginning of the pandemic has been increasing day by day, and this was causing some changes in psychiatric practices because the pandemic now functions as a new trauma or stressor for mental health professionals (Kang et al. 2020). The increased fear and anxiety of individuals due to outbreaks causes them to be more aware of various normal or benign physical symptoms and evaluate them as a precursor of a severe disease.

In this study, it is aimed to examine facilitating and complicating factors in coping with the fear of COVID-19. For this purpose, we tried to reveal the people most affected by the pandemic and the factors that people who are less affected by the pandemic use to cope.

## SUBJECTS AND METHODS

### Subjects

The average age of 530 participants in the study was 30.34, with a standard deviation of 7.13, and it ranged from 21-66. 305 (57.5%) of the participants are female, and 225 (42.5%) are male. Participants were 530 people living in 11 different cities in Turkey. Besides, more than half of the participants are single (64.5%) and undergraduate (58.1%). The demographic information of the participants is given in detail in Table 1.

### Methods

Sociodemographic data form was prepared by the researchers to obtain information of the participants such as age, gender, marital status, employment status, the number of people living with them at home, COVID-19 diagnosis status, chronic disease status, and the feel-good activities they did during the epidemic process.

**Table 1.** Sociodemographic characteristics of the sample

Age	30.34±7.13	
	n	%
Gender		
Female	305	57.5
Male	225	42.5
Highest educational level		
Primary School	21	4.0
High School	98	18.5
Undergraduate	308	58.1
Graduate	103	19.4
Marital Status		
Single	342	64.5
Married	166	31.3
Divorced	22	4.2
Living place		
Village/County	175	33.0
City	94	17.7
Metropolis	261	49.2
Occupation		
Civil Servant	216	40.8
Private Sector	129	24.3
Unemployee	74	14.0
Student	88	16.6
Housewife	23	4.3
Healthcare worker		
No	444	83.8
Yes	86	16.2
Diagnosed with COVID-19		
No	518	97.7
Yes	12	2.3
Someone diagnosed with COVID-19 in your immediate vicinity		
No	429	80.9
Yes	101	19.1
Chronic disorder		
No	465	87.7
Yes	65	12.3

Fear of Covid-19 Scale (FCV-19S) is a 7-item scale developed to measure individuals' levels of fear of COVID-19 (Ahorsu et al. 2020). The total score obtained from the scale varies between 7 and 35, and the higher the scores obtained from the scale means the increase in the level of fear accordingly. Cronbach's alpha value, which is the internal consistency coefficient of the scale, was reported as  $\alpha = 0.82$ , and the test-retest reliability value was reported as ICC = 0.72 (Ahorsu et al. 2020). The adaptation of the scale to Turkish was conducted by Satici et al. on a sample of 1304 people. Satisfactory coefficients regarding Cronbach's alpha ( $\alpha = 0.85$ ) and composite reliability (CR = 0.842) of the scale were reported in the study (Satici et al. 2020a).

Short Version of the Intolerance of Uncertainty Scale (IUS-12) was developed by Carleton et al. (2007) to measure intolerance of uncertainty, and it consists of 12 items and two sub-dimensions. The sub-dimensions are named as prospective anxiety (7 items) and inhibitory anxiety (5 items). The scale is a five-point Likert type scale. The scale was adapted to Turkish by Sarıcam and colleagues (2014). The correlation coefficient between the English and the Turkish form scores was 0.71. As a result of the exploratory factor analysis, the scale was found to explain 78.57% of the total variance with two dimensions and a 12-item structure. In the criterion validity study, the correlation between the intolerance of uncertainty and the flexibility to cope is -0.43, the correlation between the intolerance of uncertainty and education stress is 0.41. Cronbach's alpha reliability coefficient of the scale is 0.88. A higher score from the scale indicates a higher level of intolerance of uncertainty.

The State-Trait Anxiety Inventory (STAI) is a five-point Likert type scale developed by Spielberger et al. (1983) to measure state and trait anxiety levels. It consists of 40 items and two sub-dimensions. The sub-dimensions are state anxiety (20 items) and trait anxiety (20 items). The state anxiety scale determines how the individual feels at a certain moment and under certain conditions. The trait anxiety scale evaluates how the individual feels in general, regardless of the situation and conditions. The scale was adapted to Turkish by Öner and Le Compte (1985). Test-retest reliability of the Turkish form was determined between 0.71 and 0.86 for the state anxiety scale and 0.26 and 0.68 for the trait anxiety scale. The internal consistency coefficient was between 0.83 and 0.87 for the state anxiety scale and between 0.94 and 0.96 for the trait anxiety scale. A higher score from the scale indicates a higher level of anxiety.

The World Health Organization Quality of Life (WHOQOL-BREF) is a short form of the WHOQOL-100 scale. It was created by taking one question for each of the 24 sections of WHOQOL-100 and adding two questions about general health and quality of life. This tool consists of 26 questions and is scored in four dimensions: Physical health, psychological health,

social relationships, and environmental health. Cronbach alpha values were found between 0.66 and 0.84 for four subscales. Test-retest reliability for subscales was reported as 0.66 for physical health, 0.72 for psychological, 0.76 for social relationships, and 0.87 for the environment (The WHOQOL Group 1994). Fidaner et al. (1999) has adapted the WHOQOL-BREF scale to Turkish with five subscales, which is named as General Health. Cronbach alpha values were found between 0.82 and 0.92, and in the test-retest study conducted with 45 university students, the correlation values were found between 0.57 and 0.84 (Fidaner et al. 1999).

## Procedure

The data were collected from 11 different cities in Turkey using an online self-report survey delivered in Turkish by Google forms, between June 15 and July 16, 2020. All respondents provided informed consent prior to participation. The study protocol was approved by the Ethics Committee on 12 June 2020 (Session number: 40).

## Statistical Analyses

The normal distribution of the data was evaluated with the skewness and kurtosis values of the scales. Since the skewness and kurtosis values were between -1 and +1, it was accepted that the data showed a normal distribution, and parametric tests were used. Frequency and percentages for sociodemographic variables, the mean and standard deviation for age are given. Pearson Correlation analysis was used to evaluate the relationship between continuous variables, independent samples t-test to compare binary groups from continuous data, and One-Way Analysis of Variance (ANOVA) for comparing three or more groups' mean scores. Furthermore, a multiple hierarchical regression analysis was performed to find the predictors of FCV-19 scores. All statistical analyzes were performed using IBM SPSS Statistics (Statistical Package for the Social Sciences) 24.0 for Windows (SPSS Inc., Chicago, IL, USA). Statistical significance was accepted as  $p < 0.05$ .

## RESULTS

To the question "what made you feel better during lockdown period", 414 (78.1%) of the participants responded as "watching movies ", 373 (70.4%) as " being with my family ", 288 (54.3%) "reading books", 236 (44.5%) "listening to music ", 228 (43.0%) "sleeping late", 188 (35.5%) "cooking", 187 (35.3%) "taking shower", 175 (33.0%) "praying", 165 (31.1%) "watching documentaries " and 168 (29.8%) "video calling with loved ones" (Table 2).

As seen in Table 3, according to the results of analysis, there was no statistically significant difference between the FCV-19S mean scores of male and female

**Table 2.** Frequencies of answers for the question “what made you feel better during lockdown period”

	n	%
Watching movies	414	78.1
Being with family	373	70.4
Reading books	288	54.3
Listening music	236	44.5
Sleeping late	228	43.0
Cooking	188	35.5
Take shower	187	35.3
Praying	175	33.0
Watch documentary	165	31.1
Video calling with loved ones	158	29.8

participants ( $t(528) = 0.87, p=0.384$ ). Similarly, in terms of the FCV-19S mean scores, there were no statistically significant differences among participants whether there is someone diagnosed with COVID-19 in their immediate vicinity ( $t(528) = 0.21, p=0.835$ ) and whether they have a chronic disease or not ( $t(528) = -1.24, p=0.218$ ). In addition, the results of the ANOVA were not significant,  $F(2, 527) = 0.43, p=0.649$ , indicating there were no statistically significant differences among participants living in village/county, city, or metropolis (Table 4).

The results of the linear regression model were significant,  $F(8, 521)=21.16, p<0.001, R^2=0.25$ , indicating

that approximately 25% of the variance in FCV-19S is explainable by prospective anxiety, inhibitory anxiety, state anxiety, trait anxiety, physical health, psychological health, social relationships, and environment. Prospective anxiety significantly predicted FCV-19S,  $B = 0.19, t(521) = 2.97, p=0.003$ . Inhibitory anxiety significantly predicted FCV-19S,  $B = 0.17, t(521) = 2.41, p=0.016$ . State anxiety significantly predicted FCV-19S,  $B = 0.21, t(521) = 6.32, p<0.001$ . Psychological health significantly predicted FCV-19S,  $B = 0.37, t(521) = 3.70, p<0.001$ . Table 5 summarizes the results of the regression model.

## DISCUSSION

Findings about facilitating factors in dealing with the fear of COVID-19 showed that watching movies, being with family, reading books, listening to music, cooking, take shower, praying, watching documentary and video calling with loved ones were the most common activity to cope with the fear of COVID-19 during the COVID-19 pandemic. Prohibitions or some restrictions were in place across most countries to prevent the spread of the virus. With these restrictions, universities, schools and workplaces were closed. These shutdowns were a significant threat to both physical health and mental health (Metin et al. 2021, Sibley et al. 2020, Brooks et al. 2020,

**Table 3.** Independent samples t-test results for the fear of COVID-19 by gender, someone diagnosed with COVID-19 in the immediate vicinity and chronic disorder

Variable	Group	n	M	SD	t(528)	p	Cohen's d
Gender	Female	305	18.53	6.24	0.87	0.384	0.08
	Male	225	18.05	6.28			
Someone diagnosed with COVID-19 in the immediate vicinity	No	429	18.35	6.37	0.21	0.835	0.02
	Yes	101	18.21	5.79			
Chronic Disorder	No	465	18.18	6.09	-1.24	0.218	0.17
	Yes	65	19.35	7.27			

**Table 4.** Comparison of the fear of COVID-19 mean scores in terms of living region

Group	n	M	SD	F(2, 527)	p	$\eta_p^2$
Village/County	175	18.67	6.21	0.433	0.649	0.00
City	94	18.28	6.10			
Metropolis	261	18.11	6.35			

**Table 5.** Predictors of fear of COVID-19

Variable	B	SE	$\beta$	t	p	95% CI
(Intercept)	-3.66	3.95	0.00	-0.93	0.353	[-11.41, 4.09]
Prospective Anxiety	0.19	0.06	0.15	2.97	0.003	[0.06, 0.31]
Inhibitory Anxiety	0.17	0.07	0.13	2.41	0.016	[0.03, 0.30]
State anxiety	0.21	0.03	0.35	6.32	<0.001	[0.15, 0.28]
Trait anxiety	0.05	0.05	0.07	1.09	0.275	[-0.04, 0.15]
Physical health	-0.15	0.08	-0.10	-1.82	0.069	[-0.31, 0.01]
Psychological health	0.37	0.10	0.22	3.70	<0.001	[0.17, 0.57]
Social relationships	-0.21	0.13	-0.07	-1.67	0.095	[-0.46, 0.04]
Environment	0.09	0.07	0.06	1.29	0.196	[-0.05, 0.22]

Note: Results:  $F(8, 521)=21.16; p<0.001; R^2=0.25; \Delta R^2=0.23$

Fischer et al. 2020). Studies conducted during COVID-19 have provided strong evidence that staying at home due to quarantine or prohibitions increases anxiety, nervousness, stress, insomnia, anger, and uncertainty (Williams et al. 2020, Li et al. 2020b, Brooks et al. 2020, Sønderkov et al. 2020). The importance of doing activities that make individuals feel good in order to protect mental health during the pandemic process has been stated in many studies (Krause et al. 2020, Puyat et al. 2020). Based on the results of this study, it can be said that it is important to conduct informative studies by public health officials on how mental health can be protected in order to counter the negative effects of long-term isolation on physical and mental health.

Other findings of this study showed that there was no difference between COVID-19 fear scores in terms of gender, having a chronic disease or not, knowing someone diagnosed with COVID-19 in their immediate vicinity, and the region they lived in. However, many studies reported that women, those with chronic diseases, and those living in cities were more afraid. For example, in most studies using the Fear of COVID-19 scale (Ahorsu et al. 2020), it was found that women experienced more fear than men (Soraci et al. 2020, Reznik et al. 2020, Doshi et al. 2020, Tzur Bitana et al. 2020, Sakib et al. 2020, Broche-Perez et al. 2020, Tsiropoulou et al. 2020). Furthermore, Bakioğlu et al. (2020), Tzur Bitana et al. (2020) and Ping et al. (2020) have reported that individuals with chronic diseases have higher levels of fear. In addition, although some studies (Tsiropoulou et al. 2020, Bakioğlu et al., 2020, Doshi et al. 2020) indicate that there is no difference between COVID-19 fears in terms of where people live, Hossain et al. (2020) stated in their study in Bangladesh that those living in the Capital Daka had higher levels of COVID-19 fear than those living in rural areas. However, it was an expected situation that the fear of COVID-19 would not differ in terms of demographic characteristics in this study conducted in Turkey. Because there was a complete closure in Turkey at the time of conducting this study. Places such as workplaces, universities, schools, shopping malls, restaurants and cafes, cinema and theater were all closed. In addition, because of the curfew, everyone had to stay at their houses and go out just for their basic need within certain hours. Therefore, this epidemic has adversely affected all segments and every region of the country in Turkey. Also, exaggerated and misinformation of the media, which is one of the most important sources of information in this process, negatively affected all individuals in the society.

In the present study, we found that prospective anxiety and inhibitory anxiety IUS-12 subscales, state anxiety subscale of STAI, and psychological health WHOQOL-BREF subscale were significant predictors of fear of COVID-19. These results may be interpreted as prospective concerns, sense of restraint, being restricted, state anxiety, and problems with psycholo-

gical health make people feel fear about the COVID-19 outbreak more. In a study conducted by Asmundson et al. (2020) that examined if existing anxiety and mood disorders affect COVID-19 stress responses and coping or not, it is reported that people with current anxiety or mood disorders isolate themselves more and experience more distress than people without any psychological problems. The authors have stated that fear activation is likely to remain elevated in anxiety patients. Moreover, Taylor et al. (2020) reported that people with a preexisting mental health disorder had significantly higher COVID-19 stress. In the same study, it is reported that people experiencing COVID-19 stress are more likely to be those who experience health anxiety, anxiety sensitivity, intolerance of uncertainty, disgust propensity, disgust sensitivity, perceived infectability, germ aversion, and obsessive-compulsive contamination concerns and checking rituals. Similarly, the results of another study conducted by Bakioğlu et al. (2020) in Turkey, indicated fear of COVID-19 was positively and significantly correlated with prospective anxiety, inhibitory anxiety, and stress. In the same study, it is also found that there was a mediating role of intolerance of uncertainty, depression, anxiety, and stress in the relationship between the fear of COVID-19 and positivity. Satici et al. (2020b) also have found a positive relationship between fear of COVID-19 and intolerance of uncertainty, and a negative correlation between mental health and fear of COVID-19. The results of another study conducted by Satici et al. (2020a) revealed that fear of COVID-19 was also correlated with anxiety.

Moreover, the first studies on the frequency of mental health problems seen in the population after the outbreak revealed that anxiety is the most common mental health problem (Wang et al. 2020, Li et al. 2020a, Xiao 2020). However, in a recent study by Mertens et al. (2020), it is found the risk for loved ones, information through traditional media, information through social media, and health anxiety were significant predictors for fear of COVID-19. In contrast, intolerance of uncertainty, information through professional sources, online searches, general health, worry, and information through family and friends. When the results of the present study and studies in the literature are considered together, it can be said that the pandemic created a fear against the virus in the society, and this fear has been affected by intolerance to uncertainty, anxiety, and some other problems in the psychological field. Prospective uncertainties in the areas such as work, social, education, and economy that emerged in this process caused people to experience psychological problems such as anxiety and depression, which may increase fear of the virus. Therefore, it is thought that in psychotherapy sessions to be conducted with individuals who are negatively affected by the pandemic, it will be beneficial to work with individuals' symptoms of intolerance to uncertainty and anxiety to reduce the fear of COVID-19.

## Limitations

This study has some limitations. The most important limitation was that the sample consisted of people residing in only 11 provinces of Turkey. It is important to note that this situation limits the generalizability of our results to a wider population. In addition, due to the wide age range of the sample group in our study, it could not be determined what facilitating and difficult factors were in coping with the fear of COVID-19 specific to different developmental stages. In the future, we think that conducting more comprehensive studies with children, adolescents and adults will make a significant contribution to the literature.

## CONCLUSION

Although this study has some limitations, it has an important place in the literature in revealing the facilitating and complicating factors in dealing with the fear of COVID-19. In addition, it shows that almost everyone in Turkey have negatively affected by the pandemic process, regardless of gender, region of residence, having a chronic disease or being diagnosed with COVID-19. Finally, it revealed the negative effects of the COVID-19 pandemic on mental health. At this point, policy makers and mental health workers have important responsibilities in providing psychological support services.

**Acknowledgements:** None.

**Conflict of interest:** None to declare.

### Contribution of individual authors:

Mehmet Yalçın: design of the study, literature searches and analyses, statistical analyses.

Eyüp Sabır Erbiçer: design of the study, interpretation of data, manuscript writing.

Ercan Akin: design of the study, interpretation of data, manuscript writing, analyses.

## References

1. Abuhmaidan Y & Al-Majali S: The impact of the coronavirus pandemic on mental health among al ain university students in light of some demographic variables. *Psychiatr Danub* 2020; 32:482-490
2. Ahorsu DK, Lin CY, Imani V, Saffari M, Griffiths MD, Pakpour AH: The fear of COVID-19 scale: Development and initial validation. *Int J Ment Health Addict* 2020; 1-9
3. Anjum S, Ullah R, Rana MS, Khan HA, Memon FS, Ahmed Y et al.: COVID-19 Pandemic: A serious threat for public mental health globally. *Psychiatr Danub* 2020; 32:245-250
4. Asmundson GJG, Paluszek MM, Landry CA, Rachor GS, McKay D, Taylor S: Do pre-existing anxiety-related and mood disorders differentially impact COVID-19 stress responses and coping? *J Anxiety Disord* 2020; 74:102271
5. Bakioglu F, Korkmaz O, Ercan H: Fear of COVID-19 and Positivity: Mediating role of intolerance of uncertainty, depression, anxiety, and stress. *Int J Ment Health Addict* 2020; 1-14
6. Blendon RJ, Benson JM, DesRoches CM, Raleigh E, Taylor-Clark K: The public's response to severe acute respiratory syndrome in Toronto and the United States. *Clin Infect Dis* 2004; 38:925-31
7. Bogoch II, Watts A, Thomas-Bachli A, Huber C, Kraemer MUG, Khan K: Pneumonia of unknown aetiology in Wuhan, China: potential for international spread via commercial air travel. *J Travel Med* 2020; 27:taaa008
8. Braunack-Mayer A, Tooher R, Collins JE, Street JM, Marshall H: Understanding the school community's response to school closures during the H1N1 2009 influenza pandemic. *BMC Public Health* 2013; 13:344
9. Broche-Pérez Y, Fernández-Fleites Z, Jiménez-Puig E, Fernández-Castillo E, Rodríguez-Martin BC: Gender and fear of COVID-19 in a Cuban population sample. *Int J Ment Health Addict* 2020; 12:1-9
10. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N et al.: The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 2020; 395:912-920
11. Carleton RN, Norton MA, Asmundson GJ: Fearing the unknown: a short version of the intolerance of uncertainty scale. *J Anxiety Disord* 2007; 21:105-17
12. Cheng C & Cheung MW: Psychological responses to outbreak of severe acute respiratory syndrome: a prospective, multiple time-point study. *J Pers* 2005; 73:261-85
13. Chua SE, Cheung V, McAlonan GM, Cheung C, Wong JW, Cheung EP et al.: Stress and psychological impact on SARS patients during the outbreak. *Can J Psychiatry* 2004; 49:385-90
14. Chong MY, Wang WC, Hsieh WC, Lee CY, Chiu NM, Yeh WC et al.: Psychological impact of severe acute respiratory syndrome on health workers in a tertiary hospital. *Br J Psychiatry* 2004; 185:127-33
15. Doshi D, Karunakar P, Sukhabogi JR, Prasanna JS, Mahajan SV: Assessing coronavirus fear in Indian population using the fear of COVID-19 scale. *Int J Ment Health Addict* 2020; 1-9
16. Duan L & Zhu G: Psychological interventions for people affected by the COVID-19 epidemic. *Lancet Psychiatry* 2020; 7:300-302
17. Fidaner H, Elbi H, Fidaner C, Eser, SY, Eser E, Göker, E: Measuring the quality of life. WHOQOL-100 and WHOQOL-BREF. *3P Journal* 1999; 7:5-13
18. Fischer R, Bortolini T, Karl JA, Zilberberg M, Robinson K, Rabelo A et al.: Rapid Review and Meta-Analysis of Self-Guided Interventions to Address Anxiety, Depression, and Stress During COVID-19 Social Distancing. *Front Psychol* 2020; 11:563876
19. Graves C: *Invasion by virus: Can it happen again?*. Icon Books, London, 1969
20. Group WHOQOL: Development of the WHOQOL: Rationale and current status. *Int J Ment Health* 1994; 23:24-56
21. Hossain MA, Jahid MIK, Hossain KMA, Walton LM, Uddin Z, Haque MO et al.: Knowledge, attitudes, and fear of COVID-19 during the rapid rise period in Bangladesh. *PLoS One* 2020; 15:e0239646

22. Hui DSC & Zumla A: Severe acute respiratory syndrome: Historical, epidemiologic, and clinical features. *Infect Dis Clin North Am* 2019; 33:869-889
23. Huremović D: Mental health of quarantine and isolation, in Huremović D (eds.): *Psychiatry of pandemics, A mental health response to infection outbreak*. Springer, Switzerland, 2019
24. Huynh J, Li S, Yount B, Smith A, Sturges L, Olsen JC et al.: Evidence supporting a zoonotic origin of human coronavirus strain NL63. *J Virol* 2012; 86:12816-25
25. Karasar B & Canli D: Psychological resilience and depression during the Covid-19 pandemic in Turkey. *Psychiatr Danub* 2020; 32:273-279
26. Kang L, Li Y, Hu S, Chen M, Yang C, Yang BX: The mental health of medical workers in Wuhan, China dealing with the 2019 novel coronavirus. *Lancet Psychiatry* 2020; 7:e14
27. Krause AE, Dimmock J, Rebar AL, Jackson B: Music Listening Predicted Improved Life Satisfaction in University Students During Early Stages of the COVID-19 Pandemic. *Front Psychol* 2021; 11:631033
28. Leigh-Hunt N, Bagguley D, Bash K, Turner V, Turnbull S, Valtorta N et al.: An overview of systematic reviews on the public health consequences of social isolation and loneliness. *Public Health* 2017; 152:157-171.
29. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y et al.: Early transmission dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N Engl J Med* 2020a; 382:1199-1207
30. Li J, Yang Z, Qiu H, Wang Y, Jian L, Ji J et al.: Anxiety and depression among general population in China at the peak of the COVID-19 epidemic. *World Psychiatry* 2020b; 19:249-250.
31. Lu J, Gu J, Li K, Xu C, Su W, Lai Z et al.: COVID-19 Outbreak associated with air conditioning in restaurant, guangzhou, China, 2020. *Emerg Infect Dis* 2020; 26:1628-1631
32. Mak IW, Chu CM, Pan PC, Yiu MG, Chan VL: Long-term psychiatric morbidities among SARS survivors. *Gen Hosp Psychiatry* 2009; 31:318-26
33. McEvoy PM & Mahoney AE: Intolerance of uncertainty and negative metacognitive beliefs as transdiagnostic mediators of repetitive negative thinking in a clinical sample with anxiety disorders. *J Anxiety Disord* 2013; 27:216-24
34. Mertens G, Gerritsen L, Duijndam S, Saleminck E, Engelhard IM: Fear of the coronavirus (COVID-19): Predictors in an online study conducted in March 2020. *J Anxiety Disord* 2020; 74:102258. doi:10.1016/j.janxdis.2020.102258
35. Metin A, Çetinkaya A, Erbiçer ES: Subjective well-being and resilience during covid-19 pandemic in Turkey. *European Journal of Health Psychology* 2021, In press
36. Öner N & LeCompte WA: *The state-trait anxiety inventory handbook*. Boğaziçi University Press, Istanbul, 1985
37. Pappas G, Kiriakou IJ, Giannakis P, Falagas ME: Psychosocial consequences of infectious diseases. *Clin Microbiol Infect* 2009; 15:743-7
38. Ping W, Zheng J, Niu X, Guo C, Zhang J, Yang H et al.: Evaluation of health-related quality of life using EQ-5D in China during the COVID-19 pandemic. *PLoS One* 2020; 15:e0234850
39. Puyat JH, Ahmad H, Avina-Galindo AM, Kazanjian A, Gupta A, Ellis U et al.: A rapid review of home-based activities that can promote mental wellness during the COVID-19 pandemic. *PLoS One* 2020; 15:e0243125
40. Ren FF & Guo RJ: Public mental health in Post-COVID-19 era. *Psychiatr Danub* 2020; 32:251-255
41. Reznik A, Gritsenko V, Konstantinov V, Khamenka N, Isralowitz R: COVID-19 fear in Eastern Europe: validation of the fear of COVID-19 scale. *Int J Ment Health Addict* 2020; 12:1-6
42. Sakib N, Bhuiyan AKMI, Hossain S, Al Mamun F, Hosen I, Abdullah AH et al.: Psychometric validation of the Bangla fear of COVID-19 scale: Confirmatory factor analysis and rasch analysis. *Int J Ment Health Addict* 2020; 1-12
43. Saricam H, Erguvan FM, Akin A, Akça MS: The Turkish short version of the intolerance of uncertainty (IUS-12) scale: The study of validity and reliability. *Route educational and social science journal* 2014; 1: 148-157
44. Satici B, Gocet-Tekin E, Deniz ME, Satici SA: Adaptation of the Fear of COVID-19 Scale: Its association with psychological distress and life satisfaction in Turkey. *Int J Ment Health Addict* 2020a; 8:1-9
45. Satici B, Saricali M, Satici SA, Griffiths MD: Intolerance of uncertainty and mental wellbeing: Serial mediation by rumination and fear of COVID-19. *Int J Ment Health Addict* 2020b; 15:1-12
46. Shigemura J, Ursano RJ, Morganstein JC, Kurosawa M, Benedek DM: Public responses to the novel 2019 coronavirus (2019-nCoV) in Japan: Mental health consequences and target populations. *Psychiatry Clin Neurosci* 2020; 74:281-282
47. Sibley CG, Greaves LM, Satherley N, Wilson MS, Overall NC, Lee CHJ et al.: Effects of the COVID-19 pandemic and nationwide lockdown on trust, attitudes toward government, and well-being. *Am Psychol* 2020; 75:618-630
48. Šljivo A, Kačamaković M, Quraishi I, Džibur Kulenović A: Fear and Depression among Residents of Bosnia and Herzegovina during COVID-19 Outbreak - Internet Survey. *Psychiatr Danub* 2020; 32:266-272
49. Sønderkov KM, Dinesen PT, Santini ZI, Østergaard SD: The depressive state of Denmark during the COVID-19 pandemic. *Acta Neuropsychiatr* 2020; 32:226-228
50. Soraci P, Ferrari A, Abbiati FA, Del Fante E, De Pace R, Urso A et al.: Validation and psychometric evaluation of the Italian version of the fear of COVID-19 scale. *Int J Ment Health Addict* 2020; 4:1-10
51. Spielberger CD: *Understanding stress and anxiety*. Harper & Row, New York, 1979
52. Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA: *Manual for the State-Trait Anxiety Inventory*. Consulting psychologists press, USA, 1983
53. Taha S, Matheson K, Cronin T, Anisman H: Intolerance of uncertainty, appraisals, coping, and anxiety: the case of the 2009 H1N1 pandemic. *Br J Health Psychol* 2014; 19:592-605
54. Taylor S: *The psychology of pandemics: Preparing for the next global outbreak of infectious disease*. Cambridge Scholars Publishing, Newcastle upon Tyne, 2019
55. Taylor S, Landry CA, Paluszek MM, Fergus TA, McKay D, Asmundson GJG: COVID stress syndrome: Concept, structure, and correlates. *Depress Anxiety* 2020; 37:706-714

56. Tsipropoulou V, Nikopoulou VA, Holeva V, Nasika Z, Diakogiannis I, Sakka S et al.: Psychometric properties of the Greek version of FCV-19S. *Int J Ment Health Addict* 2020; 1-10
57. Tzur Bitan D, Grossman-Giron A, Bloch Y, Mayer Y, Shiffman N, Mendlovic S: Fear of COVID-19 scale: Psychometric characteristics, reliability and validity in the Israeli population. *Psychiatry Res* 2020; 289:113100
58. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS et al.: Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *Int J Environ Res Public Health* 2020; 17:1729
59. Weiss SR & Leibowitz JL: Coronavirus pathogenesis. *Adv Virus Res* 2011; 81:85-164
60. WHO: Summary of probable SARS cases with onset of illness from 1 November to 31 July 2003. [http://www.who.int/csr/sars/country/table2004\\_04\\_21/en/](http://www.who.int/csr/sars/country/table2004_04_21/en/)
61. WHO: Coronavirus disease (COVID-19) pandemic. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>. WHO, 2020
62. Williams SN, Armitage CJ, Tampe T, Dienes K: Public perceptions and experiences of social distancing and social isolation during the COVID-19 pandemic: a UK-based focus group study. *BMJ Open* 2020; 10:e039334
63. Woo PC, Lau SK, Lam CS, Lau CC, Tsang AK, Lau JH et al.: Discovery of seven novel Mammalian and avian coronaviruses in the genus deltacoronavirus supports bat coronaviruses as the gene source of alphacoronavirus and betacoronavirus and avian coronaviruses as the gene source of gammacoronavirus and deltacoronavirus. *J Virol* 2012; 86:3995-4008
64. Xiao C: A novel approach of consultation on 2019 novel coronavirus (COVID-19)-related psychological and mental problems: structured letter therapy. *Psychiatry Investig* 2020; 17:175-176

Correspondence:

Eyüp Sabır Erbiçer, MA

Department of Psychological Counseling and Guidance, Hacettepe University

Ankara, Turkey

E-mail: eyuperbicer@gmail.com