



Cross-Sectional Gene-Smoking Interaction Analysis in Relation to Subclinical Atherosclerosis—Results From the IMPROVE Study

Buamina Maitusong, MD, PhD*¹; Federica Laguzzi¹, Pharma D, PhD*¹; Rona J. Strawbridge¹, PhD; Damiano Baldassarre¹, PhD; Fabrizio Veglia¹, PhD; Steve E. Humphries¹, PhD; Kai Savonen¹, MD, PhD; Sudhir Kurl, MD, PhD; Matteo Pirro¹, MD, PhD; Andries J. Smit, MD, PhD; Philippe Giral¹, MD, PhD; Angela Silveira¹, PhD; Elena Tremoli¹, PhD; Anders Hamsten, MD, PhD; Ulf de Faire¹, MD, PhD; Bruna Gigante¹, MD, PhD; Karin Leander¹, PhD; on behalf of the IMPROVE Study group†

BACKGROUND: Smoking is associated with carotid intima-media thickness (C-IMT). However, knowledge about how genetics may influence this association is limited. We aimed to perform nonhypothesis driven gene-smoking interaction analyses to identify potential genetic variants, among those included in immune and metabolic platforms, that may modify the effect of smoking on carotid intima-media thickness.

METHODS: We used baseline data from 1551 men and 1700 women, aged 55 to 79, included in a European multi-center study. Carotid intima-media thickness maximum, the maximum of values measured at different locations of the carotid tree, was dichotomized with cut point values ≥ 75 , respectively. Genetic data were retrieved through use of the Illumina Cardio-Metabo- and Immuno- Chips. Gene-smoking interactions were evaluated through calculations of Synergy index (S). After adjustments for multiple testing, P values of $< 2.4 \times 10^{-7}$ for S were considered significant. The models were adjusted for age, sex, education, physical activity, type of diet, and population stratification.

RESULTS: Our screening of 207 586 SNPs available for analysis, resulted in the identification of 47 significant gene-smoking synergistic interactions in relation to carotid intima-media thickness maximum. Among the significant SNPs, 28 were in protein coding genes, 2 in noncoding RNA and the remaining 17 in intergenic regions.

CONCLUSIONS: Through nonhypothesis-driven analyses of gene-smoking interactions, several significant results were observed. These may stimulate further research on the role of specific genes in the process that determines the effect of smoking habits on the development of carotid atherosclerosis.

Key Words: carotid intima-media thickness ■ epidemiologic studies ■ gene-environment interaction ■ polymorphism, single nucleotide ■ smoking

Subclinical atherosclerosis is an asymptomatic, chronic condition that is easily undiagnosed until a clinical event occurs, such as myocardial infarction or stroke.¹ Carotid intima-media thickness (C-IMT), assessed with B-mode ultrasound, a noninvasive method, has been shown to be a valid surrogate marker

for subclinical atherosclerosis,² and a predictor for future cardiovascular disease (CVD).^{3,4}

Previous studies indicate that genetic susceptibility plays an important role in the pathogenesis of atherosclerosis.^{5–8} The reported proportions of heritability of carotid atherosclerosis vary between 2% and 78%.⁸ Part of this

Correspondence to: Federica Laguzzi, Pharma D, PhD, Unit of Cardiovascular and Nutritional Epidemiology, Institute of Environmental Medicine, Karolinska Institutet, Nobels väg 13, Box 210, 17177 Stockholm, Sweden. Email federica.laguzzi@ki.se

*B. Maitusong and F. Laguzzi contributed equally.

†A list of all IMPROVE study group members is given in the Supplemental Material.

Supplemental Material is available at <https://www.ahajournals.org/doi/suppl/10.1161/CIRCGEN.122.003710>.

For Sources of Funding and Disclosures, see page 245.

© 2023 The Authors. *Circulation: Genomic and Precision Medicine* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer Health, Inc. This is an open access article under the terms of the [Creative Commons Attribution Non-Commercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use, distribution, and reproduction in any medium, provided that the original work is properly cited, the use is noncommercial, and no modifications or adaptations are made.

Circulation: Genomic and Precision Medicine is available at www.ahajournals.org/journal/circgen

Nonstandard Abbreviations and Acronyms

APOB	apolipoprotein B
C-IMT	carotid intima-media thickness
C-IMT_{max}	maximum of C-IMT values measured at different locations of the carotid tree
GWAS	genome-wide association study
IMPROVE	Carotid Intima Media Thickness [IMT] and IMT-Progression as Predictors of Vascular Events in a High Risk European Population
SNP	single nucleotide polymorphism

heritability is likely to be explained by gene-environment interactions.⁸ There are hopes from the scientific community and healthcare that personalized medicine, such as knowledge of how genetic background can interact with modifiable factors and thereby influence cardiovascular risk, will be able to contribute to improved prevention of CVD.

Among the risk factors for premature atherosclerosis, smoking has been identified as a major determinant of atherosclerotic development.^{9–11} Studies have shown that smoking exposure and duration of smoking cessation can affect carotid artery structure in all phases of atherosclerosis.^{12,13} In an earlier investigation based on data from a European multi-center study IMPROVE (Carotid Intima Media Thickness [IMT] and IMT-Progression as Predictors of Vascular Events in a High Risk European Population), smoking was found to be a major determinant of C-IMT.¹⁴

Previous studies that have investigated gene-smoking interactions behind carotid atherosclerosis were generally performed with a candidate gene approach, and the results are inconclusive.^{15–33} Only 2 studies evaluated gene-smoking interactions with an explorative approach, using the whole genome, one based on 669 Hispanics, mainly women, residing in New York,³⁴ and the other based on 1776 men from West Africa.³⁵ These studies are insufficient to detect all important gene-smoking interactions due to their limited sample size. In addition, it is doubtful whether the results can be generalized to populations of other ancestries.

Hence, we aimed to explore gene-smoking interactions behind carotid subclinical atherosclerosis in a multi-center study including men and women of European ancestry. We limited the search for interactions to include genetic variants available via platforms for genetic studies of cardiovascular, metabolic, and immune traits.

METHODS

The data that support the findings of this study are available from the corresponding author upon reasonable request.

The Institutional review board at each recruitment center (Karolinska Institutet, Stockholm, Sweden; University of Milan, Milan, Italy; University of Kuopio and Kuopio Research Institute of Exercise Medicine, Kuopio, Finland; University Hospital Groningen, Groningen, The Netherlands; University of Perugia, Perugia, Italy; Groupe Hôpital Pitie-Salpetriere, Paris, France) approved the study. Written informed consents for general participation and for the genotyping were provided by all participants. The study was performed in accordance with the Helsinki Declaration.

Full materials and methods are available in [Supplemental Materials](#).

RESULTS

Baseline characteristics of all study participants and by their smoking status are presented in Table 1. The current smokers were younger, less physically active, and educated than nonsmokers. Smokers had also higher levels of total cholesterol, triglycerides, LDL-C (low-density lipoprotein cholesterol), blood glucose, and High-sensitivity C-reactive protein. However, their level of uric acid and creatinine were lower than in nonsmokers.

In total, 207 586 genetic variants were available for analyses. Results from the main analysis investigating gene-smoking interaction in relation to C-IMT_{max} cut off at the 75th percentile are shown in Table 2. We found 47 single nucleotide polymorphisms (SNPs) significant (P for Synergy index $<2.4 \times 10^{-7}$) after Bonferroni correction. All the aforementioned interaction results were synergistic, with Synergy index point estimates in the range between 3.3 and 5.8 (Table S1). Compared with the reference group of nonsmokers without the risk variant, the odds for having C-IMT >75 th percentile associated with smoking and having the risk variant were ≈ 3 to 4-fold higher (Table 2). Of the 47 significant SNPs, 28 were in protein coding genes, 2 in noncoding RNA and the remaining 17 in intergenic regions (Table 3). None of the 47 SNPs involved in the interactions identified in our study were among the published quantitative trait locus data included in the Genotype-Tissue Expression (accessed March 25, 2022).

Additional analysis that used C-IMT_{max} cutoff at the 50th percentile resulted in the identification of 146 SNPs for which a significant synergistic interaction with smoking was observed (Table S2). Among those SNPs, 75 were in protein coding genes, 21 in noncoding RNA, and the remaining 50 in intergenic regions (Table S3). Two of these significant SNPs (rs6032180 in *LOC105372631* and rs3744761 in *PLCD3*) were found both when using the 75th and the 50th percentile C-IMT_{max} cutoff values.

Analyses of gene-smoking interactions that also considered data where the number of observations for each of the possible combinations of the exposures considered are <10 resulted in the identification of additional significant results for the C-IMT_{max} cutoff 75th percentile (Table S4), and for C-IMT_{max} cutoff 50th percentile

Table 1. Characteristics of the IMPROVE Study Participants by Smoking Status

	Entire sample (N=3251)	Current smokers (n=520)	Nonsmokers (n=2731)
Male; n (%)	1551 (47.7)	273 (52.5)	1278 (46.8)
Age, y, mean±SD	64.4±5.4	61.5±5.2	64.9±5.7
Geographic gradient, n (%)			
Kuopio	928 (28.5)	166 (31.9)	762 (27.9)
Stockholm	488 (15.0)	63 (12.1)	425 (15.6)
Groningen	400 (12.3)	77 (14.8)	323 (11.8)
Paris	434 (13.3)	48 (9.2)	386 (14.1)
Milan	517 (15.9)	93 (17.9)	424 (15.5)
Perugia	484 (14.9)	73 (14.0)	411 (15.0)
Anthropometric variables, mean±SD			
BMI, kg/m ²	26.7±4.3	26.3±4.2	26.7±4.6
Waist/hip ratio	0.92±0.09	0.92±0.08	0.92±0.09
Blood pressure, mean±SD			
Diastolic blood pressure, mm Hg	81±9	81±10	81±10
Systolic blood pressure, mm Hg	140±19	140±18	140±18
Physical activity level, n (%)			
Low	613 (18.8)	114 (22.0)	499 (18.3)
Medium	1453 (44.8)	234 (45.1)	1219 (44.7)
High	1180 (36.4)	171 (32.9)	1009 (37.0)
Education, n (%)			
≤9 y	1467 (45.6)	238 (46.6)	1229 (45.4)
9–12 y	803 (25.0)	133 (26.0)	670 (24.7)
>12 y	944 (29.3)	140 (27.4)	804 (29.7)
Mediterranean diet score,* n (%)			
0	316 (9.7)	68 (13.1)	248 (9.1)
1	758 (23.3)	142 (27.3)	616 (22.6)
2	930 (28.6)	159 (30.6)	771 (28.2)
3	731 (22.5)	102 (19.6)	629 (23.0)
4	427 (13.1)	42 (8.1)	385 (14.1)
5	80 (2.5)	7 (1.3)	73 (2.7)
6	9 (0.3)	0 (0.0)	9 (0.3)
Biochemical markers, mean±SD			
Total cholesterol, mmol/L	5.44±1.11	5.53±1.12	5.43±1.25
HDL cholesterol, mmol/L	1.21±0.36	1.15±0.36	1.22±0.30
Triglycerides, mmol/L	1.29±1.01	1.41±1.13	1.27±1.17
LDL cholesterol, mmol/L	3.51±1.01	3.57±1.01	3.49±0.94
Uric acid, μmol/L	313.8±71.6	309.0±72.0	314.7±71.5
Blood glucose, mmol/L	5.50±1.54	5.60±1.45	5.50±1.42
Creatinine, μmol/L	80.7±17.7	80.1±17.3	80.8±17.7
C-reactive protein	2.79±4.30	2.98±3.60	2.76±4.42
Medical history and drug use, n (%)			
Hypercholesterolemia†	2299 (70.8)	337 (64.9)	1962 (71.9)
Hypertriglyceridemia‡	827 (25.5)	151 (29.0)	676 (24.8)
Hypertension§	2327 (71.6)	331 (63.6)	1996 (73.1)
Diabetes	775 (24.2)	125 (24.3)	650 (24.2)
Statin use	1290 (39.7)	166 (31.9)	1124 (41.2)

Results are expressed as mean and SD for continuous variables and as count and proportion (%) for categorical variables. HDL indicates high-density lipoprotein; IMPROVE, Carotid Intima Media Thickness [IMT] and IMT-Progression as Predictors of Vascular Events in a High Risk European Population; and LDL, low-density lipoprotein.

*The score indicates level of adherence; zero corresponds to the lowest level.

†Serum total cholesterol >5.17 mmol/L.

‡Serum triglycerides >1.7 mmol/L.

§Self-reported and use of antihypertensive drugs.

||Self-reported and/or use of antidiabetic drugs.

Table 2. Significant Gene-Smoking Interaction Results* After Bonferroni Adjustment for Multiple Testing in Relation to C-IMT_{max} With Cutoff at the 75th Percentile

	Number of observations								Risk allele	MAF (%)	Odds ratio (95% CI)†			P Synergy index
	Nonsmokers without the risk variant		Nonsmokers with the risk variant		Smokers without the risk variant		Smokers with the risk variant				Reference group: nonsmokers without the risk variant			
	Controls	Cases	Controls	Cases	Controls	Cases	Controls	Cases			Nonsmokers with the risk variant	Smokers without the risk variant	Smokers with the risk variant	
Chr 1														
rs12134420	1874	597	65	22	324	144	11	13	C	20	1.11 (0.67–1.83)	1.62 (1.29–2.03)	4.92 (2.09–11.57)	4.43×10 ⁻¹⁴
rs2446622	1702	538	230	75	306	131	26	25	G	6	1.02 (0.77–1.36)	1.56 (1.23–1.98)	3.86 (2.14–6.97)	9.17×10 ⁻¹⁸
rs72676073	1683	532	256	87	290	119	45	38	G	7	1.19 (0.91–1.55)	1.50 (1.17–1.92)	3.53 (2.22–5.62)	8.51×10 ⁻¹⁵
rs73009101	1742	545	197	74	315	137	20	20	G	5	1.18 (0.88–1.58)	1.60 (1.27–2.02)	4.03 (2.12–7.68)	5.96×10 ⁻¹³
Chr 2														
rs6758414	1746	558	191	61	309	134	26	23	A	5	1.00 (0.73–1.37)	1.56 (1.23–1.97)	3.60 (2.00–6.50)	6.51×10 ⁻¹⁵
rs9789490	1386	443	553	176	253	93	82	64	G	16	1.01 (0.82–1.24)	1.33 (1.01–1.74)	2.96 (2.06–4.24)	4.34×10 ⁻¹⁷
Chr 3														
rs9877192	1801	572	138	47	317	139	18	18	A	26	1.12 (0.79–1.6)	1.6 (1.27–2.02)	4.21 (2.07–8.54)	1.48×10 ⁻¹³
Chr 4														
rs11736632	1620	503	319	116	281	110	54	47	A	9	1.13 (0.89–1.44)	1.48 (1.15–1.9)	3.13 (2.05–4.78)	9.51×10 ⁻¹³
Chr 5														
rs13176964	1587	490	352	128	285	116	50	41	G	10	1.16 (0.92–1.46)	1.52 (1.18–1.94)	3.24 (2.08–5.05)	9.03×10 ⁻¹³
rs2278392	1553	495	386	124	270	108	65	49	T	11	1.01 (0.80–1.28)	1.44 (1.12–1.87)	2.86 (1.92–4.27)	7.74×10 ⁻¹³
rs4867490	1622	517	316	102	291	121	44	36	G	40	1.06 (0.82–1.36)	1.50 (1.17–1.91)	3.27 (2.05–5.21)	8.27×10 ⁻¹⁵
rs7722352	1684	533	252	86	306	132	29	25	G	7	1.04 (0.79–1.36)	1.56 (1.23–1.98)	3.45 (1.95–6.10)	3.08×10 ⁻¹³
Chr 7														
rs28695838	1580	504	353	115	290	116	44	41	G	10	1.03 (0.81–1.31)	1.43 (1.12–1.84)	3.62 (2.31–5.69)	2.21×10 ⁻²³
Chr 8														
rs12545167	1302	412	637	207	246	97	89	60	A	18	1.03 (0.84–1.25)	1.37 (1.05–1.80)	2.81 (1.96–4.03)	8.11×10 ⁻¹⁴
rs4301463	1678	530	261	89	304	127	31	30	A	7	1.14 (0.87–1.49)	1.54 (1.21–1.96)	3.77 (2.2–6.44)	9.53×10 ⁻¹⁶
rs6997802	1679	530	260	89	304	127	31	30	T	7	1.14 (0.88–1.49)	1.54 (1.21–1.96)	3.77 (2.20–6.44)	1.13×10 ⁻¹⁵
rs752039	1266	404	673	215	242	93	93	64	A	19	1.01 (0.83–1.23)	1.33 (1.01–1.76)	2.82 (1.98–4.02)	5.92×10 ⁻¹⁵
Chr 9														
rs10810371	1414	430	509	185	258	97	72	58	G	15	1.17 0.95– 1.44)	1.43 (1.10–1.88)	3.02 (2.07–4.40)	4.91×10 ⁻¹³
rs143207461	1734	549	205	70	314	134	21	23	C	5	1.12 (0.83–1.50)	1.57 (1.24–1.99)	4.00 (2.16–7.43)	2.16×10 ⁻¹⁵
Chr 10														
rs12244483	1708	540	230	79	297	120	38	37	T	6	1.05 (0.8–1.40)	1.51 (1.18–1.92)	3.27 (2.02–5.29)	1.20×10 ⁻¹³

(Continued)

Downloaded from <http://ahajournals.org> by on June 28, 2023

Table 2. Continued

	Number of observations								Risk allele	MAF (%)	Odds ratio (95% CI) [†]			P Synergy index
	Nonsmokers without the risk variant		Nonsmokers with the risk variant		Smokers without the risk variant		Smokers with the risk variant				Reference group: nonsmokers without the risk variant			
	Con-trols	Cases	Con-trols	Cases	Con-trols	Cases	Con-trols	Cases			Nonsmok-ers with the risk variant	Smokers without the risk variant	Smokers with the risk variant	
rs12251673	1721	545	217	74	304	127	31	30	C	6	1.01 (0.75–1.35)	1.54 (1.21–1.95)	3.24 (1.91–5.49)	1.16×10 ⁻¹²
rs7068194	1722	545	217	74	304	127	31	30	T	6	1.01 (0.75–1.35)	1.54 (1.21–1.96)	3.25 (1.92–5.50)	1.17×10 ⁻¹²
rs7092757	1708	541	231	77	296	120	39	37	G	6	1.01 (0.76–1.35)	1.51 (1.18–1.92)	3.19 (1.98–5.13)	1.60×10 ⁻¹³
rs72826094	1847	589	91	28	321	145	12	12	A	20	1.02 (0.65–1.59)	1.63 (1.30–2.05)	4.55 (1.92–10.81)	4.27×10 ⁻¹⁴
Chr 11														
rs1002171	1716	535	223	84	306	128	29	29	G	6	1.10 (0.84–1.46)	1.55 (1.22–1.98)	3.46 (2.02–5.94)	1.17×10 ⁻¹²
rs2434468	1654	515	285	103	307	129	28	27	C	8	1.06 (0.82–1.37)	1.55 (1.22–1.97)	3.55 (2.04–6.18)	6.56×10 ⁻¹⁵
rs2511241	1658	526	281	93	301	124	34	33	C	8	1.06 (0.82–1.38)	1.51 (1.19–1.93)	3.69 (2.21–6.16)	1.53×10 ⁻¹⁸
rs3741392	1630	520	309	99	295	117	39	40	C	8	1.01 (0.78–1.30)	1.49 (1.17–1.91)	3.16 (1.97–5.05)	3.20×10 ⁻¹⁴
rs61899280	1726	546	213	73	310	134	25	23	C	6	1.07 (0.80–1.43)	1.57 (1.24–1.98)	3.96 (2.18–7.19)	3.63×10 ⁻¹⁷
Chr 12														
rs10506726	1733	537	206	81	317	135	17	22	T	6	1.22 (0.92–1.62)	1.59 (1.26–2.01)	5.04 (2.57–9.86)	1.95×10 ⁻²¹
rs11171745	1494	470	445	149	266	106	69	51	A	12	1.08 (0.87–1.34)	1.45 (1.12–1.88)	2.91 (1.97–4.30)	1.14×10 ⁻¹²
rs11171773	1691	536	248	83	302	128	33	29	A	7	1.06 (0.81–1.40)	1.54 (1.21–1.96)	3.33 (1.98–5.61)	9.35×10 ⁻¹³
rs116378618	1718	545	219	74	307	131	28	26	A	6	1.09 (0.81–1.45)	1.56 (1.23–1.98)	3.50 (2.01–6.11)	8.02×10 ⁻¹³
rs1689512	1494	470	445	149	266	106	69	51	G	12	1.08 (0.87–1.34)	1.45 (1.12–1.88)	2.91 (1.97–4.30)	1.14×10 ⁻¹²
rs17118317	1478	469	461	150	265	106	70	51	C	13	1.03 (0.83–1.28)	1.44 (1.11–1.86)	2.85 (1.93–4.21)	8.38×10 ⁻¹³
rs35436573	1604	507	335	112	288	119	47	38	A	9	1.03 (0.81–1.32)	1.49 (1.16–1.91)	3.13 (1.99–4.92)	3.83×10 ⁻¹⁴
rs4762693	1788	571	151	48	319	141	16	16	G	27	1.03 (0.73–1.45)	1.61 (1.27–2.02)	3.90 (1.89–8.05)	7.19×10 ⁻¹³
rs773643	1623	513	316	106	285	117	50	40	A	9	1.09 (0.85–1.39)	1.49 (1.16–1.91)	3.15 (2.03–4.91)	2.49×10 ⁻¹³
rs7956913	1488	473	448	146	266	106	69	51	D	12	1.03 (0.83–1.29)	1.43 (1.10–1.85)	2.91 (1.97–4.30)	8.69×10 ⁻¹⁴
Chr 13														
rs12872592	1367	430	572	189	261	99	74	58	G	15	1.04 (0.85–1.28)	1.39 (1.07–1.81)	3.00 (2.05–4.38)	5.58×10 ⁻¹⁶
Chr 14														
rs4981312	1410	452	526	167	254	99	80	58	G	15	1.01 (0.82–1.24)	1.39 (1.06–1.81)	2.80 (1.94–4.04)	1.15×10 ⁻¹³
rs7155978	1658	523	281	96	301	128	34	29	T	7	1.03 (0.80–1.34)	1.54 (1.21–1.96)	3.30 (1.95–5.59)	2.69×10 ⁻¹³
rs915064	1746	553	193	66	311	135	24	22	C	5	1.05 (0.77–1.42)	1.58 (1.25–2.00)	3.59 (1.96–6.59)	4.86×10 ⁻¹³

(Continued)

Downloaded from <http://ahajournals.org> by on June 28, 2023

Table 2. Continued

	Number of observations								Risk allele	MAF (%)	Odds ratio (95% CI)†				P Synergy index
	Nonsmokers without the risk variant		Nonsmokers with the risk variant		Smokers without the risk variant		Smokers with the risk variant				Reference group: nonsmokers without the risk variant				
	Con-trols	Cases	Con-trols	Cases	Con-trols	Cases	Con-trols	Cases			Nonsmok-ers with the risk variant	Smok-ers without the risk variant	Smok-ers with the risk variant		
Chr 16															
rs1003341	1140	356	799	263	220	79	115	78	T	23	1.05 (0.87–1.27)	1.29 (0.96–1.73)	2.69 (1.94–3.73)	6.22×10 ⁻¹³	
Chr 17															
rs3744761	1762	559	177	60	311	131	24	26	T	5	1.05 (0.76–1.44)	1.57 (1.24–1.98)	3.52 (1.96–6.32)	1.15×10 ⁻¹²	
rs4362432	1660	516	279	103	300	121	35	36	A	8	1.17 (0.90–1.50)	1.53 (1.20–1.95)	3.52 (2.15–5.76)	4.68×10 ⁻¹⁴	
Chr 20															
rs6032180	1607	508	332	111	285	117	50	40	T	9	1.06 (0.83–1.35)	1.48 (1.16–1.90)	3.23 (2.06–5.07)	4.64×10 ⁻¹⁵	

A dominant genetic model was assumed. #C-IMT_{max} indicates maximum of carotid intima media thickness values measured at different locations of the carotid tree; and MAF, minor allele frequency.

*Synergy index results were considered significant at $P < 2.4 \times 10^{-7}$; minimum number of subjects in each group: 10.

†Model adjusted for sex, age, education (categorical), physical activity (categorical), Mediterranean diet score, and population structure (multidimensional scaling-3 continuous).

#Individuals who carry either 1 or 2 copies of the risk allele are considered to carry the risk variant.

(Table S5). All the observed interactions were synergistic. Of the SNPs that appeared in these results, 130 are located in protein coding genes, 43 in long noncoding RNA, and 84 in intergenic regions (Table S6).

We observed no significant results of interaction on the multiplicative scale.

DISCUSSION

In this population of European descent at high risk of CVD but free of clinical manifestations of CVD, our non-hypothesis-based analyses of gene-smoking interactions resulted in the identification of several genetic variants that may have a role in the process behind the effects of smoking on the development of carotid atherosclerosis. Among the 47 SNPs identified in the main analyses, 8 SNPs (Figure 1) are located in any of 7 coding genes that in previous research have been linked to atherosclerosis development: rs72676073 in the interleukin 23 receptor (*IL23R*), rs9877192 in the LIM (Lin-11, Islet-1, and Mec-3) domain containing preferred translocation partner in lipoma (*LPP*), rs2278392 in the 5-hydroxytryptamine receptor 4 (*HTR4*), rs10810371 in the tetratricopeptide repeat domain 39B (*TTC39B*), rs7068194 and rs12251673 in the 6-phosphofructo-2-kinase/fructose-2,6-biphosphatase (*PFKFB3*), rs2511241 in the purinergic receptor P2Y2 (*P2RY2*), and rs915064 in the potassium voltage-gated channel subfamily H member 5 (*KCNH5*).^{36–42} None of these coding genes were identified in 2 previous studies that evaluated gene-smoking interactions with an explorative approach in relation to

carotid atherosclerosis.^{34,35} These 2 studies were based on the whole genome and assessed interaction on the multiplicative scale only; significant findings of interaction with smoking were observed for a few genetic variants (rs112017404; rs144170770; rs4941649; rs1192824; rs77461169; rs3751383)^{34,35} that were not available in the Cardio-Metabo- and Immuno-Chips.

Scientific support for relevance of the *IL23R* gene seems to be emerging; it encodes for a protein, interleukin 23 receptor, involved in the cascade of proinflammatory mediators which may in turn play a role in the development of atherosclerosis.³⁶ Further, the *IL23R* gene has been previously related to autoimmune disease^{43,44} and smoking behavior.⁴⁵ It has been found to synergistically interact with smoking in relation to sarcoidosis, an autoimmune disease, in a Swedish population-based case-control study.⁴³ The *HTR4* and *P2RY2* genes may also possibly be of particular interest. These proteins belong to the family of serotonin and purinergic receptors, respectively. The activation of extracellular nucleotide purinergic receptors, such as ATP, has been suggested to stimulate inflammatory mediators⁴⁶ and regulate the expression of vascular cell adhesion molecule, which is thought to be important for the pathogenesis of atherosclerosis.³⁹ The *HTR4* gene has been noted to associate to C-IMT in a previous study based on the IMPROVE study material using a candidate gene approach.⁴¹

Among the 47 SNPs identified in our main analysis of interaction as well as in our additional analyses that used the 50th percentile cutoff, there is a SNP (rs3744761), located in a protein coding gene, the phospholipase C

Table 3. Genes in Proximity to the Genetic Variants Included in the Significant Gene-Smoking Interaction Results Observed for C-IMT_{max} With Cutoff at the 75th Percentile

	Position	Function	Gene in proximity to the genetic variant
Chr 1			
rs12134420	85272625	Intron variant	<i>BCL10</i>
rs2446622	161637183	Intergenic variant	<i>None</i>
rs72676073	67203930	Intron variant	<i>IL23R</i>
rs73009101	116825975	Intergenic variant	<i>None</i>
Chr 2			
rs6758414	120538909	Intergenic variant	
rs9789490	212992755	Upstream variant	<i>LOC102725082</i>
Chr 3			
rs9877192	188708468	Intron variant	<i>LPP</i>
Chr 4			
rs11736632	56306169	Intron variant	<i>CRACD; LOC105377664</i>
Chr 5			
rs13176964	175407619	Intergenic variant	<i>None</i>
rs2278392	148548662	Intron variant; upstream variant	<i>HTR4; LOC107986462; LOC105378221</i>
rs4867490	32919896	Intergenic variant	<i>None</i>
rs7722352	123450395	Intergenic variant	<i>None</i>
Chr 7			
rs28695838	52527273	Intergenic variant	<i>None</i>
Chr 8			
rs12545167	69595708	Intron variant	<i>SULF1</i>
rs752039	69601242	Intron variant	<i>SULF1</i>
rs4301463	130457363	Intergenic variant	<i>None</i>
rs6997802	130457843	Intergenic variant	<i>None</i>
Chr 9			
rs10810371	15290344	Intron variant	<i>TTC39B</i>
rs143207461	133514431	Upstream variant	<i>MYMK</i>
Chr 10			
rs12244483	30545968	Intergenic variant	<i>None</i>
rs12251673	6150108	Intron variant	<i>PFKFB3</i>
rs7068194	6149259	Intron variant	<i>PFKFB3</i>
rs7092757	30543292	Intergenic variant	<i>None</i>
rs72826094	113041729	Intron variant	<i>TCF7L2</i>
Chr 11			
rs1002171	71506525	Intergenic variant	<i>None</i>
rs2434468	43936390	Intergenic variant	<i>None</i>
rs2511241	73234296	Missense variant	<i>P2RY2</i>
rs3741392	64933558	Intron variant	<i>PPP2R5B</i>
rs61899280	46945082	Intron variant	<i>C11orf49</i>
Chr 12			
rs10506726	77073285	Intergenic variant	<i>None</i>
rs11171745	56118887	Intron variants	<i>ZC3H10</i>
rs11171773	56189702	Upstream variant	<i>SMARCC2; LOC107984468</i>
rs773643	56181404	Intron variant	<i>SMARCC2</i>
rs116378618	56166019	Intron variant	<i>SMARCC2</i>

(Continued)

Table 3. Continued

	Position	Function	Gene in proximity to the genetic variant
rs1689512	56116853	Intron variant; upstream variant	<i>RPL41; ZC3H10</i>
rs17118317	56126591	Prime UTR variant; up- stream variant	<i>ZC3H10; ESYT1</i>
rs7956913	56129931	Intron variant	<i>ESYT1</i>
rs35436573	56159225	Intron variant	<i>MYL6</i>
rs4762693	21009309	Intron variant	<i>SLCO1B3-SLCO1B7</i>
Chr 13			
rs12872592	21154616	Prime UTR ^a variant	<i>SKA3</i>
Chr 14			
rs4981312	20675000	Intergenic variant	<i>None</i>
rs7155978	68785538	Intergenic variant	<i>None</i>
rs915064	62710859	Intron variant	<i>KCNH5</i>
Chr 16			
rs1003341	25537652	Intergenic variant	<i>None</i>
Chr 17			
rs3744761	45118646	Intron variant	<i>PLCD3</i>
rs4362432	45119179	Intron variant	<i>PLCD3</i>
Chr 20			
rs6032180	45428246	Intron variant	<i>LOC105372631</i>

C-IMT_{max} indicates maximum of carotid intima media thickness values measured at different locations of the carotid tree; and UT, untranslated region.

delta 3 (*PLCD3*) gene, which may be of particular interest due to its link to hypertension. This gene has been identified in the Global Blood Pressure Genetics Consortium genome-wide association study (GWAS) including >34,000 study participants, as one of 8 genes linked to hypertension.⁴⁷ Hypertension, in turn, has been consistently associated with increased C-IMT in several studies including the IMPROVE.^{14,48} The identification of the *PLCD3* gene in the Global Blood Pressure Genetics was not confirmed in a later larger GWAS: the International Consortium for Blood Pressure (~200,000 study participants including also Global Blood Pressure Genetics participants).⁴⁹ A possible explanation for this lack of replication may relate to underlying gene-smoking interaction.

Among the 146 significant interaction results generated from analyses that used the 50th percentile C-IMT_{max} cutoff, 75 are in protein coding genes. Among those, perhaps the most interesting finding involves the *APOB* (apolipoprotein B) gene (rs550619 and rs570877). The *APOB* gene encodes for the well-known *APOB* protein involved in the transportation and metabolism of lipids such as LDL-C, which in turn seems to play a fundamental role in CVD pathophysiology.⁵⁰ Findings from recent Mendelian randomization studies suggest *APOB* as the predominant lipoprotein trait that accounts for a causal mechanism that links LDL-C to CVD.^{51,52} Also, levels of *APOB* have been noted to increase in relation to smoking tobacco,⁵³ however, not consistently.⁵⁴

The remaining significant results (not discussed above) from analyses based on the C-IMT_{max} 75th or 50th percentile cutoffs, involve SNPs located in genes previously discussed in relation to: (1) regulation of cardiometabolic factors and related diseases such as obesity, hypertension and diabetes (eg, *COBLL1; HFM1, CXCR1; COL21A1, DOCK3; DGKB, BMP1; IDE; KCNQ1*; and *KCNQ1-AS1, ZC3H10*),⁵⁵⁻⁶⁴ (2) endothelial inflammation and dysfunction (eg, *TNFAIP8L1, CCNY, GSE1*),⁶⁵⁻⁶⁷ (3) vascular smooth muscle cell proliferation (eg, *VEGFA*),⁶⁸ (4) inflammatory diseases (eg, *PSORS1C1*),⁶⁹ (5) risk of CVD hard end point such as atrial fibrillation and venous thromboembolism (eg, *ZFPM2; LMO7*),^{70,71} and (6) addiction behavior including nicotine dependence (eg, *SP140L, THSD7B*).^{72,73}

From the results of our analyses restricted to cell counts of 10 or below, the identification of a SNP located in the *PIN2/TERF1* interacting, telomerase inhibitor 1 (*PINX1*) gene is potentially interesting, because this gene was previously identified in GWAS of subclinical atherosclerosis⁶ and carotid plaque.⁷ However, it was not found to interact with smoking in a previous study on C-IMT using a candidate gene approach.³³ The study addressed multiplicative interactions only.

An important advantage of our study is that we did not limit the gene-smoking interaction analyses to involve SNPs identified in previous GWAS of C-IMT. It is possible that a gene itself is not associated with C-IMT but becomes important only when smoke exposure occurs. Interestingly, none of the




Figure 1. Visualization of 8 selected significant results from interaction analyses.

The bars show odds ratio point estimates for the risk of having carotid intima-media thickness (C-IMT) above the 75th percentile associated with (a) the genetic risk variant without the presence of smoking, (b) smoking without the presence of the genetic risk variant, and (c) the genetic risk variant in combination with smoking. Reference category is nonsmoking without the presence of the genetic risk variant. These 8 SNPs are located in coding genes previously linked to the development of atherosclerosis.

SNPs we have identified as significantly involved in smoking interaction are among the significant findings reported in previous GWAS in relation to C-IMT or smoking behavior.⁵⁷

Limitations

Our study, just like other exploratory studies, cannot determine which findings are truly positive, and as to whether there are other true effects we did not detect. However, we used the most conservative approach available to adjust for multiple testing, which increases the likelihood that reported findings are true positive. Further, to our knowledge, our study is the largest to date investigating gene-smoking interaction in relation to subclinical atherosclerosis with an explorative approach. Interactions we may have failed to identify should be of a smaller magnitude than those we have identified. Concerning our positive findings, replication analyses using an external study material would have been a good

complement. However, no suitable material for replication analyses was available. Another study limitation is that our genetic data were extracted from genetic chips which do not encompass the whole genome; our results are thus limited to genes related to cardiovascular and immunologic traits which means that some of the relevant SNPs related to smoking predisposition may not have been included. An additional limitation is that our results may not be generalized to populations other than those with European ancestry and at high risk of CVD. Finally, there is also a limitation linked to the fact that our chosen method for interaction analyses requires dichotomization of exposure variables; the results may have been diluted because we included former smokers in the same category as the current smokers. However, smoking cessation is considered a risk factor for CVD.⁷⁴ Further, studies on the relation between smoking cessation and C-IMT have not shown any clear decreased risk of C-IMT progression.⁷⁵

Conclusions

In this European population at high risk of CVD, we identified several significant gene-smoking interactions in relation to C-IMT. Further research in this field is urged to build strong scientific evidence that may open new possibilities for improving cardiovascular prevention through personalized recommendations or drug development.

ARTICLE INFORMATION

Received January 10, 2022; accepted January 29, 2023.

Affiliations

Department of Cardiology, First Affiliated Hospital of Xinjiang Medical University, Urumqi, China (B.M.). Unit of Cardiovascular & Nutritional Epidemiology, Institute of Environmental Medicine (F.L., U.d.F., K.L.) and Cardiovascular Medicine Unit, Department of Medicine Solna (R.J.S., B.G.), Karolinska Institutet, Stockholm, Sweden. Mental Health & Wellbeing, Institute of Mental Health & Wellbeing, University of Glasgow (R.J.S.). Health Data Research, United Kingdom (R.J.S.). Department of Medical Biotechnology & Translational Medicine, Università degli Studi di Milano (D.B.). Centro Cardiologico Monzino, IRCCS, Milan, Italy (D.B., F.V., E.T.). Cardiovascular Genetics, Institute Cardiovascular Science, University College London, United Kingdom (S.E.H.). Foundation for Research in Health Exercise & Nutrition, Kuopio & Research Institute of Exercise Medicine, Kuopio, Finland (K.S.) and Department of Clinical Physiology & Nuclear Medicine, Kuopio University Hospital (K.S.). Institute of Public Health & Clinical Nutrition, University of Eastern Finland, Kuopio (S.K.). Unit of Internal Medicine, Angiology & Arteriosclerosis Diseases, Department of Medicine, University of Perugia, Italy (M.P.). Department of Medicine, University Medical Center Groningen, the Netherlands (A.J.S.). Unités de Prévention Cardiovasculaire, Assistance Publique-Hôpitaux de Paris, Service Endocrinologie-Métabolisme, Groupe Hospitalier Pitié-Salpêtrière, France (P.G.). Cardiovascular Medicine Unit, Department of Medicine Solna, Karolinska Institutet & Karolinska Hospital, Stockholm, Sweden (A.S., A.H.).

Acknowledgments

The authors thank all the study participants of the IMPROVE study (Carotid Intima Media Thickness [IMT] and IMT-Progression as Predictors of Vascular Events in a High Risk European Population). The authors also thank Max Vikström and Paolo Frumento for their statistical support.

Sources of Funding

The IMPROVE study (Carotid Intima Media Thickness [IMT] and IMT-Progression as Predictors of Vascular Events in a High Risk European Population) was supported by the European Commission (contract number: QLGI-CT-2002-00896; to Drs Tremoli, Baldassarre, Giral, Kurl, Pirro); Ministero della Salute Ricerca Corrente, Italy (2017; 2018; to Dr Baldassarre). The IMPROVE was also funded by the Swedish Research Council (8691 and 0593), European Commission (contract number: QLGI-CT-2002-00896), the Swedish Heart-Lung Foundation, the Swedish Foundation for Strategic Research, the Stockholm County Council (project 562183), and the British Heart Foundation (RG2008/008). The present study was supported by the Swedish Research Council (project 2016-02815 to Dr Leander). Dr Strawbridge is supported by United Kingdom Research and Innovation-Health Data Research United Kingdom Fellowship (MR/S003061/1).

Disclosures

None.

Supplemental Material

Supplemental Methods
Tables S1–S6
References^{76–86}

REFERENCES

1. Baber U, Mehran R, Sartori S, Schoos MM, Sillesen H, Muntendam P, Garcia MJ, Gregson J, Pocock S, Falk E, et al. Prevalence, impact, and predictive value of detecting subclinical coronary and carotid atherosclerosis in asymptomatic adults: the Biolmage study. *J Am Coll Cardiol*. 2015;65:1065–1074. doi: 10.1016/j.jacc.2015.01.017
2. Baldassarre D, Hamsten A, Veglia F, de Faire U, Humphries SE, Smit AJ, Giral P, Kurl S, Rauramaa R, Mannarino E, et al; IMPROVE Study Group. Measurements of carotid intima-media thickness and of interadventitia common carotid diameter improve prediction of cardiovascular events: results of the IMPROVE (Carotid Intima Media Thickness [IMT] and IMT-Progression as Predictors of Vascular Events in a High Risk European Population) study. *J Am Coll Cardiol*. 2012;60:1489–1499. doi: 10.1016/j.jacc.2012.06.034
3. Nambi V, Chambless L, Folsom AR, He M, Hu Y, Mosley T, Volcik K, Boerwinkle E, Ballantyne CM. Carotid intima-media thickness and presence or absence of plaque improves prediction of coronary heart disease risk: the ARIC (Atherosclerosis Risk In Communities) study. *J Am Coll Cardiol*. 2010;55:1600–1607. doi: 10.1016/j.jacc.2009.11.075
4. Nambi V, Chambless L, He M, Folsom AR, Mosley T, Boerwinkle E, Ballantyne CM. Common carotid artery intima-media thickness is as good as carotid intima-media thickness of all carotid artery segments in improving prediction of coronary heart disease risk in the Atherosclerosis Risk in Communities (ARIC) study. *Eur Heart J*. 2012;33:183–190. doi: 10.1093/eurheartj/ehr192
5. Strawbridge RJ, Ward J, Bailey MES, Cullen B, Ferguson A, Graham N, Johnston KJA, Lyall LM, Pearsall R, Pell J, et al. Carotid intima-media thickness: novel loci, sex-specific effects, and genetic correlations with obesity and glucometabolic traits in UKB. *Arterioscler Thromb Vasc Biol*. 2020;40:446–461. doi: 10.1161/ATVBAHA.119.313226
6. Bis JC, Kavousi M, Franceschini N, Isaacs A, Abecasis GR, Schminke U, Post WS, Smith AV, Cupples LA, Markus HS, et al; CARDIoGRAM Consortium. Meta-analysis of genome-wide association studies from the CHARGE consortium identifies common variants associated with carotid intima media thickness and plaque. *Nat Genet*. 2011;43:940–947. doi: 10.1038/ng.920
7. Franceschini N, Giambartolomei C, de Vries PS, Finan C, Bis JC, Huntley RP, Lovering RC, Tajuddin SM, Winkler TW, Graff M, et al; MEGASTROKE Consortium. GWAS and colocalization analyses implicate carotid intima-media thickness and carotid plaque loci in cardiovascular outcomes. *Nat Commun*. 2018;9:5141. doi: 10.1038/s41467-018-07340-5
8. Forgo B, Medda E, Hernyes A, Szalontai L, Tarnoki DL, Tarnoki AD. Carotid artery atherosclerosis: a review on heritability and genetics. *Twin Res Hum Genet*. 2018;21:333–346. doi: 10.1017/thg.2018.45
9. Sugiura T, Dohi Y, Takagi Y, Yokochi T, Yoshikane N, Suzuki K, Tomiishi T, Nagami T, Iwase M, Takase H, et al. Close association between subclinical atherosclerosis and pulmonary function in middle-aged male smokers. *J Atheroscler Thromb*. 2020;27:1230–1242. doi: 10.5551/jat.55996
10. Larsson SC, Mason AM, Back M, Klarin D, Damrauer SM, Million Veteran P, Michaelsson K, Burgess S. Genetic predisposition to smoking in relation to 14 cardiovascular diseases. *Eur Heart J*. 2020;41:3304–3310. doi: 10.1093/eurheartj/ehaa193
11. Munzel T, Hahad O, Kuntic M, Keaney JF, Deanfield JE, Daiber A. Effects of tobacco cigarettes, e-cigarettes, and waterpipe smoking on endothelial function and clinical outcomes. *Eur Heart J*. 2020;41:4057–4070. doi: 10.1093/eurheartj/ehaa460
12. Kweon SS, Lee YH, Shin MH, Choi JS, Rhee JA, Choi SW, Ryu SY, Kim BH, Nam HS, Jeong SK, et al. Effects of cumulative smoking exposure and duration of smoking cessation on carotid artery structure. *Circ J*. 2012;76:2041–2047. doi: 10.1253/circj.cj-11-1353
13. Rose JE, Behm FM, Drgon T, Johnson C, Uhl GR. Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. *Mol Med*. 2010;16:247–253. doi: 10.2119/molmed.2009.00159
14. Baldassarre D, Nyyssonen K, Rauramaa R, de Faire U, Hamsten A, Smit AJ, Mannarino E, Humphries SE, Giral P, Grossi E, et al; IMPROVE study group. Cross-sectional analysis of baseline data to identify the major determinants of carotid intima-media thickness in a European population: the IMPROVE study. *Eur Heart J*. 2010;31:614–622. doi: 10.1093/eurheartj/ehp496
15. Wei Q, Doris PA, Pollizotto MV, Boerwinkle E, Jacobs DR Jr, Siscovick DS, Fornage M. Sequence variation in the soluble epoxide hydrolase gene and subclinical coronary atherosclerosis: interaction with cigarette smoking. *Atherosclerosis*. 2007;190:26–34. doi: 10.1016/j.atherosclerosis.2006.02.021
16. Wang XL, Greco M, Sim AS, Duarte N, Wang J, Wilcken DE. Effect of CYP1A1 MspI polymorphism on cigarette smoking related coronary artery disease and diabetes. *Atherosclerosis*. 2002;162:391–397. doi: 10.1016/s0021-9150(01)00723-7
17. Srinivasan SR, Li S, Chen W, Tang R, Bond MG, Boerwinkle E, Berenson GS. Q192R polymorphism of the paraoxanase 1 gene and its association with serum lipoprotein variables and carotid artery intima-media thickness in young adults from a biracial community. The Bogalusa Heart Study. *Atherosclerosis*. 2004;177:167–174. doi: 10.1016/j.atherosclerosis.2004.06.013

18. Rosner SA, Ridker PM, Zee RY, Cook NR. Interaction between inflammation-related gene polymorphisms and cigarette smoking on the risk of myocardial infarction in the Physician's Health Study. *Hum Genet.* 2005;118:287–294. doi: 10.1007/s00439-005-0052-6
19. Rios DL, D'Onofrio LO, Souza JK, Queiroz AM, Raduy-Marón L, Silva-Neto N, Carvalho HG, Santos-Filho A, Galvao-Castro B. Smoking-dependent and haplotype-specific effects of endothelial nitric oxide synthase gene polymorphisms on angiographically assessed coronary artery disease in Caucasian- and African-Brazilians. *Atherosclerosis.* 2007;193:135–141. doi: 10.1016/j.atherosclerosis.2006.05.041
20. Payne JR, Dhamrait SS, Toor IS, Cooper J, Jones A, Miller GJ, Humphries SE, Montgomery HE. The -344T>C promoter variant of the gene for aldosterone synthase (CYP11B2) is not associated with cardiovascular risk in a prospective study of UK healthy men. *Atherosclerosis.* 2004;174:81–86. doi: 10.1016/j.atherosclerosis.2004.01.004
21. Olshan AF, Li R, Pankow JS, Bray M, Tyroler HA, Chambless LE, Boerwinkle E, Pittman GS, Bell DA. Risk of atherosclerosis: interaction of smoking and glutathione S-transferase genes. *Epidemiology.* 2003;14:321–327. doi: 10.1097/01.EDE.0000059229.74889.CF
22. North KE, Carr JJ, Borecki IB, Kraja A, Province M, Pankow JS, Wilk JB, Hixson JE, Heiss G; Investigators FHS. QTL-specific genotype-by-smoking interaction and burden of calcified coronary atherosclerosis: the NHLBI Family Heart Study. *Atherosclerosis.* 2007;193:11–19. doi: 10.1016/j.atherosclerosis.2006.08.015
23. Malin R, Loimaala A, Nenonen A, Mercuri M, Vuori I, Pasanen M, Oja P, Bond G, Koivula T, Lehtimäki T. Relationship between high-density lipoprotein paraoxonase gene M/L55 polymorphism and carotid atherosclerosis differs in smoking and nonsmoking men. *Metabolism.* 2001;50:1095–1101. doi: 10.1053/meta.2001.25641
24. Lee CR, North KE, Bray MS, Avery CL, Mosher MJ, Couper DJ, Coresh J, Folsom AR, Boerwinkle E, Heiss G, et al. NOS3 polymorphisms, cigarette smoking, and cardiovascular disease risk: the Atherosclerosis Risk in Communities study. *Pharmacogenet Genomics.* 2006;16:891–899. doi: 10.1097/01.fpc.0000236324.96056.16
25. Karvonen J, Kauma H, Kervinen K, Ukkola O, Rantala M, Paivansalo M, Savolainen MJ, Kesäniemi YA. Apolipoprotein E polymorphism affects carotid artery atherosclerosis in smoking hypertensive men. *J Hypertens.* 2002;20:2371–2378. doi: 10.1097/00004872-200212000-00015
26. Jerrard-Dunne P, Sitzer M, Risleyp P, Buehler A, von Kegler S, Markus HS. Inflammatory gene load is associated with enhanced inflammation and early carotid atherosclerosis in smokers. *Stroke.* 2004;35:2438–2443. doi: 10.1161/01.STR.0000144681.46696.b3
27. Inamoto N, Katsuya T, Kokubo Y, Mannami T, Asai T, Baba S, Ogata J, Tomoike H, Ogiwara T. Association of methylenetetrahydrofolate reductase gene polymorphism with carotid atherosclerosis depending on smoking status in a Japanese general population. *Stroke.* 2003;34:1628–1633. doi: 10.1161/01.STR.0000075769.09092.82
28. Djousse L, Myers RH, Province MA, Hunt SC, Eckfeldt JH, Evans G, Peacock JM, Ellison RC. Influence of apolipoprotein E, smoking, and alcohol intake on carotid atherosclerosis: National Heart, Lung, and Blood Institute Family Heart Study. *Stroke.* 2002;33:1357–1361. doi: 10.1161/01.str.0000014325.54063.1a
29. Viiri LE, Viiri KM, Ilveskoski E, Huhtala H, Maki M, Tienari PJ, Perola M, Lehtimäki T, Karhunen PJ. Interactions of functional apolipoprotein E gene promoter polymorphisms with smoking on aortic atherosclerosis. *Circ Cardiovasc Genet.* 2008;1:107–116. doi: 10.1161/CIRCGENETICS.108.791764
30. Fan M, Raitakari OT, Kahonen M, Juonala M, Hutri-Kahonen N, Porsti I, Viikari J, Lehtimäki T. The association between cigarette smoking and carotid intima-media thickness is influenced by the -930A/G CYBA gene polymorphism: the Cardiovascular Risk in Young Finns Study. *Am J Hypertens.* 2009;22:281–287. doi: 10.1038/ajh.2008.349
31. Luo S, Wang F, Li Z, Deng J. Effect of the +781C/T polymorphism in the interleukin-8 gene on atherosclerotic cerebral infarction, and its interaction with smoking and drinking. *PLoS One.* 2013;8:e80246. doi: 10.1371/journal.pone.0080246
32. Niemiec P, Nowak T, Iwanicki T, Krauze J, Gorczynska-Kosiorz S, Grzeszczak W, Ochalska-Tyka A, Zak I. The -930A>G polymorphism of the CYBA gene is associated with premature coronary artery disease. A case-control study and gene-risk factors interactions. *Mol Biol Rep.* 2014;41:3287–3294. doi: 10.1007/s11033-014-3191-9
33. Li C, Chen W, Jiang F, Simino J, Srinivasan SR, Berenson GS, Mei H. Genetic association and gene-smoking interaction study of carotid intima-media thickness at five GWAS-indicated genes: the Bogalusa Heart Study. *Gene.* 2015;562:226–231. doi: 10.1016/j.gene.2015.02.078
34. Wang L, Rundek T, Beecham A, Hudson B, Blanton SH, Zhao H, Sacco RL, Dong C. Genome-wide interaction study identifies RCBTB1 as a modifier for smoking effect on carotid intima-media thickness. *Arterioscler Thromb Vasc Biol.* 2014;34:219–225. doi: 10.1161/ATVBAHA.113.302706
35. Boua PR, Brandenburg JT, Choudhury A, Hazelhurst S, Sengupta D, Agongo G, Nonterah EA, Odoro AR, Tinto H, Mathew CG, et al. Novel and known gene-smoking interactions with cIMT identified as potential drivers for atherosclerosis risk in West-African Populations of the AWI-Gen Study. *Front Genet.* 2019;10:1354. doi: 10.3389/fgene.2019.01354
36. Liu W, Chang C, Hu H, Yang H. Interleukin-23: a new atherosclerosis target. *J Interferon Cytokine Res.* 2018;38:440–444. doi: 10.1089/jir.2018.0006
37. Jin L, Hastings NE, Blackman BR, Somlyo AV. Mechanical properties of the extracellular matrix alter expression of smooth muscle protein LPP and its partner palladin; relationship to early atherosclerosis and vascular injury. *J Muscle Res Cell Motil.* 2009;30:41–55. doi: 10.1007/s10974-009-9173-1
38. Hsieh J, Koseki M, Molusky MM, Yakushiji E, Ichi I, Westerterp M, Iqbal J, Chan RB, Abramowicz S, Tascau L, et al. TTC39B deficiency stabilizes LXR reducing both atherosclerosis and steatohepatitis. *Nature.* 2016;535:303–307. doi: 10.1038/nature18628
39. Chen X, Qian S, Hoggatt A, Tang H, Hacker TA, Obukhov AG, Herring PB, Seye CI. Endothelial cell-specific deletion of P2Y2 receptor promotes plaque stability in atherosclerosis-susceptible ApoE-null mice. *Arterioscler Thromb Vasc Biol.* 2017;37:75–83. doi: 10.1161/ATVBAHA.116.308561
40. Poels K, Schnitzler JG, Waissi F, Levels JHM, Stroes ESG, Daemen M, Lutgens E, Pennekamp AM, De Kleijn DPV, Seijkens TTP, et al. Inhibition of PFKFB3 hampers the progression of atherosclerosis and promotes plaque stability. *Front Cell Dev Biol.* 2020;8:581641. doi: 10.3389/fcell.2020.581641
41. Sabater-Lleal M, Malarstig A, Folkersen L, Soler Artigas M, Baldassarre D, Kavousi M, Almgren P, Veglia F, Brusselle G, Hofman A, et al. Common genetic determinants of lung function, subclinical atherosclerosis and risk of coronary artery disease. *PLoS One.* 2014;9:e104082. doi: 10.1371/journal.pone.0104082
42. O'Donnell CJ, Cupples LA, D'Agostino RB, Fox CS, Hoffmann U, Hwang SJ, Ingelsson E, Liu C, Murabito JM, Polak JF, et al. Genome-wide association study for subclinical atherosclerosis in major arterial territories in the NHLBI's Framingham Heart Study. *BMC Med Genet.* 2007;8:S4. doi: 10.1186/1471-2350-8-S1-S4
43. Rivera NV, Patasova K, Kullberg S, Diaz-Gallo LM, Iseda T, Bengtsson C, Alfredsson L, Eklund A, Kockum I, Grunewald J, et al. A gene-environment interaction between smoking and gene polymorphisms provides a high risk of two subgroups of sarcoidosis. *Sci Rep.* 2019;9:18633. doi: 10.1038/s41598-019-54612-1
44. Karami J, Aslani S, Jamshidi A, Garshasbi M, Mahmoudi M. Genetic implications in the pathogenesis of rheumatoid arthritis; an updated review. *Gene.* 2019;702:8–16. doi: 10.1016/j.gene.2019.03.033
45. Doecke JD, Simms LA, Zhao ZZ, Roberts RL, Fowler EV, Croft A, Lin A, Huang N, Whiteman DC, Florin TH, et al. Smoking behaviour modifies IL23r-associated disease risk in patients with Crohn's disease. *J Gastroenterol Hepatol.* 2015;30:299–307. doi: 10.1111/jgh.12674
46. Khakh BS, North RA. P2X receptors as cell-surface ATP sensors in health and disease. *Nature.* 2006;442:527–532. doi: 10.1038/nature04886
47. Newton-Cheh C, Johnson T, Gateva V, Tobin MD, Bochud M, Coin L, Najjar SS, Zhao JH, Heath SC, Eyheramendy S, et al; Wellcome Trust Case Control Consortium. Genome-wide association study identifies eight loci associated with blood pressure. *Nat Genet.* 2009;41:666–676. doi: 10.1038/ng.361
48. Ferreira JP, Girerd N, Bozec E, Machu JL, Boivin JM, London GM, Zannad F, Rossignol P. Intima-media thickness is linearly and continuously associated with systolic blood pressure in a population-based cohort (STANISLAS Cohort Study). *J Am Heart Assoc.* 2016;5:e003529. doi: 10.1161/JAHA.116.003529
49. Ehret GB, Munroe PB, Rice KM, Bochud M, Johnson AD, Chasman DI, Smith AV, Tobin MD, Verwoert GC, Hwang SJ, et al; International Consortium for Blood Pressure Genome-Wide Association S. Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature.* 2011;478:103–109. doi: 10.1038/nature10405
50. Fulcher J, O'Connell R, Voysey M, Emberson J, Blackwell L, Mihaylova B, Simes J, Collins R, Kirby A, Colhoun H, et al; Cholesterol Treatment Trialists C. Efficacy and safety of LDL-lowering therapy among men and women: meta-analysis of individual data from 174,000 participants in 27 randomised trials. *Lancet.* 2015;385:1397–1405. doi: 10.1016/S0140-6736(14)61368-4
51. Zuber V, Gill D, Ala-Korpela M, Langenberg C, Butterworth A, Bottolo L, Burgess S. High-throughput multivariable Mendelian randomization analysis

- prioritizes apolipoprotein B as key lipid risk factor for coronary artery disease. *Int J Epidemiol*. 2021;50:893–901. doi: 10.1093/ije/dyaa216
52. Richardson TG, Sanderson E, Palmer TM, Ala-Korpela M, Ference BA, Davey Smith G, Holmes MV. Evaluating the relationship between circulating lipoprotein lipids and apolipoproteins with risk of coronary heart disease: a multivariable Mendelian randomisation analysis. *PLoS Med*. 2020;17:e1003062. doi: 10.1371/journal.pmed.1003062
 53. Lubin JH, Couper D, Lutsey PL, Yatsuya H. Synergistic and non-synergistic associations for cigarette smoking and non-tobacco risk factors for cardiovascular disease incidence in the Atherosclerosis Risk In Communities (ARIC) study. *Nicotine Tob Res*. 2017;19:826–835. doi: 10.1093/ntr/ntw235
 54. Jain RB. Impact of smoking on the observed levels of apolipoprotein B: data from NHANES 2007–2012. *Environ Toxicol Pharmacol*. 2017;53:227–233. doi: 10.1016/j.etap.2017.06.006
 55. Perez-Montarelo D, Madsen O, Alves E, Rodriguez MC, Folch JM, Noguera JL, Groenen MA, Fernandez AI. Identification of genes regulating growth and fatness traits in pig through hypothalamic transcriptome analysis. *Physiol Genomics*. 2014;46:195–206. doi: 10.1152/physiolgenomics.00151.2013
 56. Kraja AT, Chasman DI, North KE, Reiner AP, Yanek LR, Kilpelainen TO, Smith JA, Dehghan A, Dupuis J, Johnson AD, et al; Cross Consortia Pleiotropy Group. Pleiotropic genes for metabolic syndrome and inflammation. *Mol Genet Metab*. 2014;112:317–338. doi: 10.1016/j.ymgme.2014.04.007
 57. Coulson DJ, Bakhshab S, Latief JS, Weaver JU. MiR-126, IL-7, CXCR1/2 receptors, inflammation and circulating endothelial progenitor cells: the study on targets for treatment pathways in a model of subclinical cardiovascular disease (type 1 diabetes mellitus). *J Transl Med*. 2021;19:140. doi: 10.1186/s12967-021-02785-7
 58. Chiu YF, Chung RH, Lee CY, Kao HY, Hou L, Hsu FC. Identification of rare variants for hypertension with incorporation of linkage information. *BMC Proc*. 2014;8:S109. doi: 10.1186/1753-6561-8-S1-S109
 59. Surendran P, Drenos F, Young R, Warren H, Cook JP, Manning AK, Grarup N, Sim X, Barnes DR, Witkowska K, et al; CHARGE-Heart Failure Consortium. Trans-ancestry meta-analyses identify rare and common variants associated with blood pressure and hypertension. *Nat Genet*. 2016;48:1151–1161. doi: 10.1038/ng.3654
 60. Benn M, Tybjaerg-Hansen A, McCarthy MI, Jensen GB, Grande P, Nordestgaard BG. Nonfasting glucose, ischemic heart disease, and myocardial infarction: a Mendelian randomization study. *J Am Coll Cardiol*. 2012;59:2356–2365. doi: 10.1016/j.jacc.2012.02.043
 61. Banerjee S, Andrew RJ, Duff CJ, Fisher K, Jackson CD, Lawrence CB, Maeda N, Greenspan DS, Kellett KAB, Hooper NM. Proteolysis of the low density lipoprotein receptor by bone morphogenetic protein-1 regulates cellular cholesterol uptake. *Sci Rep*. 2019;9:11416. doi: 10.1038/s41598-019-47814-0
 62. Borges DO, Patarrao RS, Ribeiro RT, de Oliveira RM, Duarte N, Belew GD, Martins M, Andrade R, Costa J, Correia I, et al. Loss of postprandial insulin clearance control by Insulin-degrading enzyme drives dysmetabolism traits. *Metabolism*. 2021;118:154735. doi: 10.1016/j.metabol.2021.154735
 63. Chiou J, Zeng C, Cheng Z, Han JY, Schlichting M, Miller M, Mendez R, Huang S, Wang J, Sui Y, et al. Single-cell chromatin accessibility identifies pancreatic islet cell type- and state-specific regulatory programs of diabetes risk. *Nat Genet*. 2021;53:455–466. doi: 10.1038/s41588-021-00823-0
 64. Audano M, Pedretti S, Ligorio S, Gualdrini F, Polletti S, Russo M, Ghisletti S, Bean C, Crestani M, Caruso D, et al. Zc3h10 regulates adipogenesis by controlling translation and F-actin/mitochondria interaction. *J Cell Biol*. 2021;220:e202003173. doi: 10.1083/jcb.202003173
 65. Shao J, Li Y, Zhou C, Geng M, Zhang G, Zhang N, Jin G, Zhang L, Gao C, Liu S. TIPE1 accelerates atherogenesis by inducing endothelial dysfunction in response to oxidative stress. *Biochim Biophys Acta Mol Basis Dis*. 2020;1866:165578. doi: 10.1016/j.bbadis.2019.165578
 66. Kyselova A, Siragusa M, Anthes J, Solari FA, Loroch S, Zahedi RP, Walter U, Fleming I, Randriamboavonjy V. Cyclin Y is expressed in platelets and modulates integrin outside-in signaling. *Int J Mol Sci*. 2020;21:8239. doi: 10.3390/ijms21218239
 67. Zheng ML, Du XP, Zhao L, Yang XC. Expression profile of circular RNAs in epicardial adipose tissue in heart failure. *Chin Med J (Engl)*. 2020;133:2565–2572. doi: 10.1097/CM9.0000000000001056
 68. Yang Y, Mao W, Wang L, Lu L, Pang Y. Circular RNA circLMF1 regulates PDGF-BB-induced proliferation and migration of human aortic smooth muscle cells by regulating the miR-125a-3p/VEGFA or FGF1 axis. *Clin Hemorheol Microcirc*. 2022;80:167–183. doi: 10.3233/CH-211166
 69. Butt SA, Jeppesen JL, Torp-Pedersen C, Sam F, Gislasen GH, Jacobsen S, Andersson C. Cardiovascular manifestations of systemic sclerosis: a Danish Nationwide Cohort Study. *J Am Heart Assoc*. 2019;8:e013405. doi: 10.1161/JAHA.119.013405
 70. Klarin D, Emdin CA, Natarajan P, Conrad MF, Consortium I, Kathiresan S. Genetic analysis of venous thromboembolism in UK biobank identifies the ZFPM2 locus and implicates obesity as a causal risk factor. *Circ Cardiovasc Genet*. 2017;10:e001643. doi: 10.1161/CIRCGENETICS.116.001643
 71. Li MY, Chen HX, Hou HT, Wang J, Liu XC, Yang Q, He GW. Biomarkers and key pathways in atrial fibrillation associated with mitral valve disease identified by multi-omics study. *Ann Transl Med*. 2021;9:393. doi: 10.21037/atm-20-3767
 72. Chen J, Loukola A, Gillespie NA, Peterson R, Jia P, Riley B, Maes H, Dick DM, Kendler KS, Damaj MI, et al. Genome-wide meta-analyses of FTND and TTFC phenotypes. *Nicotine Tob Res*. 2020;22:900–909. doi: 10.1093/ntr/ntz099
 73. McGue M, Zhang Y, Miller MB, Basu S, Vrieze S, Hicks B, Malone S, Oetting WS, Iacono WG. A genome-wide association study of behavioral disinhibition. *Behav Genet*. 2013;43:363–373. doi: 10.1007/s10519-013-9606-x
 74. Duncan MS, Freiberg MS, Greevy RA Jr, Kundu S, Vasan RS, Tindle HA. Association of smoking cessation with subsequent risk of cardiovascular disease. *JAMA*. 2019;322:642–650. doi: 10.1001/jama.2019.10298
 75. Hansen K, Ostling G, Persson M, Nilsson PM, Melander O, Engstrom G, Hedblad B, Rosvall M. The effect of smoking on carotid intima-media thickness progression rate and rate of lumen diameter reduction. *Eur J Intern Med*. 2016;28:74–79. doi: 10.1016/j.ejim.2015.10.018
 76. Baldassarre D, Veglia F, Hamsten A, Humphries SE, Rauramaa R, de Faire U, Smit AJ, Giral P, Kurl S, Mannarino E, et al. Progression of carotid intima-media thickness as predictor of vascular events: results from the IMPROVE study. *Arterioscler Thromb Vasc Biol*. 2013;33:2273–2279. doi: 10.1161/atvbaha.113.301844
 77. Amato M, Veglia F, de Faire U, Giral P, Rauramaa R, Smit AJ, Kurl S, Ravani A, Frigerio B, Sansaro D, et al; IMPROVE study group. Carotid plaque thickness and common carotid IMT show additive value in cardiovascular risk prediction and reclassification. *Atherosclerosis*. 2017;263:412–419. doi: 10.1016/j.atherosclerosis.2017.05.023
 78. Andersson T, Alfredsson L, Kallberg H, Zdravkovic S, Ahlbom A. Calculating measures of biological interaction. *Eur J Epidemiol*. 2005;20:575–579. doi: 10.1007/s10654-005-7835-x
 79. Stein JH, Korcarz CE, Hurst RT, Lonn E, Kendall CB, Mohler ER, Najjar SS, Rembold CM, Post WS; American Society of Echocardiography Carotid Intima-Media Thickness Task Force. Use of carotid ultrasound to identify subclinical vascular disease and evaluate cardiovascular disease risk: a consensus statement from the American Society of Echocardiography Carotid Intima-Media Thickness Task Force. Endorsed by the Society for Vascular Medicine. *J Am Soc Echocardiogr*. 2008;21:93–111; quiz 189. doi: 10.1016/j.echo.2007.11.011
 80. Voight BF, Kang HM, Ding J, Palmer CD, Sidore C, Chines PS, Burt NP, Fuchsberger C, Li Y, Erdmann J, et al. The metabochip, a custom genotyping array for genetic studies of metabolic, cardiovascular, and anthropometric traits. *PLoS Genet*. 2012;8:e1002793. doi: 10.1371/journal.pgen.1002793
 81. Trynka G, Hunt KA, Bockett NA, Romanos J, Mistry V, Szperl A, Bakker SF, Bardella MT, Bhaw-Rosun L, Castillejo G, et al; Spanish Consortium on the Genetics of Coeliac Disease (CEGEC). Dense genotyping identifies and localizes multiple common and rare variant association signals in celiac disease. *Nat Genet*. 2011;43:1193–1201. doi: 10.1038/ng.998
 82. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*. 2007;81:559–575. doi: 10.1086/519795
 83. Rothman KJ, Lash TL. *Modern Epidemiology*. Lippincott William and Wilkins; 2008.
 84. Hosmer DW, Lemeshow S. Confidence interval estimation of interaction. *Epidemiology*. 1992;3:452–456. doi: 10.1097/00001648-199209000-00012
 85. Veglia F, Baldassarre D, de Faire U, Kurl S, Smit AJ, Rauramaa R, Giral P, Amato M, Di Minno A, Ravani A, et al. A priori-defined Mediterranean-like dietary pattern predicts cardiovascular events better in north Europe than in Mediterranean countries. *Int J Cardiol*. 2019;282:88–92. doi: 10.1016/j.ijcard.2018.11.124
 86. Ding B, Kallberg H, Klaveskog L, Padyukov L, Alfredsson L. GEIRA: gene-environment and gene-gene interaction research application. *Eur J Epidemiol*. 2011;26:557–561. doi: 10.1007/s10654-011-9582-5