

RESEARCH ARTICLE

Network approaches and interventions in healthcare settings: A systematic scoping review

Ameneh Ghazal Saatchi^{1*}, Francesca Pallotti², Paul Sullivan^{3,4}

1 Imperial College London, London, United Kingdom, **2** Department of Business, Operations and Strategy, University of Greenwich, London, United Kingdom, **3** NIHR ARC Northwest London, Imperial College London, London, United Kingdom, **4** University Sussex Hospitals NHS Foundation Trust, Sussex, United Kingdom

* AGSaatchi@gmail.com

Abstract

Introduction

The growing interest in networks of interactions is sustained by the conviction that they can be leveraged to improve the quality and efficiency of healthcare delivery systems. Evidence in support of this conviction, however, is mostly based on descriptive studies. Systematic evaluation of the outcomes of network interventions in healthcare settings is still wanting. Despite the proliferation of studies based on Social Network Analysis (SNA) tools and techniques, we still know little about how intervention programs aimed at altering existing patterns of social interaction among healthcare providers affect the quality of service delivery. We update and extend prior reviews by providing a comprehensive assessment of available evidence.

Methods and findings

We searched eight databases to identify papers using SNA in healthcare settings published between 1st January 2010 and 1st May 2022. We followed Chambers et al.'s (2012) approach, using a Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) checklist. We distinguished between studies relying on SNA as part of an intervention program, and studies using SNA for descriptive purposes only. We further distinguished studies recommending a possible SNA-based intervention. We restricted our focus on SNA performed on networks among healthcare professionals (e.g., doctors, nurses, etc.) in any healthcare setting (e.g., hospitals, primary care, etc.). Our final review included 102 papers. The majority of the papers used SNA for descriptive purposes only. Only four studies adopted SNA as an intervention tool, and measured outcome variables.

Conclusions

We found little evidence for SNA-based intervention programs in healthcare settings. We discuss the reasons and challenges, and identify the main component elements of a

OPEN ACCESS

Citation: Saatchi AG, Pallotti F, Sullivan P (2023) Network approaches and interventions in healthcare settings: A systematic scoping review. PLoS ONE 18(2): e0282050. <https://doi.org/10.1371/journal.pone.0282050>

Editor: Shahadat Uddin, The University of Sydney, AUSTRALIA

Received: August 3, 2022

Accepted: February 7, 2023

Published: February 23, 2023

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0282050>

Copyright: © 2023 Saatchi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its [Supporting Information](#) files.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

network intervention plan. Future research should seek to evaluate the long-term role of SNA in changing practices, policies and behaviors, and provide evidence of how these changes affect patients and the quality of service delivery.

Introduction

It is widely recognized that there is a gap between best achievable healthcare outcomes and those that are actually delivered, even in the best funded systems, suggesting that more is required than simply increasing available resources [1, 2]. Improving healthcare outcomes requires changes in frontline clinical practice, which in turn involves the ability to disseminate information across diverse teams, and to engender alignment of multiple groups.

The diffusion of practices and behaviors within any healthcare setting may be usefully framed as a network problem involving multiple individuals and the way they relate and interact with one another. Leaders aiming to improve healthcare outcomes would benefit from understanding how team members interact, and how interactions may be leveraged to optimize the adoption and diffusion of new practices. Information about patterns of interaction can be obtained using Social Network Analysis (SNA). SNA provides a set of tools and techniques used to investigate structural characteristics of networks [3], and understand how a broad range of behaviors may be triggered by social interaction [4]. SNA generates three main types of outputs. The first is a visual representation of networks structures, or network graphs. The second is a set of metrics providing quantitative information on properties of networks, such as density, or properties of individuals, such as centrality. The third type of output is produced by statistical models for network data, such as models for the analysis of longitudinal networks [5].

SNA outputs can be used to inform the design, implementation and monitoring of behavioral change programs, policies and practices [4, 6]. A network intervention can be defined as a structured process using social networks to accelerate behavior change or improve organizational performance [7]. Social networks are channels for information diffusion and interpersonal influence. Hence, changing the wiring of an existing social network may determine changes in how behaviors, ideas and practices spread in a social group.

Valente [7] proposed a taxonomy of four types of network intervention strategies: i) 'Individuals', based upon the identification of individuals with certain network characteristics who are recruited to act as change proponents; ii) 'Segmentation', involving the identification of subgroups in a network on which to focus behavioral change; iii) 'Alteration', whereby an existing network is changed by adding or removing ties or nodes in order to alter patterns of interaction and diffusion, and finally iv) 'Induction', whereby peer-to-peer interactions are encouraged through, for example, the use of meetings or training events bringing previously unconnected people together.

While a large body of research is available that relies on SNA to examine networks of health professionals in healthcare settings, much of this research has been descriptive, with limited reporting of the relationship between network interventions and clinical or organizational outcomes. This is confirmed by recent systematic reviews. For example, Chambers et al.'s [8] systematic scoping review of SNA-based studies in healthcare settings found very little evidence of the use of SNA as part of an intervention. Cunningham et al.'s [9] review (1995–2009) included 40 eligible studies. Only one described an SNA-based intervention using survey data to identify opinion leaders, but did not measure its impact. Bae et al.'s [10] systematic review

included 28 eligible studies (up to 2013), none of which reported on outcomes of SNA-based interventions. A recent umbrella review by Hu et al. [11] included 13 reviews between 2010 and 2019 and demonstrated a wide applicability of SNA to study health professional networks. Of the 330 papers included in the reviews, only one reported on a network intervention.

The aim of the present review is threefold. First, provide an update of prior reviews by searching for papers using SNA to investigate networks of healthcare professionals in healthcare settings. Second, identify research reporting about network-based interventions and their outcomes. Third, identify the component elements and discuss the main challenges of a network intervention strategy to call attention on its potential in healthcare settings. The primary research question that this review seeks to address is what evidence is available on the adoption of network interventions and evaluation of their effect on care processes and outcomes.

Methods

Protocol

The literature review was undertaken in accordance with the protocol (S1 File) followed by Chambers' et al. in their 2012 review [8]. We used the Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) statement and guidelines (S2 File) [12].

Information sources and search strategy

The literature search focused on identifying studies performing SNA on networks of healthcare professionals in healthcare settings. We used the same search strategy, inclusion and exclusion criteria and keywords as those used by Chambers et al. [8]. We performed a systematic electronic database search of OVID MEDLINE (R) ALL first, using free text terms, synonyms and subject headings associated with social networks and the methods used to investigate them including 'sociometrics', 'sociograms' and 'sociomaps'. We also used words associated with SNA software, such as NetDraw and UCINET. Finally, the search strategy included the subject headings inter-professional relations, inter-disciplinary communication and physician-nurse relationships. The search strategy was later adapted for other databases in our search. Specifically, for the period 1st January 2010 to 1st May 2022, we searched the following databases: OVID MEDLINE (R) ALL, EMBASE Classic+EMBASE, APA PsycINFO, Health Management Information Consortium (HMIC), the Cochrane Library (Cochrane Database of Systematic Reviews, Cochrane Protocols and Cochrane Central Register of Controlled Trials), CINAHL Plus, Business Source Ultimate, Social Science Citation Index (SSCI) and Conference Proceedings Citation Index—Social Science & Humanities (CPCI-SSH) databases. Reference lists of relevant reviews and studies were searched, as was the website of the International Network for Social Network analysis (www.insna.org) and its linked sites. The index of contents of the Social Networks journal was also searched. The online search was run on 5th January 2021 and later updated on 1st May 2022 to include papers published up to this date. The search strategy had no study design filters or restrictions to language as long as the paper could be found in English. Records were managed within a Mendeley library.

Eligibility criteria. The review included studies undertaken in any healthcare setting that reported the results of an SNA performed on networks among healthcare professionals (e.g., doctors, nurses, etc.) and other individuals involved in their professional networks (e.g., management, administrative support etc.). Examples of these networks include discussion networks, advice and knowledge sharing, and working on projects together. The healthcare setting was not restricted to a single geographical or organizational location, and could include wider interpersonal networks, such as the Parkinson network [13]. Veterinary or dental

professionals were not included. Studies of networks linking organizations, rather than individuals, were excluded. We excluded studies where network relations were defined solely by patient sharing, as this predicts person-to-person communication only in minority of instances [14].

We built upon Chambers et al.'s [8] classification method. We divided papers into three groups, which we termed level 1 to 3. Level 1 included studies reporting on the impact of an SNA-based intervention. Level 2 included studies describing existing social networks among healthcare professionals without reporting any follow-up action. Level 3 included descriptive studies that went on to suggest an SNA-based intervention intended to affect outcomes and behaviors. We added this additional category to shed light on the significant number of papers acknowledging the value of using SNA to inform the design of intervention plans, and the benefits associated with it.

Study selection and data extraction. Two Authors independently screened studies by title and disregarded those that they agreed to exclude. Studies where there was agreement for inclusion were independently screened by abstract by three Authors. Studies that appeared to meet the review inclusion criteria were forwarded to full-text evaluation and data extraction. The Cochrane EPOC (Effective Practice and Organisation of Care) Group criteria were used to assess the risk of bias by two Authors. Disagreements were discussed with a third Author.

Results

The search returned 31,2867 unique papers, of which 102 met the eligibility criteria. Ten of these [15–24] were also included in Chambers et al.'s [8] review due to a crossover of search periods. We excluded these papers. The PRISMA diagram in Fig 1 below outlines the study selection process, and S1 Table outlines the number of records identified by database with a comparison to Chambers et al.'s [8] review. The comparison seems to suggest an increased use of social network approaches in healthcare studies over the past few years.

Four included studies met the level-1 [13, 25–27], 74 the level-2 [15–17, 19–21, 23, 28–94], and 24 the level-3 [18, 22, 24, 95–115] criteria.

Of the 102 papers, one third ($n = 33$) was conducted in the USA, 22 in Europe (excluding UK), 16 in low- and middle-income countries (LMIC), 11 in Australia, eight in the UK, seven in Canada, two in Japan, two in China and one in Malaysia. The Netherlands and Italy produced the largest number of papers in Europe. Compared to previous reviews mentioned earlier, we found an increased number of studies conducted in LMIC. The largest number of studies ($n = 59$) had participants from multidisciplinary teams, and were conducted in secondary care settings ($n = 64$). The number of participants ranged from 10 [71] to 16,171 [66]. The largest number of studies used surveys/questionnaires ($n = 57$), followed by direct observations ($n = 7$), mixed methods ($n = 13$), process logs or other administrative data ($n = 9$), interviews ($n = 7$), online platforms or forums ($n = 5$), and interaction data collected through sensors ($n = 4$).

We summarized the types of ties examined in the included papers into 10 categories to standardize the language (see S2 Table). We also grouped network measures into 36 categories (see S3 Table). These measures were used across studies to describe or analyze networks at the individual, dyadic, group, and whole network levels. We also created a distinct category for those papers performing only statistical analysis of network data, such as Exponential Random Graph Models (ERGMs), Multiple Quadratic Assignment Procedure (MQAP), and Stochastic Actor Oriented Models (SAOMs). Network visualization was included as a distinct category when it was the only social network method used.




Fig 1. Flow diagram of study selection process.

<https://doi.org/10.1371/journal.pone.0282050.g001>

Level-1 studies

[Table 1](#) below includes the level-1 studies, followed by a descriptive summary.

The four level-1 studies report on the results of SNA as part of an intervention, which we classified according to Valente [7]. Benton et al. [26] employed ‘alteration’ and ‘induction’ strategies by using shared project work to form new connections and increase interactions among network members. van der Eijk et al. [13] employed ‘induction’ through training events. The remaining two studies [25, 27] focused on ‘individuals’, by using social network methods to identify individuals who would act as champions. The impacts reported in the papers included structural network changes as well as changes in working practices and, in one study, staff safety outcomes. None of the studies reported on the impact on patient outcomes. The overall aim of the reported interventions was to improve organizational performance [26], patient care across the Parkinson’s network [13], safe patient manual handling [25], and hand hygiene [27]. All four papers used the information from SNA to improve connectedness within the networks. A summary of the level-1 studies is provided in turn below.

Table 1. The level-I studies.

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure(s)	Key network findings	Network intervention	Network strategy	Recommendations
Benton 2015 [26]	Scotland, UK	46 Nurse leaders	Secondary and community care	Survey	Communication	Density; average path length; network diameter	Information exchange network was mapped before and after an intervention bringing unconnected nurse leaders together to work on projects. Six months after there was an increase in network density and a reduction in average path length and more ties spanning different areas of work. Participants with low initial connectedness improved their number of ties. Connectedness and closeness improved considerably for those doing projects but not for individuals not involved in projects.	Results of SNA are fed back to participants. A subgroup of participants was then allocated to projects based on their interest in topics, and their low level of pre-existing connection. The projects required participants to communicate and work together to agree on actions to strengthen organizational strategy.	Alteration and induction	Use SNA followed by visual feedback to staff to stimulate positive change in the network. Bring disparate staff together in project teams to facilitate a sustained increase in connectiveness. Include staff with low baseline connectiveness.
van der Eijk 2015 [13]	Netherlands	101 Multidisciplinary healthcare professionals involved with Parkinson care	Secondary and primary	Questionnaire and interview	Knowing each other; professional contacts	Number of connections; density; reciprocity	Participants completed a survey at baseline and one year after the training. Connections increased substantially in both networks from baseline to year one. Positive changes being associated with a central role of neurologists and nurse specialists committed to multidisciplinary care. Perceived team performance did not change.	Multidisciplinary professionals received training in technical and discipline-specific aspects of care, and in communication. Participants are granted access to an online community. There were semi-annual meetings and an annual national conference.	Induction	Provide shared training for multidisciplinary healthcare professionals treating a common disease. Invite participants to be involved in an online community. This may result in increased numbers of connections.

(Continued)

Table 1. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure(s)	Key network findings	Network intervention	Network strategy	Recommendations
Hurtado 2020 [25]	USA	38 (pre) and 55 (post) nurses and nursing assistants	Community hospital	Survey	Advice	7 Network centrality measures	Deployment of champions who had received technical and leadership training was associated with an increase in equipment use, safety compliance and incident reporting. There was a reduction in injuries to staff which was significantly different from 2 control sites.	SNA used to identify influencers in the area of safe patient handling. The top quintile nominations were invited to be champions. They were trained in technical aspects of patient handling and team leadership. They also participated in a number of quality improvement meetings.	Individuals	Use surveys to identify individuals who are already seen as 'go to' people for a particular topic. Identify champions, make them visible, and provide training in both technical skills and knowledge, and leadership. Maintain connection with champions through regular meetings.
Lee 2019 [27]	Malaysia	111 Health care workers	Secondary	Questionnaire	Communication	Geodesic distance; density; reciprocity; degree; closeness; betweenness	Hand hygiene compliance improved by a similarly degree in both peer and manager nominated champion arms. There was an improvement in hand hygiene practice and a preference for top-down leadership structure.	In one study arm staff nominated and ranked peers to become change agents. In the second arm, managers selected champions. Change agents and champions promoted hand hygiene in their local workplace.	Individuals	In order to change workplace behavior, select and train local champions; peer nominated or manager nominated champions have similar impact.

<https://doi.org/10.1371/journal.pone.0282050.t001>

Benton et al.'s [26] research was set in the National Health Service, Scotland. This was a quasi-experimental, pre-post intervention design. Analysis of the communication network of a group of nurse leaders was performed. Forty-six nurse participants from the acute and community setting participated to a baseline survey, which identified 18 participants for the intervention. Participants were selected because SNA data showed they were relatively weakly connected within the network. They were placed into one of three working groups based on their area of expressed expertise or interest. The aim was to influence the existing communication network by encouraging less connected participants to work together. To facilitate this, SNA data from the initial survey was fed back to all participants. The communication network was measured six months after the first data collection. Following involvement in the working groups, the selected 18 individuals showed substantial increase in number of ties. This was evidenced by a rise in connectedness score, which improved from 15.72 to 33.9, and closeness centrality which improved from 8.76 to 13.17. There were also improvement in global network efficiency and density, while the average path length reduced from 1.58 to 1.48. Network visualization showed more connections between professional groups. The Authors suggested that the wider network effects may have been affected by the feedback of the results of the first survey, which made people aware of their own position, and prompted curiosity about how they could change it. It also made people aware of the expertise available in peers. One weakness of the paper is that increase in connectedness among the 18 project participants was based on a survey done six months after the completion of the project groups. Hence, it is unclear whether the impact on network topology would be continued long term.

van der Eijk et al. [13] conducted a parallel group, mixed-methods study in the Netherlands. The study aimed to evaluate the Parkinson network, a nationwide organization with regional networks of health professionals. The study involved 101 multidisciplinary healthcare workers involved with Parkinson's care. Participants, who were based in hospital, nursing home or primary care settings, were selected to take part in a program on the basis of their location and 'motivation' (the latter term is not explicitly defined in the paper). They underwent a training course on multidisciplinary aspects of Parkinson's disease, and were given access to a database of expert therapists in their geographical location. There were also semi-annual meetings and an annual conference. Participants completed a survey on network connections and perceived team performance at baseline. One year later, a subsample was interviewed. There was a substantial increase in the number of 'knowing each other' connections from 1,431 to 2,175 ($p < 0.001$) and in 'professional contact' connections from 664 to 891 ($p < 0.001$). Neurologists and nurse specialists had a central position and were very well connected one year after the program implementation. Overall team performance did not change, but satisfaction with multidisciplinary collaboration increased significantly. There were no data on the impact of network characteristics on either patient outcome measures—such as symptom control or patient satisfaction, or process measures—such as rate of provision of evidence-based elements of care.

Hurtado et al. [25] used social network survey data to identify highly influential co-workers who were recruited as local champions in a safe patient handling education program. The Authors reported that previous studies in this context showed variable short- and long-term impact and that this may be due to a lack of proper methods for selecting workers best suited to exert influence. The study was carried out in critical care areas in one US hospital, and used a survey to collect data on advice seeking for safe patient handling. Individuals showing high centrality in the network were chosen as champions and were trained in safe handling. They were identified to other staff through announcements and wearing of ribbons. The results showed an increase in safety incident reporting, correct equipment use and safety compliance,

as well as reduction in staff injuries. Individual injury profile was significantly different from that of the two control hospitals in the same system.

Lee et al. [27] performed a parallel group study comparing two strategies to influence a behavior, hand hygiene compliance, through the use of local champions. The strategies were deployed on two similar medical wards. SNA showed there were few ties between the wards, suggesting that cross contamination was unlikely to occur. Staff on both wards were asked to nominate and rank peers in terms of their suitability to be hand hygiene champions. In one study arm, champions were selected on this basis. In the other study arm, managers selected champions without reference to the peer ranking. The champions themselves did not know how they had been selected. Trained observers used a validated approach to measure hand hygiene compliance during the study. Compliance increased substantially, from 48% to 66% in the peer selected champion arm, and from 50% to 65% in the manager selected champion arm. There was no statistical difference between the groups.

Level-2 studies

Table 2 below includes the level-2 studies, followed by a descriptive summary.

Seventy-four studies were classified as Level-2. These are studies using SNA solely for descriptive or analytic purposes, without discussing about possible interventions aimed at changing or improving the structure or functioning of the networks. Twenty-three studies were from the USA, 14 from Europe (excluding UK), 14 from LMIC, seven from UK, nine from Australia, five from Canada and two from Japan. Forty-five studies used teams or mixed groups of healthcare professions as participants, 15 papers featured doctors only, 10 papers involved nurses, one study radiologists, one study psychologists, one also involved patients, and one had other types of healthcare professionals.

The majority of the studies ($n = 46$) were set in secondary care settings, followed by community ($n = 9$) and primary care settings ($n = 5$). Eight studies were conducted in mixed secondary and community, and primary and secondary settings. Finally, three studies were set in virtual settings, one in a university hospital, one in a cross sector and one in a nursing home. Twenty-six papers relied on surveys to collect network data, 17 used questionnaires, 10 used logs or administrative data, seven were based on mixed methods, six on observation, four on interviews, two on online platforms or forums, and two on interaction data from sensors.

Ten different types of ties were examined, the commonest being information and knowledge exchange. Nine papers described more than one tie [15, 34, 36, 44, 60, 64, 72, 75, 92]. Twenty-nine different network measures were used to describe the networks at the individual, dyadic, group and whole network levels. Statistical analysis was performed as the only analytical method in 10 studies. Burt et al. [42] is a theoretical paper suggesting different types of questions for name generators. Forty papers (60%) were published between 2010 and 2015, and thirty-four (40%) between 2016 to 1st May 2022.

Level-3 studies

Table 3 below includes the level-3 studies, followed by a descriptive summary.

Twenty-four studies were classified as Level-3. Nine were conducted in the USA, seven in Europe (excluding UK), two in Australia, two in LMIC, two in Canada and two in China. Twelve used teams or mixed groups of healthcare professionals as participants, nine studies used doctors, two had other health professionals and one used healthcare providers and patients. The majority of the studies ($n = 17$) collected data in a secondary care setting, four in the community, one in primary care, one in primary and secondary care, and one in public health.

Table 2. The level-2 studies.

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Kim 2021 [91]	Korea	222 Nursing students	University	Survey	Personal and social support	Indegree; outdegree; betweenness	A high level of subjective happiness is associated with a strong social network. Students with a high level of subjective happiness showed high network centrality. SNA can be used to improve nursing students' happiness by utilizing team-learning social networks within programs.
Haruta 2021 [90]	Japan	52 Multidisciplinary healthcare workers	Secondary	Questionnaire	Advice	Clustering; density; degree; reciprocity; betweenness	Advice seeking network structures differed by topic areas. Nurses had highest centrality for all areas. The effect of feeding back the findings to healthcare professionals may have helped them to reflect on, and act upon their own networks.
Mukinda 2021 [92]	South Africa	42 Managers and healthcare providers involved with maternal, newborn and child health	Primary and secondary	Questionnaire	Communication; social support	Degree; betweenness; density	Governance structures can support collaborative networks to improve cohesion between multidisciplinary teams by integrating missing links to improve information sharing and strengthen teamwork between frontline providers.
Bertoni 2022 [94]	Brazil	133 Multidisciplinary or intensive care unit workers	Secondary	Questionnaire and interviews	Advice	In-degree; closeness; betweenness	Key players are not the same across the four ability-based networks. Thus, if responding, anticipating, learning, and monitoring are core activities that a resilient system displays, different individuals may take the lead on each of those roles. It is possible to investigate the contribution of individual players to resilience from a system perspective.
Smit 2021 [89]	Netherlands	55 Multidisciplinary healthcare professionals	Primary	Survey	Collaboration	Degree; reciprocity	It is feasible to implement an interprofessional collaboration in practice (IPCP) program. Secondary data on the reporting of network metrics showed an increase in the number of contacts among the program participants. After the program, the program and non-program participants gained more collaborative, and diverse inter-professional networks.
Hayward 2021 [93]	Australia	19 Multidisciplinary professionals involved in disability services	Cross sectors	Survey	Advice	Outdegree; indegree; betweenness	Nineteen individuals are identified who occupy positions of either boundary spanning (those linking people and groups) and/or opinion leadership (those that are sought for advice). Boundary spanners meet all criteria while opinion leaders do not.

(Continued)

Table 2. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Durojaiye 2022 [88]	USA	1647 Multidisciplinary pediatric trauma healthcare workers	Secondary	Electronic health records and interviews	Patient sharing	Network graph	Networks dealing with individual trauma cases are different between day and night. Network patterns for collaborative working are different during day versus night shifts.
Tasselli 2015 [85]	Netherlands	118 Hospital professionals (65 nurses and 53 doctors)	Secondary	Survey	Knowledge transfer	Average degree centrality; hierarchy; average betweenness centrality	There are disciplinary cliques for knowledge transfer. Clinical directors facilitate knowledge transfer through their central network position. Junior doctors and nurse managers display both inter-professional and intra-professional centrality positions and are more likely to access valuable knowledge.
Wagter 2012 [61]	Netherlands	108 ICU/MCU staff (senior doctors, nurses, residents and facilitating jobs)	Secondary	Questionnaire	Knowledge sharing	Densities; tie strength; reciprocity	ICU/MCU nurses formed cliques. There are unilaterally directed relations of senior doctors with nurses and patients.
Malik 2014 [35]	Pakistan	48 Primary physicians and 5 district health administrators and line managers	Primary	Interviews and questionnaire	Advice seeking	Network graph	Primary physicians are aware of available expert knowledge, but advice-seeking behavior is dependent upon existence of informal social interaction with the senior specialists.
Patterson 2011 [28]	USA	3 Emergency medical technician teams (EMT) (size of staff: N = 41; N = 67; N = 81)	Secondary	Administrative data	Familiarity (having worked previously together during shifts)	Number of partnerships; means; rates; proportions	On average, an EMT works with 19 different partners over the course of the year and there is significant variation in EMT partner familiarity across agencies. These patterns are considered an indicator of poor emergency medical services outcomes.
Groenen 2017 [55]	Netherlands	214 Healthcare workers from 8 different professions	Secondary and community	Questionnaire	Patient-related contacts	Density; centrality	Almost all professionals in the network can reach other professionals in two steps. Only community-based midwives have connections with all other groups of professionals and represent 51% of all measured connections. The youth health doctors and nurses are mostly positioned on the edge, and are less connected. Obstetricians and community midwives have the highest score for betweenness centrality.

(Continued)

Table 2. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Yuce 2014 [59]	Netherlands	394 Hospital physicians	Secondary	Questionnaire	Advice	Density; average degree centrality	Advice seeking networks among doctors differ for medical and IT related issues. Trainees are just as likely to approach faculty on medical issues as peers, but more likely to approach peers on IT issues. Faculties go to peers for advice in medical practice, but not to trainees for technology-related advice due to the mentor system. Opinion leaders are different for the two domains.
Sibbald 2013 [50]	Canada	6 Multidisciplinary healthcare teams from 2 primary health care team (PHCT) Practices	Primary	Questionnaire and interviews	Information exchange	Density; indegree	Respondents in the sample of PHCTs generally provide research information to only a few individuals on their teams and, overall, only a few individuals are providing the information. Key players in the knowledge uptake and dissemination process are residents, senior physicians, and nurse practitioners.
Benham-Hutchins 2010 [20]	USA	25 Hospital staff and hand-overs (11 to 20 providers over 5 handoffs)	Secondary	Observation Snowball sampling	Communication	Betweenness; closeness; eigenvector; betweenness centralization; hierarchy	Each handoff network exhibits unique communication patterns and coordination. Most participants prefer verbal communication.
Burt 2012 [42]	USA	25 Hospital physicians at quality improvement sites	Secondary	Survey	Different types of ties and name generator questions	Comparison of name generators	Some physicians maintain a social network organized around a specific colleague who perform multiple roles, while others maintain highly differentiated networks. A set of 5 of the 8 name generators used is needed to distinguish the networks of these physicians. Multiple survey questions are needed to elucidate networks of knowledge sharing among physicians.
Shokoohi 2013 [73]	Iran	140 Students (70 clerks, 45 interns and 25 residents) in an educational hospital	Secondary	Questionnaire	Knowledge transfer	Density; indegree; outdegree; reciprocity	Residents are consulted with almost as same as attends on diabetic foot ulcers, hence showing a prominent role in knowledge transfer. The density of clerks-residents and interns-residents is higher than clerks-attends and interns-attends. Indegree centralization in attends-related networks is greater than residents-related networks.

(Continued)

Table 2. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Fuller 2012 [38]	Australia	Two case studies of chronic illness service partnerships (42 partnership staff and 19 informants) in 2 Australian sites	Community	Survey	Communication	Degree; betweenness	Participants in both research groups considered that the network survey accurately described the links between workers related to the exchange of clinical and cultural information, team care relationships, involvement in service management, planning and policy development. Aboriginal workers have a high number of direct links in the exchange of cultural information—suggesting a role of cultural resource—but have fewer direct links in the exchange of clinical information and team care.
Patterson 2013 [39]	USA	103 Clinicians and non-clinician staff in a multidisciplinary Emergency Department (ED) team	Secondary	Survey	Communication	Density; centralization; indegree	There is wide variation in the magnitude of communication cohesion (density) and concentration of communication between clinicians (centralization) by day/night shift and over time. There is also variation in indegree centrality (a measure of power/influence) by day/night shift and over time.
Venkatesh 2011 [86]	USA	1,120 Hospital physicians and other staff (doctors, paraprofessionals, administrative personnel)	Secondary	Survey	Advice	Degree	Ingroup and outgroup ties play a critical role in influencing e-healthcare system use. Further, such use has a positive effect on a variety of quality-of-care metrics that in turn influence patient satisfaction.
Barth 2015 [53]	UK	Pediatric surgery team in 40 pediatric cardiac surgical procedures	Secondary	Observation	Communication	Degree centralization; density; closeness centralization; betweenness centralization; reciprocity	In complex surgical procedures, communication patterns are more decentralized and flatter. In critical transition phases of the procedure, communication is characterized by higher information sharing and participation.
Tsang 2012 [30]	Taiwan	60 Nurses in a dialysis department of a medical centre	Secondary	Survey	Work-related information exchange	Degree; closeness; betweenness	Organizational citizenship behavior (OCB) increases with centrality in both work and friendship networks. Experienced nurses show high centrality in the work networks. In the friendship network, those with high centrality are not necessarily of higher rank in the organization. OCB induced by social ties is satisfactory. It directly increases work satisfaction and alleviates work stress.

(Continued)

Table 2. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Tavakoli Taba 2016 [64]	Australia	31 Breast imaging radiologists	Secondary	Survey	Professional interaction and knowledge sharing	Degree; density; effective size; efficiency; constraint; hierarchy; mean tie strength	There is a positive relationship between diagnostic performance and degree centrality and network size, but a negative relationship with constraint and hierarchy. Overall, the results suggest that radiologists interacting with a closely knit cluster through multiple primary ties—resulting in higher constraints for them—performed worse than radiologists with effective, less constrained (or non-redundant) contacts.
Walton 2010 [21]	Canada	6 Teams in a pediatric ward (doctors, residents and medical students)	Secondary	Observation and questionnaire	Patterns of team interaction	Betweenness	Three different patterns of verbal interaction are observed. In most cases, the attending physician are most talkative and many students and residents spoke infrequently.
Paul 2014 [32]	USA	33 Primary physicians	Community	Survey	Knowledge sharing	Reciprocity; triadic dependence	A physician influential discussion, and a patient-sharing networks are analyzed. Patterns of influential discussions among physicians exhibit triadic dependence. Reduction in reciprocity due to triadic and other higher-order forms of clustering. Geographically proximal physicians are more likely to share patients.
Tighe 2012 [63]	USA	55 Members of Anesthesiology department and 29 patients	Secondary	Service schedule	Communication	Various measures for size and structure of the network, and information flow are used. Many node-level measures are also used	The network exhibits a relatively low density and clustering coefficient, suggesting a low level of redundancy. The high Krackhardt hierarchy score suggests multiple levels of responsibility and supervision between attending, fellow, and resident anesthesiologists. Despite the relatively small size of the core regional anesthesia and perioperative pain medicine) team, its interactions with a large number of services over multiple geographic locations lead to considerable network complexity.
Hinami 2019 [31]	USA	2280 Prescribers of opioid analgesic	Secondary and community	Prescription claim data	Shared benefactors	K-shell centrality	SNA identifies two small, interconnected prescriber communities of high-volume pain management specialists, and three sparsely connected groups of predominantly low-volume primary or emergency medicine clinicians. The sparsely connected clinicians are a risk factor for uncoordinated opioid prescribing.

(Continued)

Table 2. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Long 2014 [52]	Australia	68 Cancer research networks of hospital-based clinicians and university-based researchers	Secondary	Online Survey	Collaboration	Density; components; External-Internal (E-I) index; clustering coefficient	Geographic proximity and past working relationships have significant effects on the choice of current research collaboration partners. Future intended collaborations include a significant number of weak ties and ties based on other members' reputations.
Dauvrin 2017 [72]	Belgium	575 Healthcare professionals working in inpatient and outpatient services	Secondary	Survey	Professional relationships	Degree	At the dyadic level, no significant associations are found between ego cultural competence and alter cultural competence, except for subjective exposure to intercultural situations. No significant associations between centrality and cultural competence, except for subjective exposure to intercultural situations: The most central healthcare professionals are not more culturally competent than less central health professionals.
Altalib 2019 [37]	USA	66 Epilepsy care facilities and 165 providers	Secondary and primary	Secondary data and interviews	Patient sharing	Degree; betweenness; closeness	Across Veterans Affairs Healthcare System (VA) facilities, neurologists are found to be higher on average node degree, betweenness, and closeness centrality measured followed by mental health professionals, then primary. Providers, across disciplines, have higher centrality measures in Epilepsy Centres of Excellence (ECOE) hubs compared to spoke referral facilities and non-affiliated networks. Facilities had a variety of network configurations.
Stewart 2012 [57]	Thailand	46 Pediatric pain practitioners	Secondary	Online discussion forum	Knowledge sharing	Degree; closeness; betweenness; coreness	The network is dominated by one institution and a single profession. There is also evidence of a varied relationship between reading and posting content to the discussion forum. SNA reveals a network with strong communication patterns and users who are central to facilitating communication. SNA also reveals that there is a strong interprofessional and interregional communication, but a dearth of non-nurse participants are identified as a shortcoming.

(Continued)

Table 2. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Blanchet 2013 [62]	Ghana	12 Ghanaian districts; 53 individuals (hospital managers, nurses and district/regional/national health officers, district education officers, community health volunteers, coordinators)	Secondary	Interviews	Coordination and collaboration	Density; distance; degree; betweenness	The departure of an international organization, caused a big shock to the health system, resulting in a change in relationships and power structures within the network. The system shifts from a centralized and dense hierarchical network, to an enclaved network made up of five sub-networks. The sub-networks are less able to respond to shock, circulate information and knowledge across scales or implement solutions. The network is less resilient, yet it responds better to management's need to access information.
Alexander 2015 [70]	USA	12 Certified nursing assistants and registered nurses	Nursing home	Observation	Communication	Network graph	Direct interaction between nurses is higher in the low IT sophistication home and occur in more centralized locations compared to the high IT sophistication home.
Laapotti 2016 [71]	Finland	10 Healthcare professionals with managerial roles, a chair and a secretary	Secondary	Observation	Interactions between team members	Network graphs	The structure of the interaction network reveals that interactions reflect the organizational roles of the participants, as they are focused on the chair.
Lai 2020 [79]	Taiwan	50 Nurses of surgical wards	Secondary	Questionnaire	Friendship	Network graphs; regression	Perceived usefulness, perceived ease of use, and social influence affect behavioral intention to use cloud sphygmomanometer. Besides, perceived ease of use and social influence positively influence perceived usefulness of cloud sphygmomanometer. Peers are helpful in motivating medical staff to use the cloud sphygmomanometer.
Mascia 2014 [40]	Italy	104 Primary physicians and pediatricians	Primary and Secondary	Questionnaire	Knowledge exchange	Outdegree	The number of relationships with hospital colleagues is associated with use of evidence-based medicine.
Shafiei 2018 [82]	Iran	64 Nurses	Secondary	Interviews	Work-related interactions	Degree; closeness; betweenness; eigenvector	Interactions within a department are strong but those between nurses of different departments are not.
Kawamoto 2020 [76]	Japan	76 Intensive Care Unit (ICU) healthcare professionals (HCP)	Secondary	Wearable sensors	Face-to-face interactions	Degree; betweenness; eigenvector	Wearable sociometric sensor badges show nurses have a pivotal role in communication amongst the ICU HCP.

(Continued)

Table 2. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Cavalcante 2018 [77]	Brazil	3 Healthcare professionals (1 doctor and 2 nurses) and their networks' members (19 people)	Mobile Urgent Care Service	Interviews	Work-related interactions	Size; Density	The networks consist of (mutual) relationships that satisfy the demands and needs of service users in an integrated manner while attempting to respect the knowledge and autonomy of each member. Nevertheless, the networks are characterized by poor collaboration ("star" shape) with few transposition points (bridges). This leads to problems in the performance of tasks and mental suffering at work.
Lazzari 2019 [51]	UK	42 Dementia professionals in 3 teams, and 42 patients with Alzheimer's disease	Secondary	Observation	2-mode networks of professionals by services provided	Degree	All professional roles are involved in the case of patients' biological and sociologic personhood. The nurse is the most central figure in the case of biological personhood.
Currie 2012 [87]	UK	36 Pediatric nephrology multidisciplinary teams	Secondary and community	Survey	Knowledge exchange	Degree; betweenness; brokerage roles; density	Knowledge-brokering roles are influenced by professional hierarchy, particularly in the case of clinical knowledge and even more so with medical knowledge.
Chung 2014 [84]	Australia	107 General Practitioners	Primary	Questionnaire	Advice	Density; inclusiveness; components	Considering the GP-patient encounter as a complex system, the interactions between the GP and their personal network of peers give rise to "aggregate complexity," which in turn influences the GP's decisions about patient treatment. GPs in simple profiles (i.e. with low components and interactions) in contrast to those in nonsimple profiles, indicate a higher responsibility for the decisions they make in medical care.
Yuan 2020 [49]	USA	207 Nurses in 6 clinical units in an academic hospital	Secondary	Survey	Advice	Mean peer belief (ego-network analysis)	Although mean beliefs across the entire peer network have no effect on individuals' system use, shared peer beliefs were associated with nurses' increased use of the IT system. Reinforcement by the social network appears to influence whether individuals' own beliefs translate into system use, providing further empirical support that social networks play an important role in the implementation of health information technology.

(Continued)

Table 2. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Uddin 2013 [69]	Australia	85 Physicians networks	Secondary	Health insurance claim dataset	Collaboration	Density Degree; betweenness centralization Exponential random graph models (ERGMs)	Collaboration structures among physicians affect hospitalization cost and hospital readmission rate.
Benton 2014 [48]	Scotland, UK	27 Senior nurses	Virtual	Survey	Communication	Degree; betweenness; eigenvector; density; average path length; network diameter	The majority of nurse leader who participated in the Global Nursing Leadership Institute 2013 Programme are poorly connected in social media, i.e., they have low indegree and outdegree scores. Existing connections are centered on geographic proximity and participation in regional and global bodies.
Mundt 2015 [54]	USA	155 Primary health care professionals from 31 teams at 6 primary care clinics	Community	Survey	Communication	Density; centralization	Teams with dense interactions are associated with fewer hospital days and lower medical care costs. Conversely, teams with interactions revolving around a few central individuals are associated with increased hospital days and greater costs.
Quinlan 2013 [74]	Canada	49 Nurse Practitioners in primary healthcare teams	Primary	Survey	Knowledge transfer	Within-team reciprocity; within-team degree centrality	Mutual understanding increases from one clinical decision to another in some teams and decreases in others. The new Nurse Practitioners play a crucial role in facilitating mutual understanding and knowledge exchange in the newly created multidisciplinary teams. A well-functioning team has effective intrateam knowledge exchange.
Li 2016 [58]	Netherlands	621 Healthcare Professionals (users) and 723 threads over 40 forums	Virtual	Online discussion forum	2-mode network of forum users by discussion threads	Density; centralization; diameter; average path length; SAOMs	The participation level in the discussion within the online community is low in general. A change of lead contributor results in a change in learning interaction and network structure. Health professionals are reluctant to share knowledge and collaborate in groups, but are interested in building personal learning networks or simply seeking information.
Sullivan 2019 [65]	UK	39 Trainee doctors in an acute medical unit	Secondary	Survey	Advice	Degree; betweenness; density	Information and influence relating to different aspects of practice have different patterns of spread within teams of trainee doctors. Influencers in clinical teams have particular characteristics, and this knowledge could guide leaders and teachers.

(Continued)

Table 2. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Zappa 2011 [16]	Italy	711 Physicians	Secondary	Survey/ Questionnaire	Knowledge sharing	ERGMs	Knowledge flows informally in mutual information-seeking relationships. Physicians tend to cluster in small groups of proximate and similar peers. The propensity to share knowledge is affected by individual-specific characteristics.
Aylward 2012 [47]	USA	286 Pediatric Psychologists	Primary and Secondary	Survey	Mentoring	Density; indegree; outdegree; closeness; betweenness; average geodesic distance	The field of pediatric psychology is interconnected with professionals learning from multiple mentors in multiple settings. The average “degrees of separation” between individuals in the network is 5.30.
Benammi 2019 [78]	Morocco	58 Members of an Acute Care Unit (ACU) in a university hospital	Secondary	Survey	Communication	Density; degree and betweenness centralization; degree and betweenness centrality	ACU network shows a moderate degree centralization, and lower betweenness centralization. The team is connected by well-positioned members to support inter-team communication, and is dominated by a number of gatekeepers, with low degree of communication among different function team members.
Bachand 2018 [66]	USA	8338 Women with breast cancer in 157 physician peer groups (made up of 16,171 physicians)	Secondary	Surveillance, epidemiology, and end results-Medicare data	Patient-sharing	Ingroup density; transitivity	Surgical delays vary substantially across physician peer groups, and are associated with provider density and patient racial composition. Women in physician peer groups with the highest provider density are less likely to receive delayed surgery.
Bae 2017 [33]	England, UK	54 Nurses in an acute care hospital unit	Secondary	Survey	Mutual support	Degree; closeness; betweenness; eigenvector density; shortest path; reciprocity; transitivity	Providers of mutual support claim to give their peers more help than these peers gave them credit for. Those who work overtime provide more mutual support.
Fong 2017 [43]	Taiwan	100 Multidisciplinary staff members in 3 Intensive Care Unit (ICU) in an academic teaching hospital	Secondary	Questionnaire	Communication	Cluster analysis (k-means)	Distinct patterns and categories of influencers (well-rounded, relational, and knowledge-based) are identified using a clustering approach. Knowledge of how influence is distributed across the care team could lead to a better planning of change initiatives.
Creswick 2010 [19]	Australia	45 Health professionals in a renal ward	Secondary	Questionnaire	Advice seeking	Geodesic distance; density; average strength of ties; reciprocity; degree; betweenness	On average, there is little interaction between each of the staff members in the medication advice-seeking network, with even less interaction between staff from different professional groups. Nurses are mainly located on one side of the network and doctors on the other. However, the pharmacist is quite central in the medication advice seeking network as are some senior nurses and a junior doctor.

(Continued)

Table 2. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Dauvrin 2015 [60]	Belgium	507 Healthcare professionals	Secondary and primary	Questionnaire	Problem-solving, advice-seeking, and socialization	Indegree	Cultural competence of the healthcare staff is associated with the cultural competence of the leaders. The leadership effect varied with the degree of cultural competence of the leaders.
Wong 2015 [36]	USA	98 Pediatric Intensive Care Unit staff	Secondary	Survey	Information seeking, social influence and social support	Degree; density	Amongst the 3 networks, there are no weakly connected groups. Few individuals report no links to a colleague. The number of links among colleagues is greatest for the information seeking network, followed by social influence, and social support. Five individuals, three of whom have formal leadership roles, are amongst the 10 most influential team members in all 3 networks.
Hurtado 2018 [67]	USA	38 Patient care workers	Community hospital	Survey	Advice seeking	Degree; reciprocity	There is a positive correlation between identifying more peers for safe patient handling advice and using equipment more frequently. Nurses with more reciprocal advice seeking nominations use safe patient handling equipment more frequently. However, nurses consulted more do not use equipment more frequently than nurses with fewer nominations.
van Beek 2013 [29]	Netherlands	391 Nursing staff from 37 long-term care dementia units	Community	Questionnaire	Communication	In-group density	In units with more networks between nursing staff and relatives of residents, staff treated residents with more respect and were more at ease with residents. Social networks were also positively related to staff's organizational identification which, in turn, related to their work motivation and their behavior towards residents.
Boyer 2010 [23]	France	104 Healthcare professionals in a hospital	Secondary	Questionnaire	Information sharing	Ingroup centrality; prestige; clique indicators	Centrality, prestige and clique indicators are highly correlated. Physicians have the highest scores for the three indicators. Older age is found to be associated with higher centrality and clique scores.
Anderson 2011 [56]	USA	Operating room staff (n = 733 interdisciplinary members) of 2 surgical specialties	Secondary	Staffing data on surgical cases in the 29 operating rooms	Individual affiliation to surgical cases	Degree; closeness; betweenness; eigenvector; core/periphery	Both surgical services show a core/periphery network structure. Team coreness is associated with the length of the case. Procedure start time predicts the team coreness measure, with cases starting later in the day less likely to be staffed with a high core team. Registered nurses constitute the majority of core interdisciplinary team members in both groups.

(Continued)

Table 2. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Brewer 2020 [75]	USA	268 Nursing staff in 24 Patient Care Unit (PCUs)	Secondary	Web-based questionnaire	Information sharing and advice seeking	Average distance; betweenness; clique count; clustering; density; diffusion; eigenvector; fragmentation; hierarchy; isolates; size; degree	In clinical workplaces with high day-to-day staff variation, several network characteristics remain stable over time. Hierarchy, fragmentation and cliques are unstable.
Lower 2010 [15]	Australia	13 Multi-disciplinary teams in hearing services	Community	Questionnaire and interviews	Information exchange; referrals; working relationships	Degree; average number of ties	Nurse audiometrists, WorkCover and agricultural retailers have the lead role in disseminating information on hearing health within the network. For client referrals the nurse audiometrists, private audiology services, general practitioners, ear, nose and throat specialists and industry groups play the major roles.
Quinlan 2010 [17]	Canada	29 Nurse practitioners in primary-care teams	Community	Survey	Mutual understanding	Within-team density; flow-betweenness centralization	In two teams mutual understanding increases with time. In the other two teams, it decreases. As the overall mutual understanding within the team decreases, the facilitation of mutual understanding becomes more centralized among few team members; conversely, as mutual understanding increases, the facilitation becomes more equally distributed. The inverse relationship exists in all teams, except in team.
Edge 2019 [45]	UK	138 Foundation doctors in one NHS trust	Secondary	Observational study	Physical contact	Degree; density; density by groups; assortativity	Direct network links to vaccinated colleagues increase an individual's likelihood of being vaccinated.
Espinoza 2018 [44]	Chile	53 Inter-professional teams (409 professionals) at a university hospital	Secondary	Questionnaire and interview	Advice and personal support	Density; isolates; centrality; Within-group cohesion	For the work advice network, when a team structures itself around one professional, this allows its members to approach and be approached easily and facilitates information exchange. Teams with the least satisfaction reveal a fragmented structure with members organized as subgroups. The organization of social support networks is even more fragmented, with half of them being isolated from the rest of the team.
Crockett 2018 [80]	Canada	22 Healthcare professionals in 18 general Emergency Departments	Secondary	Interviews	Information seeking	Content analysis	Health care professionals sought information both formally and informally, by using guidelines, talking to colleagues, and attending pediatric related training sessions. Network structure and processes were found to increase connections, support practice change, and promote standards of care.

(Continued)

Table 2. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure (s)	Key network findings
Pomare 2019 [34]	Australia	23 and 27 Clinical and non-clinical staff members in 2 headspace centres	Youth mental health service	Survey	Collaboration, advice, problem solving	Degree; sub-group cohesion; density; centralization	Staff of headspace (clinical and non-clinical) show a tendency to collaborate with colleagues outside of their professional group, compared to within. Networks are well connected when staff collaborate in routine work and when faced with uncertainty in decision-making. There are fewer interactions during times of role uncertainty. The headspace centre that had been in operation for longer show greater indicators of cohesiveness.
Choudhury 2018 [81]	USA	3 Large-sized integrated delivery networks; 14 hospitals; 288 physicians; 353 prescriptions	Secondary	Medical prescriptions and affiliations datasets	Affiliation	Diffusion models	Physicians affiliated to same hospital and integrated delivery network contribute highly in the diffusion process. The weighted edge approach is better able to explain diffusion of influence in terms of prescribing patterns.
Palazzolo 2011 [68]	USA	3 Multidivisional healthcare teams (n = 126 individuals) in 1 hospital	Secondary	Email archives	Communication	Betweenness; contribution index; group betweenness; core/periphery; density; structural holes; connectivity	SNA of email communications of three teams caring for patients with different complex long-term conditions reveal distinct patterns and structures. Team metrics varied over time. Teams' network characteristics may explain their functioning.
Hornbeck 2012 [46]	USA	Healthcare workers (HCW) and patients in 1 Medical intensive care unit	Secondary	Mote-based sensor network	Physical contact	Agent-based simulation	Electronic sensor derived data on HCW interactions with other HCW's and patients reveal that a small number of HCWs were responsible for a large number of interactions.
Shoham 2015 [41]	USA	69 Co-workers listed by 48 clinical team members in a burn intensive care unit	Secondary	Questionnaire	Communication	Degree; betweenness; density	The analysis revealed three distinct sets of team members caring for two sets of patients. The five clinical team members most central to the network included three physicians, a social worker, and a dietitian.
Zappa 2014 [83]	Italy	106 Oncologists	Virtual community	Emails	Cooperation	SAOMs	Emergent network effectively represented by a small number of local rules, i.e., actors' behaviors of counterpart's selection in their neighborhood.

<https://doi.org/10.1371/journal.pone.0282050.t002>

Ten studies used questionnaires to collect data, three relied on mixed methods, four used surveys, three interviews, two collected interaction data from sensors, one used direct observation, and one an online platform or forum. Seven different types of ties were analyzed across studies. Two studies analyzed more than one tie [108, 113, 114]. Twelve different network measures were used to describe or analyze networks at the individual, dyadic, group and whole network levels. Statistical analysis relying on ERGMs and MRQAP were used five times.

Table 3. The level-3 studies.

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure(s)	Key network findings	Recommendations	Network strategy
Xu 2021 [115]	China	5247 Healthcare Workers	Secondary	Survey	Discussion	Density; degree	A vaccination consulting network of 1817 members is reconstructed. The network shows low density. Twenty-two influential members are identified. Lack of discussion is associated with vaccine hesitancy. Department leads are particularly influential as promoters of vaccination.	Use influential individuals as role models to encourage vaccine uptake.	Individuals
Jippes 2010 [18]	Netherlands	81 Gynecologists and 63 residents in O&G and Pediatrics	Secondary	Questionnaire and interviews	Communication	Degree; closeness; betweenness	Social connections are more important than training for uptake of a new practice. A strong association is found between closeness centrality and adoptive behavior, and a moderate effect of degree centrality.	Incorporate individuals who have both strong and weak ties in 'teacher-the-teacher' courses.	Individuals
Mascia 2018 [111]	Italy	97 Pediatricians in 2 Local Health Authorities (LHAs)	Community	Questionnaire	Advice	ERGMs	In both LHAs, physicians tend to reciprocate advice ties; there is considerable clustering in advice-seeking.	Create new opportunities for knowledge exchange, such as taskforces or training programs.	Induction
Llupia 2016 [97]	Spain	235 Healthcare workers in 1 hospital	Secondary	Interviews	Information exchange	ERGMs	Similarity in vaccination behavior does not play a significant role in the probability of being connected to another healthcare worker.	Use SNA to guide the design, implementation, evaluation of a health promotion campaign. For example, messages could be tailored by professional category or strategy could be implemented to foster communication among different professional categories.	Segmentation and induction

(Continued)

Table 3. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure(s)	Key network findings	Recommendations	Network strategy
Meltzer 2010 [102]	USA	56 Physicians attending on the general medical services in 1 hospital	Secondary	Questionnaire	Communication	Degree; Net degree of team; betweenness; density	Connections of team members outside the team are important for dissemination of information or influence. Connections of team members inside the team are important for within-team coordination, knowledge sharing and communication.	Use SNA to decide whom to select for a quality improvement team, and how to structure the team. When influence through direct social interaction is important, choose individuals who can reach the largest number of persons outside the team. The use of degree alone to select team members may produce many redundant ties.	Individuals
Polgreen 2010 [103]	USA	148 Multidisciplinary healthcare workers in 1 hospital	Secondary	Observational data and simulation	Physical contact	Number of contacts	Preferentially vaccinating healthcare workers in more connected job categories yield a lower attack rate and fewer infections in a simulation.	Identifying workers with many contacts might aid targeting vaccinations to optimize impact on flu spread.	Individuals
Mascia 2011 [24]	Italy	297 Hospital physicians in 6 hospitals	Secondary	Questionnaire	Advice	MRQAP	Physicians reporting similar attitudes toward evidence-based medicine (EBM) are more likely to exchange information and advice.	Foster heterophily when multidisciplinary cooperation is required. Identify groups exhibiting desired attitudes and behaviors. Adopt organizational arrangements, processes and informal meetings to foster collaboration.	Induction and segmentation
Sykes 2011 [22]	USA	151 Hospital physician in 1 hospital	Secondary	Survey	Advice	First degree centrality; second degree centrality	Both first-degree (direct) and second-degree (indirect) centrality negatively influence electronic medical records (EMR) system use. Physicians with more connections are less likely to be early users of EMR.	Be aware that resistance to EMR systems is greater among physicians with high centrality who should then be the target of resources to reduce such resistance.	Individuals

(Continued)

Table 3. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure(s)	Key network findings	Recommendations	Network strategy
Pinelli 2015 [105]	USA	72 Multidisciplinary healthcare professionals in 1 hospital	Secondary	Interviews	Communication	Size; density; strength of tie; betweenness	Most communication is synchronous. Most communication events occur between the primary nurse and patient, and the care coordinator and primary nurse.	Improvements in discharges are possible by reorganizing systems to optimize communication. SNA could offer a cost-effective way to improve patient care provision.	Alteration
Mascia 2015 [107]	Italy	297 Hospital physicians in a Local Health Authority (LHA)	Secondary	Questionnaire	Information exchange	MRQAP	Institutional and professional homophily affect inter-physician networks. Professional homophily is more relevant than institutional affiliation for collaborative ties.	Foster collaboration across heterogeneous groups of physicians from different specializations.	Induction
Shoham 2016 [99]	USA	71 Multidisciplinary healthcare professionals in a hospital burn unit	Secondary	Survey	Discussion	Density; degree; ERGMs	Members of all roles are involved in a higher percentage of inter- than intra-professional ties. Physicians are most central to the network. Nurses are significantly more likely to connect with other nurses.	Consider purposefully developing the role of nurses within the team.	Segmentation
Gorley 2016 [106]	Canada	227 Participants in a BC Sepsis network	Secondary	Questionnaire and interviews	Knowledge sharing	Density; Centrality	Eleven participants stand out as hubs (high degree centrality). These individuals have many connections with people who trust them.	When launching a new network or strengthening an existing network for quality improvement, several recommendations are offered (e.g. to seek and include distributed leaders in the network).	Individuals and Induction

(Continued)

Table 3. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure(s)	Key network findings	Recommendations	Network strategy
Mascia 2013 [100]	Italy	297 Hospital physicians in 6 hospitals; 1 Local health unit	Secondary	Questionnaire	Advice	Coreness; network authority	The overall network shows a core-periphery structure. There is a negative association between physicians' attitudes toward evidence-based medicine (EBM) and the coreness they exhibited in the professional network. Network centrality indicators confirm a negative association between physicians' propensity to use EBM and their structural importance in the professional network.	Policy makers can foster collaboration across staff with different propensities to use EBM by relying on organizational arrangements, informal meetings, and use of medical leaders to persuade other professionals to collaborate more with EBM user.	Individuals and Induction
Creswick 2015 [101]	Australia	101 Hospital staff members in 1 teaching hospital	Secondary	Questionnaire	Advice	Density; reciprocation; indegree	Medication advice-seeking networks among staff on hospital wards are sparse, information sharing across professional groups is modest, and rates of reciprocation of advice is low. Senior physicians are weakly integrated into medication advice networks; pharmacists and junior physicians play central roles.	Policies to advance the advice-giving networks between senior and junior physicians may improve medication safety as one ward with stronger networks had lower prescribing error rate.	Segmentation
Marques-Sanchez 2018 [98]	Spain	196 Multidisciplinary healthcare professionals	Secondary and primary	Questionnaire	Internal and external advice	Outdegree (internal and external ties)	For physicians, external ties improve the performance at an individual and team level, yet external ties are not relevant for nurses' work performance.	Use SNA to facilitate healthcare professionals sharing information within and across organizations.	Alteration

(Continued)

Table 3. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure(s)	Key network findings	Recommendations	Network strategy
Kothari 2014 [114]	USA	13 Public health practitioners	Community	Questionnaire and interviews	Interaction, support, and professional relationships	Cliques; degree; closeness; betweenness	Participants' report on their experience with SNA.	Use SNA as a reflective practice tool for professionals to assess their networks and strengthen collaborations. Assess team arrangements to identify the absence of key players or to recognize critical gaps in communication links that are necessary to work collaboratively.	Alteration and Individuals
Mundt 2019 [96]	USA	143 Healthcare professionals at 5 primary clinics	Primary	Survey	Communication	Core/periphery	Clinic employees in the core of the communication network have significantly greater job satisfaction than those who are on the periphery.	To increase clinicians' job satisfaction, foster face-to-face communication among all team members.	Alteration
Assegaai 2019 [110]	South Africa	Community health workers (CHW) (n = 37), ward-based outreach team (WBOT) leaders (N = 3), primary healthcare facility (PHC) managers (N = 5) and local area managers (N = 2)	Community	Questionnaire	Interaction (supportive supervision)	Network graphs (indegree; density)	The supportive supervision system revolves around team leaders, who are nurse cadres and who ensure internal cohesion and support among WBOT members. The network patterns also show the extent of peer support between CHWs and WBOTs.	Relationships within teams work better than those between teams. Use SNA to identify relationships that could be strengthened.	Alteration
Tighe 2014 [112]	USA	A single day operating room (OR) schedule encompassing 32 anesthetizing sites	Secondary	Simulation and interviews	Interaction	Degree; betweenness; eigenvector	The OR is a scale-free network with small-world characteristics. There are differences in degree centrality between nurses and anesthesiologists and surgeons. Attendings have greater degree centrality than residents.	Use SNA to improve communication within ORs (e.g., by protecting a few highly-connected individuals; by placing senior staff into roles based on communication volumes).	Segmentation

(Continued)

Table 3. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure(s)	Key network findings	Recommendations	Network strategy
Sykes 2015 [109]	Australia	171 Operating room (OR) staff members in 4 surgical teams in 1 hospital	Secondary	Electronic database	Interaction	Network graphs	Eighteen staff members are regularly shared across teams, including 12 nurses, five anesthetists, and one registrar. Weak but significant correlations is found between the number of staff, procedure start time, length of procedure, and patient acuity.	Use SNA to identify change champions who can support initiatives across multiple teams.	Individuals
Hossain 2012 [104]	USA	204 Outpatient departments (OPDs) and 458 emergency departments (EDs)	Secondary	National survey	Coordination	Degree; Density; centralization	The nurse is the actor with highest degree, followed by physician and lab technician. There is a significant relationship between degree and performance of coordination.	Use SNA to understand the possible causes of inefficient coordination performance and coordination quality resulting in access blocks.	Alteration
Li 2020 [108]	China	102 Hospital doctors	Secondary	Online forum	Communication and information exchange	Density; degree centralization; geodesic distance; centrality; reciprocity	Doctors are more closely connected, and information is easily spread. Doctors with higher professional titles show high levels of reciprocity. They are more likely to influence the behavior of other doctors.	Introduce clinical educational meetings to increase the frequency of doctor interaction at different levels.	Induction
Yousefi Nooraie 2017 [95]	Canada	14 Multidisciplinary public health staff	Public health centers	Interviews	Information exchange	Indegree	Information seeking networks evolve towards more centralized structures. Staff who are already central at baseline gain even more centrality.	Use SNA to support and inform the design, process and evaluation of the evidence informed decision-making training interventions.	Induction

(Continued)

Table 3. (Continued)

Ref	Country	Participants	Setting	Data collection	Type of tie	Network measure(s)	Key network findings	Recommendations	Network strategy
Cannavacciuolo 2017 [113]	Romania	28 Multidisciplinary rehabilitation unit staff.	Rehab	Questionnaire	Advice and knowledge exchange	Centrality; frequency of interactions; in-group/out-group interactions	Knowledge is shared in a centralized network characterized by the presence of a few hubs and some marginal actors. The team members consult with a high number of external experts but these sources tend to belong to personal networks and are not shared. Interpersonal knowledge exchange is mostly vertical than lateral.	Redesign the team network to improve the efficiency and effectiveness of knowledge sharing. The re-design interventions concern three main features of knowledge network: "knowledge centralization," "over-reliance on external experts", and "unshared knowledge tools and sources." Different strategies are discussed.	Alteration

<https://doi.org/10.1371/journal.pone.0282050.t003>

The four types of network interventions were mentioned as recommended strategies to be designed and implemented in order to improve the overall structure and functioning of the networks. Nine studies recommended to use 'individuals' [18, 22, 100, 102, 103, 106, 109, 114, 115], eight studies recommended 'induction' [24, 95, 97, 100, 106–108, 111], seven studies discussed possible 'alteration' strategies [96, 98, 104, 105, 110, 113, 114], and four recommended 'segmentation' [97, 99, 101, 112]. Four studies recommended more than one strategy [97, 100, 106, 114]. Thirteen papers were published between 2010 to 2015, and 11 between 2016 to 1st May 2022.

Discussion

We updated previous reviews by including papers published since 2010 that have used SNA to investigate networks among healthcare professionals. Our search strategy included a wide range of databases and placed no restrictions on professional groups, healthcare setting, country, or study design. We found 102 papers that used SNA to examine networks of healthcare professionals. We confirmed the findings of prior systematic reviews: The majority of published studies were descriptive, with only four papers discussing the outcomes of an SNA-based intervention. We defined network intervention as a set of actions aimed at modifying the main elements of a network system (i.e., nodes and relations) so as to generate behavior change and improve system performance. The main idea behind network intervention is that if networks affect outcomes of interest, change in network structure could lead to change in relevant outcomes.

A possible explanation for the limited number of studies on network interventions concerns the practical difficulties in designing and implementing network-based interventions in general, and in healthcare contexts more specifically. Valente et al. [4] discuss the main challenges associated with network interventions in the domains of public health and medicine. In what follows, we will briefly describe the main challenges that we believe arise when an intervention is designed and implemented within an organizational context, such as a hospital or other healthcare organizations. Healthcare organizations present additional challenges over and above those identified by Valente et al. [4] for the public health domain. We organize our discussion by using the four-stage model of program implementation suggested by Valente et al. [4].

Exploration

The first stage involves the assessment of a community in terms of needs, vision and opportunity for change [4]. In practice, this implies identifying: (i) a well-defined network (i.e., community boundaries); (ii) the relations among community members (i.e., social capital); (iii) the specific interests of various stakeholders, and (iv) the behavior under investigation. A number of specific challenges may arise at this stage when social network research is conducted within organizations [116]. First, network identification. This may be facilitated by the natural boundaries that organizations provide for the network of interest. Problems typically arise in collecting the non-anonymous data needed for network research. The management of the organization (which is often also the commissioner of the research) may provide partial commitment or discontinued support to the research, or even restricted access to data. Access to network and other types of data may also be problematic due to the specific nature of the population under investigation. Intervention programs within healthcare organizations are likely to involve multiple professional groups (e.g., hospital administrators, medical doctors, nurses, etc.) whose interdependencies may be difficult to manage or predict thoroughly *ex ante*. The actual use of output data from hospital administrators, participants' protection of ethical

rights, as well as the existence of ethical codes for professionals are all factors that may make data collection within healthcare organizations particularly challenging [117]. A solution to this problem may be a clear identification and communication of the goals and objectives of the research. The four studies that we identified as reporting the results of a network intervention (level-1), or those recommending a follow-up intervention in their conclusion section (level-2), mainly focused on improving specific structural features of the networks. Of the four level-1 studies, only two measured the impact of network intervention on health-related outcomes [25, 27]. The reason for this may lie in the difficulty of envisioning clear-cut causal links between behaviors at one level (e.g., health professionals) and outcomes at another level (e.g., patients). More direct evidence of measurable outcomes of network interventions at the patient or organizational level is needed. Finally, ethical challenges should also be considered at this stage. Cronin et al. [118] and Borgatti and Molina [119] offer explicit guidance on how to deal with specific ethical issues such as protecting anonymity, presenting output data in aggregated form, and offering participants multiple opportunities for opting-out.

Adoption

The second stage involves the creation and adoption of an intervention program to address a behavioral problem [4]. The use of network analysis is particularly helpful at this stage, as it provides valuable information that can be used to tailor an intervention to the specific needs of the population under investigation. High response rates and lack of missing data are crucial as they allow to design an intervention based on more complete information. The identification of opinion leaders within a network who may act as change agents has been used in a large number of studies. Also, network analysis may be useful at this stage to identify other roles or positions, cohesive subgroups, or important cleavages within a network structure. Within an organizational setting, the existence of a formal reporting structure is particularly relevant in that it provides additional information on power structures and formal roles that can also be leveraged in a network-based intervention.

Implementation

The third stage involves implementing the program with adherence and competence [4]. Within healthcare organizations, pressures to improve outcomes (e.g., clinical, operational, financial and managerial) are frequently generated by policy changes that produce top-down initiatives proposed by senior management and implemented through the involvement of various organizational change agents such as medical doctors, hospital administrators and, occasionally, technical and support staff. Research has recognized that the success of change initiatives hinges on the ability of change agents to overcome potential resistance from other organizational members, and encourage them to adopt or develop new practices [120]. In professional organizations, such as healthcare organizations, the coexistence of many professional groups with strong identity and role boundaries may represent the biggest obstacle to organizational change. Furthermore, not all change initiatives are equivalent, and recent research has pointed to the need of establishing the extent to which a change initiative diverges from the institutional status quo in order to better identify factors enabling adoption [120]. Other than resistance to, and extent of, change, challenges that may arise at this stage include availability of resources needed to implement a change program, lack of evidence of successful research designs to use in non-experimental, organizational settings, and lack of clarity about outcome variables to be monitored during the implementation stage.

Sustainment

The fourth, and last stage involves checking that the program continues to be implemented as intended over time, and is continuing to exert the anticipated effects [4]. The main challenge at this stage concerns the slow-moving nature of network and organizational variables, compounded by the often-far too high turnover rates within organizational units. This could make particularly difficult predicting with a reasonable level of certainty how long a social structure would take to affect a behavior, or an outcome of interest. As this usually takes time, problems may arise that are related to changes in the composition of a network structure, which should ideally remain unchanged for the duration of an intervention program. In non-experimental, naturalistic settings this is unlikely to occur. Research has also shown that changes in the composition of a network structure led to changes in the attitudes and behaviors of those who remain in the organization [121].

We have not offered specific solutions to the various issues highlighted above. Rather, our aim was to shed light on the main challenges of implementing a change initiative within an organizational setting. A possible solution to some of the challenges associated with implementing an intervention and measuring its effects over time is the adoption of a simulation-based analytic approach. This approach involves data collected on an existing network to simulate a number of alternative scenarios resulting from altering specific characteristics of the nodes and ties within a network. An example of application of a simulation-based approach to a longitudinal network dataset can be found in Schaefer et al. [122]. The authors use the results of Stochastic Actor-Oriented Models to simulate the coevolution of friendship ties and smoking behavior under potential intervention scenarios. Currently available statistical models for network data have the advantage of being particularly well-suited for simulation analyses. This is an approach that we believe may provide realistic and interpretable evidence of the possible outcomes of a change initiative, and may justify the long-term resource commitment that network-based interventions usually require.

While a number of studies are available that describe network structure, it is important to consider that research informing on how to make positive changes in networks is likely to be closer to having practical impact. There is an urgent need for more research into which healthcare network interventions work in different contexts and how they can be best designed and employed. Similarly pressing is a need for further work to identify experimental design options that are more effective at identifying and maximizing control over relevant variables and outcomes, and that are more efficient in terms of time and resource needed. We may conclude that this is an important opportunity for the field to coalesce on terminology, measures, and applications, after establishing priority areas for researchers in how to do so to advance work on the application of SNA to the design, dissemination, implementation and sustainability of behavior change interventions.

Limitations

We used a comprehensive broad approach to searching but may have missed some research results such as unpublished conference proceedings, papers not available in English language, negative findings or studies that did not complete and were not submitted, and grey literature.

Conclusion

Studies of network intervention remain scant and devoid of implications for the impact of intervention initiatives on patient care. There is a need for evidence on which kinds of network interventions work, in which contexts, and under what conditions—or for whom. It is possible to measure the effect of an intervention on network effectiveness, for example, by measuring

the number of new links or increased volume of communication. However implicitly, this approach assumes a causal link between inter-professional communication and patient benefits. The complexity of healthcare, and the ubiquitous nature of barriers to best practice, implies that this is often a conjecture too far, and a more direct evidence of patient benefit should be preferred. The most important test of the effectiveness of network intervention would be assessing its impact on patient level outcomes, or, when this is difficult to determine, on the delivery of processes of care that are supported by good evidence.

Supporting information

S1 Table. Database results.

(PDF)

S2 Table. Types of network ties.

(PDF)

S3 Table. Network measures.

(PDF)

S1 File. Search strategy.

(PDF)

S2 File. PRISMA-ScR checklist.

(PDF)

Acknowledgments

Disclaimer. The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

Author Contributions

Conceptualization: Ameneh Ghazal Saatchi, Francesca Pallotti, Paul Sullivan.

Data curation: Ameneh Ghazal Saatchi.

Formal analysis: Ameneh Ghazal Saatchi, Francesca Pallotti, Paul Sullivan.

Funding acquisition: Ameneh Ghazal Saatchi.

Investigation: Ameneh Ghazal Saatchi, Francesca Pallotti, Paul Sullivan.

Methodology: Ameneh Ghazal Saatchi, Francesca Pallotti, Paul Sullivan.

Project administration: Ameneh Ghazal Saatchi.

Validation: Ameneh Ghazal Saatchi, Francesca Pallotti, Paul Sullivan.

Writing – original draft: Ameneh Ghazal Saatchi, Francesca Pallotti, Paul Sullivan.

Writing – review & editing: Ameneh Ghazal Saatchi, Francesca Pallotti, Paul Sullivan.

References

1. Baker A. Crossing the Quality Chasm: A New Health System for the 21st Century. *BMJ*. 2001 Nov 17; 323:1192.
2. Schneider EC, Sarnak DO, Squires D, Shah A, Doty. Mirror, Mirror 2017: International comparison reflects flaws and opportunities for better U.S. health. *Care*. Commonwealth Fund. 2017. Available from: <https://doi.org/10.26099/0mh5-a632>

3. Freeman LC. The development of social network analysis. Vol. 1, A Study in the Sociology of Science. New York: Empirical Press; 2004.
4. Valente TW, Palinkas LA, Czaja S, Chu KH, Brown CH. Social network analysis for program implementation. *PLoS One*. 2015; 10(6):e0131712. <https://doi.org/10.1371/journal.pone.0131712> PMID: 26110842
5. Snijders TAB. Models for Longitudinal Network Data. *Models and Methods in Social Network Analysis*. 2005; 11:215–247.
6. Hunter RF, McAneney H, Davis M, Tully MA, Valente TW, Kee F. “Hidden” social networks in behavior change interventions. *Am J Public Health*. 2015; 105:513–516. <https://doi.org/10.2105/AJPH.2014.302399> PMID: 25602895
7. Valente TW. Network interventions. *Science*. 2012; 337:49–53. <https://doi.org/10.1126/science.1217330> PMID: 22767921
8. Chambers D, Wilson P, Thompson C, Harden M. Social network analysis in healthcare settings: a systematic scoping review. *PLoS One*. 2012; 7:e41911. <https://doi.org/10.1371/journal.pone.0041911> PMID: 22870261
9. Cunningham FC, Ranmuthugala G, Plumb J, Georgiou A, Marks D, Westbrook JI, et al. Social-Professional Networks of Health Professionals: A Systematic Review. Centre for Clinical Governance Research in Health Australian Institute of Health Innovation. 2010. Available from: <https://www.yumpu.com/en/document/read/28077911/social-professional-networks-of-health-professionals-australian->
10. Bae SH, Nikolaev A, Seo JY, Castner J. Health care provider social network analysis: A systematic review. *Nurs Outlook*. 2015; 63:566–584.
11. Hu H, Yang Y, Zhang C, Huang C, Guan X, Shi L. Review of social networks of professionals in health-care settings—where are we and what else is needed? *Global Health*. 2021; 17:139. <https://doi.org/10.1186/s12992-021-00772-7> PMID: 34863221
12. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med*. 2018; 169:467–473. <https://doi.org/10.7326/M18-0850> PMID: 30178033
13. van der Eijk M, Bloem BR, Nijhuis FAP, Koetsenruijter J, Vrijhoef HJM, Munneke M, et al. Multidisciplinary collaboration in professional networks for PD: A Mixed-method analysis. *Journal of Parkinson's Disease*. 2015; 5:937–945. <https://doi.org/10.3233/JPD-150673> PMID: 26444096
14. Barnett ML, Landon BE, O'Malley AJ, Keating NL, Christakis NA. Mapping physician networks with self-reported and administrative data. *Health Serv Res*. 2011; 46:1592–1609. <https://doi.org/10.1111/j.1475-6773.2011.01262.x> PMID: 21521213
15. Lower TE, Fragar L, Depczynski J, Fuller J, Challinor K, Williams W. Social network analysis for farmers' hearing services in a rural community. *Aust J Prim Health*. 2010; 16:47–51. <https://doi.org/10.1071/py09043> PMID: 21133298
16. Zappa P. The network structure of knowledge sharing among physicians. *Quality & Quantity. International Journal of Methodology*; 2011. p. 1109–26.
17. Quinlan E, Robertson S. Mutual understanding in multi-disciplinary primary health care teams. *J Interprof Care*. 2010; 24(5):565–578. <https://doi.org/10.3109/13561820903520385> PMID: 20102267
18. Jippes E, Achterkamp MC, Brand PLP, Kiewiet DJ, Pols J, van Engelen JML. Disseminating educational innovations in health care practice: training versus social networks. *Soc Sci Med*. 2010; 70:1509–1517. <https://doi.org/10.1016/j.socscimed.2009.12.035> PMID: 20199840
19. Creswick N, Westbrook JI. Social network analysis of medication advice-seeking interactions among staff in an Australian hospital. *International Journal of Medical Informatics*. 2010; 79:e116–125. <https://doi.org/10.1016/j.ijmedinf.2008.08.005> PMID: 19008147
20. Benham-Hutchins MM, Effken JA. Multi-professional patterns and methods of communication during patient handoffs. *International Journal of Medical Informatics*. 2010; 79(4):252–67. <https://doi.org/10.1016/j.ijmedinf.2009.12.005> PMID: 20079686
21. Walton JM, Steinert Y. Patterns of interaction during rounds: implications for work-based learning. *Med Educ*. 2010 Jun; 44:550–558. <https://doi.org/10.1111/j.1365-2923.2010.03626.x> PMID: 20604851
22. Sykes TA, Venkatesh V, Rai A. Explaining physicians' use of EMR systems and performance in the shakedown phase. *Journal of the American Medical Informatics Association: JAMIA*. 2011 Mar; 18:125–130. <https://doi.org/10.1136/jamia.2010.009316> PMID: 21292704
23. Boyer L, Belzeaux R, Maurel O, Baumstarck-Barrau K, Samuelian JC. A social network analysis of healthcare professional relationships in a French hospital. *Int J Health Care Qual Assur*. 2010; 23:460–469. <https://doi.org/10.1108/09526861011050501> PMID: 20845677

24. Mascia D, Cicchetti A, Fantini MP, Damiani G, Ricciardi W. Physicians' propensity to collaborate and their attitude towards EBM: a cross-sectional study. *BMC Health Serv Res.* 2011; 11:172. <https://doi.org/10.1186/1472-6963-11-172> PMID: 21787395
25. Hurtado DA, Greenspan SA, Dumet LM, Heinonen GA. Use of Champions Identified by Social Network Analysis to Reduce Health Care Worker Patient-Assist Injuries. *Joint Commission Journal on Quality & Patient Safety.* 2020 Nov; 46:608–616. <https://doi.org/10.1016/j.jcjq.2020.08.003> PMID: 32893178
26. Benton DC. Mapping and changing informal nurse leadership communication pathways in a health system. *Asian Nurs Res (Korean Soc Nurs Sci).* 2015; 9:28–34. <https://doi.org/10.1016/j.anr.2014.10.006> PMID: 25829207
27. Lee Y, McLaws ML, Ong LM, Husin S, Chua H, Wong S, et al. Hand hygiene—Social network analysis of peer-identified and management-selected change agents. *Antimicrob Resist Infect Control.* 2019; 8.
28. Patterson PD, Arnold RM, Abebe K, Lave JR, Krackhardt D, Carr M, et al. Variation in emergency medical technician partner familiarity. *Health Services Research.* 2011; 46:1319–1331 <https://doi.org/10.1111/j.1475-6773.2011.01241.x> PMID: 21306367
29. Beek APA van, Wagner C, Frijters DHM, Ribbe MW, Groenewegen PP. The ties that bind? Social networks of nursing staff and staff's behaviour towards residents with dementia. *Social Networks.* 2013; 35:347–356.
30. Tsang SS, Chen TY, Wang SF, Tai HL. Nursing work stress: the impacts of social network structure and organizational citizenship behavior. *J Nurs Res.* 2012; 20:9–18. <https://doi.org/10.1097/JNR.0b013e318247bf11> PMID: 22333962
31. Hinami K, Ray MJ, Doshi K, Torres M, Aks S, Shannon JJ, Trick WE. Prescribing Associated with High-Risk Opioid Exposures Among Non-cancer Chronic Users of Opioid Analgesics: a Social Network Analysis. *J Gen Intern Med.* 2019; 34:2443–2450. <https://doi.org/10.1007/s11606-019-05114-3> PMID: 31420823
32. Paul S, Keating NL, Landon BE, O'Malley AJ. Results from using a new dyadic-dependence model to analyze sociocentric physician networks. *Soc Sci Med.* 2014; 117:67–75. <https://doi.org/10.1016/j.socscimed.2014.07.014> PMID: 25047711
33. Bae SH, Farasat A, Nikolaev A, Seo JY, Foltz-Ramos K, Fabry D, et al. Nursing teams: behind the charts. *J Nurs Manag.* 2017; 25:354–365. <https://doi.org/10.1111/jonm.12473> PMID: 28294446
34. Pomare C, Long JC, Ellis LA, Churruca K, Braithwaite J. Interprofessional collaboration in mental health settings: a social network analysis. *J Interprof Care.* 2019; 33:497–503. <https://doi.org/10.1080/13561820.2018.1544550> PMID: 30411988
35. Malik AU, Willis CD, Hamid S, Ulikpan A, Hill PS. Advancing the application of systems thinking in health: Advice seeking behavior among primary health care physicians in Pakistan. *Health research policy and systems.* *BioMed Central.* 2014; 12:43.
36. Wong J. Structure and function of teams in the PICU: A social network analysis. *Critical Care Medicine.* 2015; 43(12 SUPPL. 1):216.
37. Altalib HH, Lanham HJ, McMillan KK, Habeeb M, Fenton B, Cheung KH, et al. Measuring coordination of epilepsy care: A mixed methods evaluation of social network analysis versus relational coordination. *Epilepsy Behav.* 2019; 97:197–205. <https://doi.org/10.1016/j.yebeh.2019.05.023> PMID: 31252279
38. Fuller J, Hermeston W, Passey M, Fallon T, Muyambi K. Acceptability of participatory social network analysis for problem-solving in Australian Aboriginal health service partnerships. *BMC Health Serv Res.* 2012; 12:152. <https://doi.org/10.1186/1472-6963-12-152> PMID: 22682504
39. Patterson PD, Pfeiffer AJ, Weaver MD, Krackhardt D, Arnold RM, Yealy DM, et al. Network analysis of team communication in a busy emergency department. *BMC Health Serv Res.* 2013; 13:109. <https://doi.org/10.1186/1472-6963-13-109> PMID: 23521890
40. Mascia D, Dandi R, di Vincenzo F. Professional networks and EBM use: A study of inter-physician interaction across levels of care. *Health Policy.* 2014; 118:24–36. <https://doi.org/10.1016/j.healthpol.2014.06.003> PMID: 25022323
41. Shoham DA, Mundt MP, Gamelli RL, McGaghie WC. The social network of a burn unit team. *Journal of Burn Care and Research.* 2015; 36:551–557. <https://doi.org/10.1097/BCR.0000000000000218> PMID: 25501788
42. Burt RS, Meltzer DO, Seid M, Borgert A, Chung JW, Colletti RB, et al. What's in a name generator? Choosing the right name generators for social network surveys in healthcare quality and safety research. *BMJ Qual Saf.* 2012; 21:992–1000. <https://doi.org/10.1136/bmjqs-2011-000521> PMID: 22942400

43. Fong A, Clark L, Cheng T, Franklin E, Fernandez N, Ratwani R, et al. Identifying influential individuals on intensive care units: using cluster analysis to explore culture. *J Nurs Manag.* 2017; 25:384–391. <https://doi.org/10.1111/jonm.12476> PMID: 28568480
44. Espinoza P, Peduzzi M, Agreli HF, Sutherland MA. Interprofessional team member's satisfaction: a mixed methods study of a Chilean hospital. *Hum Resour Health.* 2018; 16:30. <https://doi.org/10.1186/s12960-018-0290-z> PMID: 29996936
45. Edge R, Keegan T, Isba R, Diggle P. Observational study to assess the effects of social networks on the seasonal influenza vaccine uptake by early career doctors. *BMJ Open.* 2019; 9:e026997. <https://doi.org/10.1136/bmjopen-2018-026997> PMID: 31471430
46. Hornbeck T, Naylor D, Segre AM, Thomas G, Herman T, Polgreen PM. Using sensor networks to study the effect of peripatetic healthcare workers on the spread of hospital-associated infections. *J Infect Dis.* 2012; 206:1549–1557. <https://doi.org/10.1093/infdis/jis542> PMID: 23045621
47. Aylward BS, Odar CC, Kessler ED, Canter KS, Roberts MC. Six degrees of separation: an exploratory network analysis of mentoring relationships in pediatric psychology. *J Pediatr Psychol.* 2012 Oct; 37:972–979. <https://doi.org/10.1093/jpepsy/jss078> PMID: 22739360
48. Benton DC, Ferguson SL. How nurse leaders are connected internationally. *Nurs Stand.* 2014; 29:42–48. <https://doi.org/10.7748/ns.29.16.42.e9060> PMID: 25515483
49. Yuan CT, Nembhard IM, Kane GC. The influence of peer beliefs on nurses' use of new health information technology: A social network analysis. *Social Science and Medicine.* 2020; 255:113002. <https://doi.org/10.1016/j.socscimed.2020.113002> PMID: 32353652
50. Sibbald SL, Wathen CN, Kothari A, Day AMB. Knowledge flow and exchange in interdisciplinary primary health care teams (PHCTs): an exploratory study. *J Med Libr Assoc.* 2013; 101:128–137. <https://doi.org/10.3163/1536-5050.101.2.008> PMID: 23646028
51. Lazzari C, Kotera Y, Thomas H. Social Network Analysis of Dementia Wards in Psychiatric Hospitals to Explore the Advancement of Personhood in Patients with Alzheimer's Disease. *Curr Alzheimer Res.* 2019; 16:505–517. <https://doi.org/10.2174/1567205016666190612160955> PMID: 31195945
52. Long JC, Cunningham FC, Carswell P, Braithwaite J. Patterns of collaboration in complex networks: the example of a translational research network. *BMC Health Serv Res.* 2014; 14:225. <https://doi.org/10.1186/1472-6963-14-225> PMID: 24885971
53. Barth S, Schraagen JM, Schmettow M. Network measures for characterising team adaptation processes. *Ergonomics.* 2015; 58:1287–1302. <https://doi.org/10.1080/00140139.2015.1009951> PMID: 25677587
54. Mundt MP, Gilchrist VJ, Fleming MF, Zakletskaia LI, Tuan WJ, Beasley JW. Effects of primary care team social networks on quality of care and costs for patients with cardiovascular disease. *Ann Fam Med.* 2015; 13:139–148. <https://doi.org/10.1370/afm.1754> PMID: 25755035
55. Groenen CJM, van Duijnhoven NTL, Faber MJ, Koetsenruijter J, Kremer JAM, Vandenbussche FPHA. Use of social network analysis in maternity care to identify the profession most suited for case manager role. *Midwifery.* 2017; 45:50–55. <https://doi.org/10.1016/j.midw.2016.12.007> PMID: 28024229
56. Anderson C, Talsma A. Characterizing the structure of operating room staffing using social network analysis. *Nurs Res.* 2011; 60:378–85. <https://doi.org/10.1097/NNR.0b013e3182337d97> PMID: 22048555
57. Stewart SA, Abidi SSR. Applying social network analysis to understand the knowledge sharing behaviour of practitioners in a clinical online discussion forum. *J Med Internet Res.* 2012; 14:e170. <https://doi.org/10.2196/jmir.1982> PMID: 23211783
58. Li X, Verspoor K, Gray K, Barnett S. Analysing Health Professionals' Learning Interactions in an Online Social Network: A Longitudinal Study. *Stud Health Technol Inform.* 2016; 227:93–99. PMID: 27440295
59. Yuce YK, Zayim N, Oguz B, Bozkurt S, Isleyen F, Gulkesen KH. Analysis of social networks among physicians employed at a medical school. *Stud Health Technol Inform.* 2014; 205:543–547. PMID: 25160244
60. Dauvrin M, Lorant V. Leadership and cultural competence of healthcare professionals: a social network analysis. *Nurs Res.* 2015; 64:200–210. <https://doi.org/10.1097/NNR.000000000000092> PMID: 25871625
61. Wagter JM, van de Bunt G, Honing M, Eckenhausen M, Scherpbier A. Informal interprofessional learning: visualizing the clinical workplace. *J Interprof Care.* 2012; 26:173–182. <https://doi.org/10.3109/13561820.2012.656773> PMID: 22332642

62. Blanchet K, James P. The role of social networks in the governance of health systems: the case of eye care systems in Ghana. *Health Policy Plan.* 2013; 28:143–156. <https://doi.org/10.1093/heapol/czs031> PMID: 22411882
63. Tighe PJ, Smith JC, Boezaart AP, Lucas SD. Social network analysis and quantification of a prototypical acute pain medicine and regional anesthesia service. *Pain Med.* 2012; 13:808–819. <https://doi.org/10.1111/j.1526-4637.2012.01379.x> PMID: 22568636
64. Tavakoli Taba S, Hossain L, Heard R, Brennan P, Lee W, Lewis S, et al. Personal and Network Dynamics in Performance of Knowledge Workers: A Study of Australian Breast Radiologists. *PLoS One.* 2016 Feb 26; 11:e0150186. <https://doi.org/10.1371/journal.pone.0150186> PMID: 26918644
65. Sullivan P, Saatchi G, Younis I, Harris ML. Diffusion of knowledge and behaviours among trainee doctors in an acute medical unit and implications for quality improvement work: A mixed methods social network analysis. *BMJ Open.* 2019; 9:e027039. <https://doi.org/10.1136/bmjopen-2018-027039> PMID: 31826886
66. Bachand J, Soulos PR, Herrin J, Pollack CE, Xu X, Ma X, et al. Physician peer group characteristics and timeliness of breast cancer surgery. *Breast Cancer Res Treat.* 2018; 170:657–665. <https://doi.org/10.1007/s10549-018-4789-8> PMID: 29693229
67. Hurtado DA, Dumet LM, Greenspan SA, Rodriguez YI. Social Network Analysis of peer-specific safety support and ergonomic behaviors: An application to safe patient handling. *Appl Ergon.* 2018; 68:132–137. <https://doi.org/10.1016/j.apergo.2017.11.009> PMID: 29409627
68. Palazzolo M, Grippa F, Booth A, Rechner S, Bucuvalas J, Gloor P. Measuring Social Network Structure of Clinical Teams Caring for Patients with Complex Conditions. *Procedia-Social and Behavioral Sciences*; 2011;17–29. (*Procedia Social and Behavioral Sciences*; vol. 1;26).
69. Uddin S, Hossain L, Hamra J, Alam A. A study of physician collaborations through social network and exponential random graph. *BMC Health Serv Res.* 2013; 13:234. <https://doi.org/10.1186/1472-6963-13-234> PMID: 23803165
70. Alexander GL, Steege LM, Pasupathy KS. Case studies of IT sophistication in nursing homes: A mixed method approach to examine communication strategies about pressure ulcer prevention practices. *International Journal of Industrial Ergonomics.* 2015; 49:156–66.
71. Laapotti T, Mikkola L. Social interaction in management group meetings: a case study of Finnish hospital. *J Health Organ Manag.* 2016; 30:613–629. <https://doi.org/10.1108/JHOM-02-2015-0040> PMID: 27296882
72. Dauvrin M, Lorant V. Cultural competence and social relationships: a social network analysis. *Int Nurs Rev.* 2017; 64:195–204. <https://doi.org/10.1111/inr.12327> PMID: 27859147
73. Shokoohi M, Nedjat S, Majdzadeh R. A social network analysis on clinical education of diabetic foot. *J Diabetes Metab Disord.* 2013; 12:44. <https://doi.org/10.1186/2251-6581-12-44> PMID: 24330538
74. Quinlan E, Robertson S. The communicative power of nurse practitioners in multidisciplinary primary healthcare teams. *J Am Assoc Nurse Pract.* 2013; 25:91–102. <https://doi.org/10.1111/j.1745-7599.2012.00768.x> PMID: 23347245
75. Brewer BB, Carley KM, Benham-Hutchins M, Effken JA, Reminga J. Exploring the stability of communication network metrics in a dynamic nursing context. *Social Networks.* 2020; 61:11–19. <https://doi.org/10.1016/j.socnet.2019.08.003> PMID: 32863552
76. Kawamoto E, Ito-Masui A, Esumi R, Ito M, Mizutani N, Hayashi T, et al. Social Network Analysis of Intensive Care Unit Health Care Professionals Measured by Wearable Sociometric Badges: Longitudinal Observational Study. *J Med Internet Res.* 2020; 22:e23184. <https://doi.org/10.2196/23184> PMID: 33258785
77. Cavalcante JB, Da-Silva-Junior GB, Bastos MLA, Costa MEM, Santos A de L, Maciel RHM de O. Relationship network at a mobile urgent care service unit: analysis of a work team. *Revista brasileira de medicina do trabalho: publicacao oficial da Associacao Nacional de Medicina do Trabalho-ANAMT.* 2018; 16(2):158–66.
78. Benammi S, Madani N, Abidi K, Dendane T, Zeggwagh A, Belayachi J. Team communication in an acute medical unit: A Social network analysis. *Annals of Intensive Care.* 2019; 9(SUPPL. 1).
79. Lai YH. The social network analysis on the behavioral intention to use cloud sphygmomanometer. *Health and Technology.* 2020; 10:787–794.
80. Crockett LK, Leggett C, Curran J, Knisley L, Brockman G, Scott S, et al. Knowledge sharing between general and pediatric emergency departments: Connections, barriers, and opportunities. *Canadian Journal of Emergency Medicine.* 2018; 20(4):523–31. <https://doi.org/10.1017/cem.2018.7> PMID: 29467040

81. Choudhury A, Kaushik S, Dutt V. Social-network analysis in healthcare: analysing the effect of weighted influence in physician networks. *Network Modeling Analysis in Health Informatics and Bioinformatics*. 2018; 7:17.
82. Shafiei S, Azar A. Mapping and social network analysis of the nurses of Razi hospital. *Iranian Red Crescent Medical Journal*. 2018; 20:e58321.
83. Zappa P. Assessing Cooperation in Open Systems: An Empirical Test in Healthcare. In: *Analysis And Modeling Of Complex Data In Behavioral And Social Sciences*. Berlin: Springer. 2014:293–301.
84. Chung KSK. Understanding Decision Making through Complexity in Professional Networks. *Advances in Decision Sciences*. 2014: 215218.
85. Tasselli S. Social networks and inter-professional knowledge transfer: The case of healthcare professionals. *Organization Studies*. 2015: 36:841–72.
86. Venkatesh V, Zhang X, Sykes TA. “Doctors do too little technology”: A longitudinal field study of an electronic healthcare system implementation. *Information Systems Research. Institute for Operations Research & the Management Sciences (INFORMS)*; 2011; 22: 523–546.
87. Currie G, White L. Inter-professional barriers and knowledge brokering in an organizational context: The case of healthcare. *Organization Studies*. 2012; 33:1333–1361.
88. Durojaiye A, Fackler J, McGeorge N, Webster K, Kharrazi H, Gurses A. Examining Diurnal Differences in Multidisciplinary Care Teams at a Pediatric Trauma Center Using Electronic Health Record Data: Social Network Analysis. *Journal of Medical Internet Research*. 2022; 24:e30351. <https://doi.org/10.2196/30351> PMID: 35119372
89. Smit LC, Dikken J, Moolenaar NM, Schuurmans MJ, de Wit NJ, Bleijenberg N. Implementation of an interprofessional collaboration in practice program: a feasibility study using social network analysis. *Pilot Feasibility Stud*. 2021; 7:7. <https://doi.org/10.1186/s40814-020-00746-3> PMID: 33407919
90. Haruta J, Tsugawa S. What Types of Networks Do Professionals Build, and How Are They Affected by the Results of Network Evaluation? *Front Public Health*. 2021; 9:758809. <https://doi.org/10.3389/fpubh.2021.758809> PMID: 34888285
91. Kim EJ, Lim JY, Kim GM, Kim SK. Nursing students’ subjective happiness: A social network analysis. *International Journal of Environmental Research and Public Health*. 2021; 18:11612. <https://doi.org/10.3390/ijerph182111612> PMID: 34770124
92. Mukinda FK, van Belle S, Schneider H. Local Dynamics of Collaboration for Maternal, Newborn and Child Health: A Social Network Analysis of Healthcare Providers and Their Managers in Gert Sibande District, South Africa. *Int J Health Policy Manag*. 2021;1–11.
93. Hayward BA, McKay-Brown L, Poed S, McVilly K. Identifying important persons in the promotion of positive behaviour support (pbs) in disability services: A social network analysis. *Journal of Intellectual and Developmental Disability*. 2021; 47:292–307.
94. Bertoni VB, Saurin TA, Fogliatto FS. How to identify key players that contribute to resilient performance: A social network analysis perspective. *Safety Science*. 2022; 148:105648.
95. Yousefi Nooraie R, Lohfeld L, Marin A, Hanneman R, Dobbins M. Informing the implementation of evidence-informed decision making interventions using a social network analysis perspective; a mixed-methods study. *BMC Health Serv Res*. 2017; 17:122. <https://doi.org/10.1186/s12913-017-2067-9> PMID: 28178958
96. Mundt MP, Zakletskaia LI. Professional Communication Networks and Job Satisfaction in Primary Care Clinics. *Annals of Family Medicine*. 2019; 17:428–435. <https://doi.org/10.1370/afm.2442> PMID: 31501206
97. Llupia A, Puig J, Mena G, Bayas JM, Trilla A. The social network around influenza vaccination in health care workers: a cross-sectional study. *Implement Sci*. 2016; 11:152. <https://doi.org/10.1186/s13012-016-0522-3> PMID: 27881186
98. Marques-Sanchez P, Munoz-Doyague MF, Martinez Y v, Everett M, Serrano-Fuentes N, van Bogaert P, et al. The Importance of External Contacts in Job Performance: A Study in Healthcare Organizations Using Social Network Analysis. *Int J Environ Res Public Health*. 2018; 15:1345. <https://doi.org/10.3390/ijerph15071345> PMID: 29954054
99. Shoham DA, Harris JK, Mundt M, McGaghie W. A network model of communication in an interprofessional team of healthcare professionals: A cross-sectional study of a burn unit. *Journal of Interprofessional Care*. 2016; 30:661–667. <https://doi.org/10.1080/13561820.2016.1203296> PMID: 27436781
100. Mascia D, Cicchetti A, Damiani G. “Us and them”: A social network analysis of physicians’ professional networks and their attitudes towards EBM. *BMC Health Serv Res*. 2013; 13:429. <https://doi.org/10.1186/1472-6963-13-429> PMID: 24148207

101. Creswick N, Westbrook JI. Who Do Hospital Physicians and Nurses Go to for Advice About Medications? A social network analysis and examination of prescribing error rates. *J Patient Saf.* 2015; 11:152–159. <https://doi.org/10.1097/PTS.000000000000061> PMID: 24583953
102. Meltzer D, Chung J, Khalili P, Marlow E, Arora V, Schumock G, et al. Exploring the use of social network methods in designing healthcare quality improvement teams. *Soc Sci Med.* 2010; 71:1119–1130. <https://doi.org/10.1016/j.socscimed.2010.05.012> PMID: 20674116
103. Polgreen PM, Tassier TL, Pemmaraju SV, Segre AM. Prioritizing healthcare worker vaccinations on the basis of social network analysis. *Infection Control and Hospital Epidemiology.* 2010; 31:893–900. <https://doi.org/10.1086/655466> PMID: 20649412
104. Hossain L, Kit Guan DC. Modelling coordination in hospital emergency departments through social network analysis. *Disasters.* 2012; 36:338–364. <https://doi.org/10.1111/j.0361-3666.2010.01260.x> PMID: 22409650
105. Pinelli VA, Papp KK, Gonzalo JD. Interprofessional Communication Patterns During Patient Discharges: A Social Network Analysis. *Journal of General Internal Medicine.* 2015; 30:1299–1306. <https://doi.org/10.1007/s11606-015-3415-2> PMID: 26173532
106. Gorley C, Lindstrom RR, McKeown S, Krause C, Pamplin C, Sweet D, et al. Exploring distributed leadership in the BC Sepsis Network. *Healthcare Management Forum.* 2016; 29:63–66. <https://doi.org/10.1177/0840470415606451> PMID: 26872797
107. Mascia D, di Vincenzo F, Iacopino V, Fantini MP, Cicchetti A. Unfolding similarity in interphysician networks: the impact of institutional and professional homophily. *BMC Health Serv Res.* 2015; 15:92. <https://doi.org/10.1186/s12913-015-0748-9> PMID: 25890319
108. Li Z, Xu X. Analysis of network structure and doctor behaviors in e-health communities from a social-capital perspective. *Int J Environ Res Public Health.* 2020; 17:1136. <https://doi.org/10.3390/ijerph17041136> PMID: 32053913
109. Sykes M, Gillespie BM, Chaboyer W, Kang E. Surgical team mapping: implications for staff allocation and coordination. *AORN J.* 2015; 101:238–248. <https://doi.org/10.1016/j.aorn.2014.03.018> PMID: 25645040
110. Assegaai T, Schneider H. The supervisory relationships of community health workers in primary health care: social network analysis of ward-based outreach teams in Ngaka Modiri Molema District, South Africa. *BMJ Glob Health.* 2019; 4:e001839. <https://doi.org/10.1136/bmjgh-2019-001839> PMID: 31908861
111. Mascia D, Pallotti F, Dandi R. Determinants of knowledge-sharing networks in primary care. *Health Care Management Review.* 2018 Dec; 43:104–114. <https://doi.org/10.1097/HMR.000000000000139> PMID: 27984404
112. Tighe PJ, Davies L, Lucas SD, Bernard HR. Connections The Operating Room: It's a Small World (and Scale Free Network) After All. 2014; 34:1.
113. Cannavacciuolo L, Iandoli L, Ponsiglione C, Maracine V, Scariat E, Nica AS. Mapping knowledge networks for organizational re-design in a rehabilitation clinic. *Business Process Management Journal.* 2017; 23:329–348.
114. Kothari A, Hamel N, MacDonald JA, Meyer M, Cohen B, Bonnenfant D. Exploring community collaborations: Social network analysis as a reflective tool for public health. Vol. 27, *Systemic Practice and Action Research.* 2014;123–137.
115. Xu B, Zhang Y, Chen L, Yu L, Li L, Wang Q. The influence of social network on COVID-19 vaccine hesitancy among healthcare workers: a cross-sectional survey in Chongqing, China. *Human Vaccines and Immunotherapeutics.* 2021; 17:5048–5062. <https://doi.org/10.1080/21645515.2021.2004837> PMID: 34982646
116. Agneessens F, Labianca G (Joe). Collecting survey-based social network information in work organizations. *Social Networks.* 2022; 68:31–47.
117. Pomare C, Churruca K, Long JC, Ellis LA, Braithwaite J. Organisational change in hospitals: A qualitative case-study of staff perspectives. *BMC Health Services Research.* 2019; 19:1–10.
118. Cronin B, Perra N, Rocha LEC, Zhu Z, Pallotti F, Gorgoni S, et al. Ethical implications of network data in business and management settings. *Social networks.* 2021; 67:29–40.
119. Borgatti SP, Molina JL. Ethical and Strategic Issues in Organizational Social Network Analysis. *The Journal of Applied Behavioral Science.* 2003 Sep 1; 39:337–49.
120. Battilana J, Casciaro T. Change agents, networks, and institutions: A contingency theory of organizational change. *Academy of Management Journal.* 2012; 55:381–398.
121. Krackhardt D, Porter LW. When Friends Leave: A Structural Analysis of the Relationship between Turnover and Stayers' Attitudes. *Administrative Science Quarterly.* 1985 Feb 20; 30:242–261.

122. Schaefer DR, Adams J, Haas SA. Social Networks and Smoking: Exploring the Effects of Peer Influence and Smoker Popularity Through Simulations. *Health Education and Behavior*. 2013; 40(1 SUPPL.):1–15. <https://doi.org/10.1177/1090198113493091> PMID: 24084397