

The Use of Plant Steroids in Viral Disease Treatments: Current Status and Future Perspectives

Pinar Obakan Yerlikaya^{1,2} , Elif Damla Arisan³ , Leila Mehdizadehtapeh⁴ ,
Pinar Uysal-Onganer⁵ , Ajda Coker-Gurkan⁶ 

¹Istanbul Medeniyet University, Faculty of Engineering and Natural Sciences, Department of Molecular Biology and Genetics, Istanbul, Turkiye

²Istanbul Medeniyet University, Science and Advanced Technology Research Center, Istanbul, Turkiye

³Gebze Technical University, Institute of Biotechnology, Kocaeli, Turkiye

⁴Istanbul Sabahattin Zaim University, Department of Food and Nutrition, Halkali Campus, Istanbul, Turkiye

⁵University of Westminster, Cancer Research Group, School of Life Sciences, London, United Kingdom

⁶Marmara University, Faculty of Science and Literature, Department of Biology, Istanbul, Turkiye

ORCID IDs of the authors: P.O.Y. 0000-0001-7058-955X, E.D.A. 0000-0002-4844-6381, L.M. 0000-0001-8759-5016, P.U.O. 0000-0003-3190-8831, A.C.G. 0000-0003-2298-5755

Please cite this article as: Obakan Yerlikaya P, Arisan ED, Mehdizadehtapeh L, Uysal-Onganer P, Coker-Gurkan A. The Use of Plant Steroids in Viral Disease Treatments: Current Status and Future Perspectives. Eur J Biol. Advance Online Publication. DOI: 10.26650/EurJBiol.2023.1130357

ABSTRACT

Plants have been used for the prevention and treatment of diseases since the early days of humankind and constitute the natural sources of today's modern medicine. Approximately one-quarter of approved drugs are derived from plants. Plant steroids are a group of biologically active secondary metabolites with a 5 α and 5 β gonane carbon skeleton. There is immense chemical diversity in plant steroids due to the side chains, oxidation status of the carbons in the tetracyclic core, and methyl groups. Plant steroids are classified into several groups based on their biological functions and structures, also on their mechanism of biosynthesis. All subtypes have been investigated for their anti-cancer, immunomodulatory, anti-inflammatory, and anti-viral properties. The novel coronavirus disease (COVID-19) is caused by severe acute respiratory syndrome coronavirus (SARS-CoV-2), which carries an RNA genome. An intense effort has been made in terms of effective treatment strategies and vaccine development since it was declared a pandemic. Nucleoside analogs such as favipiravir and remdesivir are used to block RNA-dependent RNA polymerase enzymes. Other strategies including neuraminidase inhibitors, chloroquine, and hydroxychloroquine as immunomodulatory agents, stem cell and cytokine-based therapies are being conducted. One part of the therapies against SARS-CoV-2 is focused on the spike (S) protein of the virus that binds to the host receptor, angiotensin-converting enzyme 2 (ACE2). It has been suggested that SARS-CoV-2 S protein has a free fatty acid-binding pocket, and according to molecular simulations, steroids are ligands that bind to this pocket. Therefore, this review summarizes the plant steroid biological actions as well as their anti-viral potential against SARS-CoV-2 infection.

Keywords: *In silico*, Plant steroids, anti-viral drug, SARS-CoV-2

INTRODUCTION

PLANT STEROIDS AND THEIR BIOLOGICAL ACTIONS

Plant steroids are unique compounds found throughout the plant kingdom, which exert critical physiological effects including plant growth, development, and reproduction. They are also important micronutrients commonly found

in a diet.¹ Plant steroids have a definite chemical structure composed of four carbon rings called the steroid nucleus (Figure 1). The nucleus typically contains the tetracyclic 5 α or 5 β -gonane carbon skeleton with methyl substituents at C-10, C-13, and an alkyl side chain at C-17 (Figure 1).

Plant steroids are quite similar to those of well-known animal steroids in terms of the regulation of the ex-



Corresponding Author: Pinar Obakan Yerlikaya

E-mail: pinar.obakan@medeniyet.edu.tr

Submitted: 14.06.2022 • **Revision Requested:** 27.10.2022 • **Last Revision Received:** 07.03.2023 •

Accepted: 15.03.2023 • **Published Online:** 13.04.2023

Content of this journal is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.



pression of specific genes and complex physiological processes involved in cell growth and division. The addition of distinct chemical groups at different positions to the nucleus leads to the formation of many types of steroids categorized into seven main groups based on their biological functions, structures, and taxonomic considerations such as brassinosteroids, bufadienolides, cardenolides, cucurbitacins, ecdysteroids, saponins, steroidal alkaloids, and withanolides (Table.1).^{2,3}

Along with their important physiological functions within plants, they exhibit pharmacological activities helpful to humankind. Due to the knowledge about the pharmacological activities of these herbs, they are gaining increasing attention all over the world as herbal medicine supplements.⁴ The pharmacological actions of these various types of plant steroids indicated that they have growth-promoting, cardiostimulant, anti-microbial, anti-tumor, hepatoprotective, anti-fungal, antioxidant, anti-inflammatory, and anti-viral roles.⁵⁻⁸




Figure 1. Plant steroids backbone structure. Created by ChemSpider.

Brassinosteroids

One of the most studied classes of plant steroids is brassinosteroids (BRs). BRs were discovered in 1979 by Grove et al. and have been isolated from the pollen of *Brassica napus*.⁹ The most bioactive BR was referred to as brassinolide, followed by castasterone (CS) among more than fifty brassinosteroids identified to date.¹⁰ BRs regulate growth in very low concentrations in plants and have the ability to make impacts on physiological processes (germination, growth, reproductive development, etc), modulation of the antioxidative enzyme cascade and protection from various stress conditions, including drought, heavy metals, herbicidal injury, thermotolerance, and salinity^{11,12}. Antigenotoxicity effect of 24-epibrassinolide has been verified employing *Allium cepa* chromosomal aberration assay.¹³ *In vitro* analyses in neuronal PC12 mammalian cell line demonstrated that the potent antioxidant and neuroprotective role of 24-epibrassinolide reduce DNA fragmentation, Bax/Bcl-2 protein rate and cleaved caspase-3 and inhibit the MPP⁺-induced (1-methyl-4-phenylpyridinium) apoptosis in dopaminergic neurons.¹⁴ BRs have numerous impacts on plant physiological processes including germination, growth, reproductive development, and protection from environmental stresses including drought, heavy metals, and salinity.¹⁵ For instance, 24-epibrassinolide (EBR) has been shown to increase thermotolerance of tomato plants via heat-shock proteins.¹⁶ BRs, apart from their well-known roles in the improvement in quality and yield of crops, have been shown to inhibit cancer cell survival and replication of viruses. 28-homocastasterone (homoCS) and EBR were first suggested as nuclear hormone receptor blockers and to cause cell cycle arrest in prostate and breast cancer cells. Recently, Coskun et al showed that EBR can trigger mitochondria-mediated apoptosis in an endoplasmic reticulum stress-inducing manner in prostate and colon cancer cell lines without causing the same effect in normal epithelial cells due to the autophagic induction¹⁷. Their findings showed that calreticulin alteration, an ER-resi-

Table 1. Major groups of plant steroids.

Plant steroid group	Examples
Phytosterols	β -Sitosterol , campesterol , stigmasterol
Brassinosteroids	Brassinolide , castasterone
Withanolides (withasteroids)	Withanolide A ,withanolides Q
Phytoecdysteroids	20-Hydroxyecdysone
Steroidal alkaloids /glycoalkaloids	Jervine , α -solanine , conessine
Steroidal saponins /sapogenins	Tigogenin,25-Epi-ruizgenin
Mammalian steroidal hormones	Androstenone , progesterone
Bufadienolides	19-nor bufadienolides
Cardenolides	Quabain
Cucurbitacins	Cucurbitacins(A,B,C,D)
Ecdysteroids	Dacryhainansterone

dent chaperone protein, was the main phenomenon in EBR-induced apoptosis. The anti-angiogenic roles of BRs were also identified. Both EBR and homoCS inhibited the proliferation, adhesion, and migration of HMEC-1/HUVEC endothelial cells.^{18,19} BRs were shown to be effective against several disorders apart from cancer, e.g., Alzheimer's and Huntington's Diseases, sexual differentiation disorders, steroid-induced osteoporosis and cataract, hyperadrenocorticism, etc.²⁰

Bufadienolides

Plant steroid subgroups, bufadienolides, and cardenolides, are described as cardiac glycosides because they increase the contractile force of the heart via the inhibition of cardiac Na⁺-K⁺-ATPase.^{21,22} They consist of a steroid nucleus bearing a pentadienolide ring at C17 β . Bufadienolides and cardenolides containing plants can cause severe toxicity to livestock. Besides, they have a wide range of bioactivities including blood pressure stimulating, immunoregulatory, etc.²³ They also exert cytotoxic effects on HCT116 colon, A549 lung, and HepG2 hepatocellular, A-375a skin, MCF-7 breast cancer cell lines.^{23,24} Bufadienolides and cardenolides are also found in other organisms such as *Bufo marinus* and *Chrysolina coeruleans*, Lampyridae, and Colubridae as well as mammalian tissues.²⁴

Cucurbitacins

Cucurbitacins are steroid-like molecules found in the plants of the Cucurbitaceae family, as well as Cruciferae, Rubiaceae, etc. They possess a lanostane skeleton with hydroxy, methyl, and oxo substituents with unsaturation at positions 5 and 23. Their medicinal and toxic properties have been identified. Certain species-rich in cucurbitacins have been used in traditional medicines against metabolic diseases like diabetes *via* activating the 5' AMP-activated protein kinase pathway.²⁵ The anti-tumor, anti-inflammatory, antidiabetic, and immunosuppressant roles of these metabolites were investigated in several studies. The anti-cancer effect of cucurbitacins, such as elatericin A and B, was discovered more than fifty years ago *via* inhibition of cell proliferation, induction of apoptosis *in vitro*, and tumor growth in tumor-bearing mice. Oi et al cited cucurbitacin as a signal transducer activator of transcription 3 (STAT3 inhibitor and a novel therapeutic agent in osteosarcoma, lung, laryngeal, and breast cancer cell lines.^{25,26} Cucurbitacins can inhibit the expression of tumor necrosis factor (TNF) and proinflammatory mediators such as cyclooxygenase-2 to exert an anti-inflammatory effect.²⁷

Phytoecdysteroids

The natural polyhydroxylated plant steroids group, phytoecdysteroids, appears in many plants as toxins for protection against herbivore insects. More than 300 different phytoecdysteroids have been described, and they show a wide distribution in the plant kingdom.²⁸ Besides, their presence is not restricted to plants, several aquatic plants and fungi also contain ecdysteroids.²⁹ A broad spectrum of these compounds with medicinal properties in mammals has been described, including antimicrobial, hepatoprotective, hypoglycemic, and hypocholesterolemic. The ecdysteroids have very low toxicity in mice with LD50 approximately 6 g kg⁻¹ of body mass.^{30,31} They have been

suggested as skin collagenase inhibitors and accelerators of wound healing. More recently, ecdysteroids isolated from *Vitex doniana* have exhibited antidepressant effect via monoaminergic transmission interference in mice studied by Isohola et al. Besides, the antioxidant and antiproliferative action of phytoecdysteroids have been also reported both *in vitro* and *in vivo* studies.^{32,33} The fact that ecdysteroids influence various cellular processes was explained by their Akt/Protein kinase B signaling targeting.³⁰

Saponins

Saponins are a group of plant steroids with a classical steroid nucleus to which sugar groups are attached. They are subdivided into triterpenoid and steroid glycosides. Saponins are used in many applications in the pharmaceutical industry as a starting material for the synthesis of steroidal hormones such as vitamin D and cardioactive glycosides. They are also used as food additives or firefighting foams due to their foaming and surface tension reducing properties,³⁴ in other words, they are natural surfactants and emulsifiers. Saponins have been proposed for the treatment of diabetes, obesity, and osteoporosis.³⁵ Their anti-inflammatory, hypocholesterolemic, and immune-stimulating roles are also described. They have been suggested as anti-cancer agents because of their ability to inhibit cell proliferation, angiogenesis, oxidative stress and promote apoptosis^{36,37} in a variety of cancer cells *in vitro* including brain, breast, colon, pancreas, prostate, and ovary without causing toxicity and changes and body weight in rats, mice, and rabbits.³⁸⁻⁴⁰

Steroidal Alkaloids

Steroidal alkaloids (SAs) are a diverse class of plant steroids with a steroid nucleus containing nitrogen atoms attached to a ring or side chains.⁴¹ Some examples of steroidal alkaloids are α -solanine, solamargine, tomatidine, solasonine, α -solanine, brioifil-in, and α -chaconine with anti-cancer activity⁴² *via* induction of either intrinsic or extrinsic pathways of apoptosis.^{43,44} The cell cycle arrest and anti-metastatic effect of some SAs have been also reported. Solanidine and spirosolane types are also known to possess pregnane skeleton and exhibit anti-fungal and anti-viral roles.⁴⁵ Antifungal, antiviral, antitumor and embryotoxic activities of steroidal alkaloid glycosides (spirosolane and solanidine) have been proven via various spectroscopic analyses.^{46,47} Recently, both *in vitro* and *in vivo* investigation on *Sarcococca saligna*, the hepatoprotective and immunosuppressive potential of isolated SAs was defined. According to the results in a dose-dependent treatment, human T-cells proliferation, IL-2 production and phytohemagglutinin stimulated T-cell proliferation have been suppressed on a significant scale.⁴⁸

Withanolides

Withanolides are steroidal lactones carrying oxidized C22 and C26 ergostane skeleton and have been found to show immunomodulatory, anti-inflammatory, and anti-microbial activities by altering NF- κ B and JAK/STAT signaling. Therefore, their application in inflammation-mediated chronic diseases has been studied in several studies. Besides, like other plant steroids, withanolides showed antitumor activity in several cancer cell

lines for example, the most potent withanolide, withaferin A caused cell cycle arrest in MCF-7, SUM159, and SK-BR-3 breast cancer cell lines.⁴⁹ Withaferin A exhibited an antiproliferative effect in melanoma, head and neck squamous cell carcinoma, and human glioblastoma by altering Akt/mTOR and MAPK signaling pathways. Another important effect is that withanolides can target breast cancer stem cells by decreasing mammospheres and aldehyde dehydrogenase activity.⁵⁰⁻⁵²

Phytosterols

Phytosterols (plant sterols) are a ubiquitous group of plant sterols. Phytosterols (PS), a member of the triterpene family, are steroidal alcohols. They are known as natural plant metabolites that exhibit functional similarity with sterol-structured cholesterol in mammalian cells. Unlike cholesterol, they are not synthesized endogenously in the human body. More than 250 phytosterols of dietary origin such as legumes, fruits, vegetables, tubers, beans, nuts, wheat germ, whole grains, sunflower seeds and vegetable oil have been defined. Among them, campesterol, β -Sitosterol, and stigmasterol are the most sterols present in edible fats of PSs. Sterols and stanols both in PSs represent a class of sterol composites that modulate essential physiological functions such as membrane fluidity, permeability and signal transduction in plant cells. While sterols are in "free" unbound form, they can be covalently bonded through their ester or glycosidic bonds additionally.⁵³⁻⁵⁵ Various studies emphasize that phytosterol glucosides are critical in the functional organization of plasma membrane lipid rafts, plasma membrane enzymes and receptor proteins. PSs also act as precursors in the synthesis of main bioactive compounds such as steroidal glycoalkaloids, steroidal saponins, phytoecdysteroids, and brassinosteroids.^{2,56,57} As a new functional food group, extensive research has been conducted on the anticancer, anti-diabetes, anti-obesity, cytotoxic, insecticidal/larvicidal, anti-atherosclerosis, anti-Alzheimer and hepatoprotective effects of PSs related to the management of human health and metabolic disorders.⁵⁷ Plant sterols have been proven to increase mitochondrial ATP content, reduce dyslipidemia, insulin resistance and dysfunction of cells, improve gut microbiota dysbiosis and barrier dysfunction, increase adipose inflammatory signal, reduce oxidative stress, and modulate inflammatory signals. The fact that PSs interact directly and / or indirectly with free radicals and specific proteins in redox signaling pathways involved in various physiological processes is actually thought to be due to their antioxidant properties.⁵⁸ Ergosterol, campesterol, β -sitosterol, and stigmasterol, inhibit the production of inflammatory enzymes and pro-inflammatory cytokines in different cell lines. In LPS-stimulated HaCaT human keratinocytes and J774A.1 mouse macrophages, TNF- α , interleukin (IL)-8, IL-1 β and IL-6 secretion have been reduced after the application of β -sitosterol isolated from *moringa oleifera*.⁵⁹ Scientific evidence reveals that PSs modulate the gut microbiota composition and play a protective role against pathologies. In addition, functionality has a beneficial effect on gut inflammation and barrier integrity. *In vitro* faeces analysis of morbidly obese individuals during the fermentation process determined that sterol supplementation decreased the population of *Erysipelotrichaceae*, a family of bac-

teria associated with lipidemic imbalances.⁶⁰ The same microbial changes have been confirmed in mouse and hamster models of hypercholesterolaemia.⁶¹

ANTI-VIRAL ACTIVITIES OF PLANT STEROIDS

Due to the inevitable reality of viral infection risks, modern medical science aims to prevent the transmission and spread of viral infection and also to minimize the treatment and financial burden. In this regard, scientific research and pharmaceutical companies have been examining their potential compounds for the identification and production of new antivirals. Most plants have hormonal functionality and are considered to be steroid synthesis factories. Recent research proves that although there is a structural distinction of plant sterols, they have a viral transcription / replication alteration effect, along with their inhibitory, antioxidant, and immunomodulatory activities, support host defense mechanisms and cell survival in the event of viral infection.^{62,63} With the known details of the preventive role of functional foods and medicine herbs with natural nutraceutical ingredients such as polyphenols, terpenoids, flavonoids, alkaloids, sterols in noncommunicable diseases (NCDs), their preventive and / or therapeutic roles associated with the immune system in viral infectious diseases (CDs) in particular remain a mystery in many respects.^{63,64}

In a recent study, 4 biomarker extracts of *Guiera senegalensis*, a phytoestrogen family were examined against hepatitis B virus in HuH7 liver cell lines, and their antioxidative, antiviral and hepatoprotective potential have been confirmed.⁶⁵ Molecular investigations of β -sitosterol as a common phytosterol in Chinese traditional medicine have been designed in influenza (IAV) infected mouse models in a dose dependent manner β -sitosterol (150–450 μ g/mL) that is able to suppress inflammation *via* p38 mitogen-activated protein kinase (MAPK) and NF- κ B signaling. Additionally, RIG-I signaling, harmful IFN production and acute lung injury inhibition have been reported.⁶⁶

Activation of the resistance mechanisms of BRs against DNA and RNA viruses is the main characteristic of their defense strategies. As a result of the host cell's special sensors (PRRs) encountering with expressed structural motifs (PAMPs) unique to the pathogen virus, immune response mechanisms that inactivate bacteria, viruses, fungi and oomycetes are triggered.^{20,67,68} Recent studies indicated that brassinosteroids and their derivatives inhibited the *in vitro* replication of herpes simplex type 1 (HSV-1) thymidine kinase (TK)+ and TK- strains, and the arenaviruses Junin (agent of Argentine hemorrhagic fever), Pichinde, and Tacaribe viruses.⁶⁹ The structural alterations determine their efficacy on antiviral responses through increasing host recognition mechanisms. BR treatment in tobacco plants substantially reduced *Tobacco mosaic virus* (TMV) viral infection and increased crop yields by 56% with induced resistance to TMV.⁷⁰ The exogenous application of brassinolides to *Nicotiana benthamiana*, was shown as an antiviral mechanism during TMV infection. Brassinosteroids could activate different cellular targets such as MEK2 (MAPKK)-SIPK (salicylic acid-induced protein

kinase) and RBOHB (respiratory burst oxidase homolog protein B)-dependent ROS burst. Additional targets such as BES1/BZR1 could inhibit RBOHB-dependent ROS generation and maintain growth function as a hallmark of plant immunity.²⁰

In a case study, it was shown that the *Arabidopsis thaliana* plant belonging to the cruciferous family, BAK1 and/or BKK1 proteins in the steroid signaling are mandatory to maintain plant immunity against RNA viruses' infection such as Turnip crinkle virus (TCV).⁷¹ Enhancement of antioxidative enzymatic performance and gene expression regulation have been observed in *A. thaliana* infected with cucumber mosaic virus (CMV) after BR treatment.⁷² Shamsabadipour et al. determined the structural properties of triterpene and steroid compounds obtained from *E. denticulate* plant endemic to Iran by NMR and mass spectroscopic approaches. Toxicity and antiviral analysis results proved that both types of compounds have protective effects against Herpes Simplex Type 1 (HSV-1) virus infection.⁷³

The anti-viral activity of bufadienolides and cardenolides has been reviewed by Kamano et al. Buadienolide derivatives showed selective inhibition of Rhinovirus at very low concentrations. In addition, bufadienolides isolated from *Kalanchoe pinnata* were shown to inhibit the activation of Epstein Barr virus.⁷⁴ Besides, a potential anti-bovine viral diarrhoea virus (BVDV) activity of nine members of cucurbitacins was found. In addition, some of the members, namely cucurbitacin B, D and E, have been shown to inhibit Hepatitis C replicon replication in HuH-7 cells.^{75, 76}

Tomatidine in green tomatoes is a steroidal alkaloid with enzymatic activity. An in vitro research focused on antiviral reaction of tomatidine versus Chikungunya virus (CHIKV), highlighting the strong controlling role of this compound on CHIKV replication processes and a significant decrease in the count of infected cells both early or post infection.⁷⁷ The quest for effective anti-PEDV drugs to control the swine epidemic diarrhoea virus (PEDV), defined by the fatal diarrhoea symptom in piglets, continues. Wang and colleagues, by designing in silico and in vitro assays, draws attention to the inhibitory role of tomatidine during PEDV replication cycle by targeting the 3CL protease in IPEC-J2 and Vero cell lines. Findings suggest that tomatidine has an antiviral activity against porcine reproductive and respiratory syndrome virus (PRRSV), infectious gastroenteritis virus (TGEV), seneca virus A (SVA), and encephalomyocarditis virus (EMCV) strongly.⁷⁸

SARS-CoV-2 AND COVID-19

Coronavirus family members (CoVs), which cause respiratory infections of human beings and other mammals, represent an enveloped and large group of single-stranded RNA viruses. Since the genome of these viruses can be directly translated into viral proteins by the host cell's ribosomes, they are defined as positive-sense viruses (viral mRNA). It is known that CoVs belong to the Nidovirales family including Roniviridae, Arteriviridae, Coronaviridae and Mesoniviridae main four members.⁷⁹⁻⁸¹ Phylogenetic analysis disclosed that the viral genome ranging from 27–32 kb has 29,903 nucleotides and shows 89.1% nucle-

otide similarity with SARS-like coronaviruses.^{82,83} The viral ORF1ab gene encodes the envelope glycoprotein or spike protein (S), membrane (M) protein, nucleocapsid (N) protein, and envelope (E) proteins, which are defined as structural proteins.⁸⁴ Follow-up of clinical reports reveals that coronavirus members, SARS-CoV, SARS-CoV-2 and MERSCoV, show the potential to infect mammals, causing very serious symptoms. The SARS-CoV-2 virus is the seventh in this regard.⁸⁵ SARS-CoV-2 was raised in December 2019 in China from an unidentified zoonotic source, which might be generated from pangolin or bat coronaviruses. The disease is characterized by hypoxemic respiratory failure, which requires air-ventilation support at the severe stages of the disease. COVID-19 patients can also show asymptomatic or mild symptoms of the disease such as fever, dry cough, fatigue, lung failure mainly acute respiratory distress syndrome (ARDS), and cytokine storm at the very late stage of the disease. Until now, different candidates have been suggested for therapy including novel or repurposed antibodies, anti-inflammatory drugs, and corticosteroids. WHO guidelines that were published in September 2020 recommend the systemic corticosteroids rather than no corticosteroids for the treatment of patients with severe and critical COVID-19 who required mechanical ventilation.⁸⁶ On the other hand, the same guideline also indicated not to use corticosteroids in the treatment of patients with non-severe which was based on certainty evidence. Conversely, guidelines from the Infectious Diseases Society of America that were published in April 2020 issued a weak recommendation against corticosteroids, except for patients with COVID-19 and ARDS treated in the context of a clinical trial.⁸⁶

While early observational data from China suggested a potential mortality benefit of corticosteroids in COVID-19, previous studies of corticosteroids in other viral pneumonia, especially severe acute respiratory syndrome (SARS) and the Middle East respiratory syndrome (MERS), found an association with delayed viral clearance and reinforced concerns that corticosteroids may impair host response to SARS-CoV-2. Furthermore, a meta-analysis of observational studies suggested increased mortality with corticosteroid treatment in influenza pneumonia. As the COVID-19 pandemic spread across the world, clinicians struggled to weigh the potential benefits of corticosteroids against the many potential harms associated with these drugs. Despite being overwhelmed with critically ill patients, multiple clinical trial groups around the world launched high-quality RCTs of corticosteroids for severe COVID-19. Additionally, recognizing the urgency of aggregating data from these trials to guide management, the World Health Organization (WHO) coordinated a prospective meta-analysis of these ongoing RCTs (PROSPERO CRD42020197242). The clinical trial groups agreed to share data, even before acceptance of their trial data for primary publication. With a press released on June 16, 2020, reporting the results of the UK-based RECOVERY trial, the existing approach for treating and studying patients with COVID-19 underwent a major change. In this large open-label randomized trial enrolling 6425 patients (2104 randomized to receive dexamethasone and 4321 randomized to receive usual care), treatment with dexamethasone (6 mg/d for 10 days) reduced mortality by one-third in patients receiving mechanical

ventilation (29.3% vs 41.4%, respectively; rate ratio, 0.64 [95% CI, 0.51-0.81]) and by one-fifth in patients receiving supplemental oxygen (23.3% vs 26.2%, respectively; 0.82 [95% CI, 0.72-0.94]) compared with usual care alone.¹³ However, there was no benefit among patients not receiving respiratory support (1.19 [95% CI, 0.91-1.55]), and the possibility of harm could not be excluded.⁸⁶

Potential Effects of Plant Steroids as Natural Therapeutics Against Coronaviruses

With the lack of a specific/successful drug in the treatment of SARS-CoV-2 infection, most researchers focus on the antiviral effects of phytomedicines against SARS coronaviruses simultaneously. In the process defined as the term 'molecular farming', plants' secondary metabolites are preferred in the production of alternative drugs and recombinant vaccines, since they exhibit fewer side effects compared to allopathic drugs.^{87,88}

Drug design studies are mainly focused on S protein RBD (receptor binding region) region of SARS-CoV-2, which targets angiotensin converting enzyme 2 (ACE2) receptors for the initial attachment process.^{89,90} Understanding the lipid rafts and lipid-activated molecular troughs in the plasma membrane areas that support the entry of the SARS-CoV-2 virus into the host cell (endocytosis) and infection suggests valuable clues to the emergence of powerful antiviral strategies.⁹¹ Recent studies on lipophilic naturally derived phytosterols emphasize that these substances interact with lipid rafts, destabilize the structure of membrane cholesterol and impress biochemical activities associated with lipid rafts.⁹² Experimental research on *Lycoris radiata* has shown that four herbal steroidal alkaloids derived from this plant may be candidates for the development of anti-SARS-CoV therapeutics.⁹³ In an *in vitro* study targeting the anti-coronaviral activity and mechanisms of action of saicosaponins (A, B2, C and D), it has been proven that saicosaponin B2 significantly inhibits human coronavirus 229E infection by interfering with viral absorption and replication processes.⁹⁴ Research results examining the antiviral properties and therapeutic potential of cardenolides and bufadienolides against SARS-CoV-2, HCoV-229E and HCoV-OC43 belonging to the coronavirus family report that these steroid compounds are the remarkable inhibitors of selected coronaviruses.⁹⁵

Very recent comprehensive *in silico* analysis on the interaction between various plant secondary metabolites (4,704 ligands) and the four structural target proteins of SARS-CoV-2 reveals in detail the possible physicochemical, bioavailability, and binding energy scores during interaction. The virtual screening and molecular docking results demonstrate that the steroidal lactones, triterpene sterols, coagulins, steroidal saponins, triterpene glycosides, triterpene saponins and steroidal glycoalkaloids were similar, showing lower and favorable binding energy scores against the target amino acid residues (AARs) of the S protein.⁹⁶ Molecular Simulation approach using the University of Bristol Insertion Engine (BUDE) recommends steroids, vitamins, retinoids as ligands that bind the free fatty acid pocket of SARS-CoV-2 S Protein.⁹⁷

The pathophysiology of COVID-19 is complex as various signaling pathways and molecules are involved in the inflammatory state. SARS-CoV-2 initiates pro-IL-1 secretion, inflammation and IL-1 beta hyper activation and up-regulation by binding to toll-like receptors. This hyperactive pro-inflammatory response is described as Cytokine Storm (CS). The severity of the infection that has occurred is marked by the levels of circulating cytokines and chemokines (monocyte chemotactic protein1, IL-2, IL-6, IFN γ , TNF). CS is known to cause severe fibrosis induction in the lung tissues of infected patients, which can result in death unfortunately.^{98,99} Obtained results of *in vitro* and *in vivo* studies focusing on the anti-inflammatory properties of Cucurbitacin against viral infections prove that Cucurbitacin II B decreased the expression of IFN γ , TNF α and IL-6. In addition, Cucurbitacin E and R, respectively, downregulate IL-1 and TNF α expression by suppressing NF-kB translocation and the STAT3 signaling pathways.^{100,101}

CONCLUSION

Viral diseases are very common in spreading, like in the case of COVID-19, and become pandemic. To treat or prevent viral infections is a challenge since it is usually impossible to control the replicative cycle of the virus without interfering the host one, or to estimate the mutations that it will carry after each replicative cycle. Plant steroids are divided in to several subclasses, and some of them have been shown as good candidates to treat viral infections since they have few side effects to the host. The exemplary studies described above indicate that plant steroid family members have a high potential for clinical efficacy, especially against SARS-CoV-2 infection. It is thought that the pharmacological manipulation of these compounds may show promising results in the course of the pandemic including COVID-19.

Peer Review: Externally peer-reviewed.

Author Contributions: Conception/Design of Study- P.O.Y.; Data Analysis/Interpretation- P.O.Y.; Drafting Manuscript- P.O.Y., E.D.A., L.M., P.U.O., A.C.G.; Critical Revision of Manuscript- P.O.Y., E.D.A., L.M., P.U.O., A.C.G.; Final Approval and Accountability- P.O.Y., E.D.A., L.M., P.U.O., A.C.G.

Conflict of Interest: Authors declared no conflict of interest.

Financial Disclosure: Authors declared no financial support.

REFERENCES

1. Trautwein EA, McKay S. The role of specific components of a plant-based diet in management of dyslipidemia and the impact on cardiovascular risk. *Nutrients*. 2020; 12(9), 71-91.
2. Kreis W, Müller-Uri F. Biochemistry of sterols, cardiac glycosides, brassinosteroids, phytoecdysteroids and steroid saponins. *Annu Rev Plant Biol*. 2010;304-363.
3. Beg S, Hasan H, Hussain MS et al. Systematic review of herbals as potential anti-inflammatory agents: Recent advances, current clinical status and future perspectives. *Pharmacogn Rev*. 2011;5(10):120-137.

4. Gunaherath GMK, Gunatilaka AAL. Plant steroids: occurrence, biological significance and their analysis. *Encyc of Anal Chem*. 2014;1–26.
5. Li Y, Li J, Zhou K et al. A review on phytochemistry and pharmacology of cortex periplocae. *Molecules*. 2016;21(12):2702-2718.
6. Bhandari J, Muhammad B, Thapa P et al. Study of phytochemical, anti-microbial, anti-oxidant, and anti-cancer properties of *Allium wallichii*. *BMC Complement Altern Med*. 2017;17(1):102.
7. Moosavi B, Liu S, Wang N et al. The anti-fungal β -sitosterol targets the yeast oxysterol-binding protein Osh4. *Pest Management Science*, 2019; 76(2):4–11.
8. Yang L, He J. Traditional uses, phytochemistry, pharmacology and toxicological aspects of the genus *Hosta* (Liliaceae): A comprehensive review. *J Ethnopharmacol*. 2021; (265):113-123.
9. Grove MD, Spencer GF, Rohwedder WK et al. Brassinolide, a plant growth-promoting steroid isolated from *Brassica napus* pollen. *Nature*, 1979;281(5728):216–217.
10. Yokota T, Takahashi N. Chemistry, physiology and agricultural application of brassinolide and related steroids. *Proceed in Life Sci*. 1986;129–138.
11. Kour J, Kohli SK, Khanna K et al. Brassinosteroid signaling, crosstalk and, physiological functions in plants under heavy metal stress. *Front Plant Sci*. 2021;24(12):608-061.
12. Khamsuk O, Sonjaroon W, Suwanwong S et al. Effects of 24-epibrassinolide and the synthetic brassinosteroid mimic on chili pepper under drought. *Acta Physiol. Plant*. 2018;(40):106-118.
13. Sondhi N, Bhardwaj R, Kaur S et al. Isolation of 24-epibrassinolide from leaves of *Aegle marmelos* and evaluation of its antigenotoxicity employing *Allium cepa* chromosomal aberration assay. *Plant Growth Reg*. 2007;54(3):217–224.
14. Carange J, Longpre F, Daoust B et al. 24-epibrassinolide, a phyto-steroid from the brassinosteroid family, protects dopaminergic cells against MPP-induced oxidative stress and apoptosis. *J Toxicol*. 2011;392859.
15. Clouse SD, Sasse, JM. Brassinosteroids: Essential regulators of plant growth and development. *Ann. Rev. Plant Physiol. Plant Mol. Biol*. 1998;(49): 427-451.
16. Singh I, Shono M. Physiological and molecular effects of 24-epibrassinolide, a brassinosteroid on thermotolerance of tomato. *Plant Grow Reg*. 2005; 47(2–3): 111-119.
17. Coskun D, Obakan P, Arisan ED et al. Epibrassinolide alters PI3K/ MAPK signaling axis via activating Foxo3a-induced mitochondria-mediated apoptosis in colon cancer cells. *Exp Cell Res*. 2015; 15;338(1):10-21.
18. Ramirez JA, Michelini FM, Galagovsky LR, Berra A, Alche LE. Antiangiogenic brassinosteroid compounds. U.S. Patent 2013088400, 17 November 2015.
19. Zhabinskii VN, Khripach NB, Khripach VA. Steroid plant hormones: effects outside plant kingdom. *Steroids*. 2015;(97):87-97.
20. Kaur Kohli S, Bhardwaj A, Bhardwaj V, Sharma A, Kalia N, Landi M, et al. Therapeutic potential of brassinosteroids in biomedical and clinical research. *Biomolecules*. 2020;10(4):572.
21. Puschett JB, Agunanne E, Uddin MN. Emerging role of the bufadienolides in cardiovascular and kidney diseases. *Am J Kidney Dis*. 2010;56(2):359-370.
22. Tang HJ, Ruan LJ, Tian HY, et al. Novel stereoselective bufadienolides reveal new insights into the requirements for Na(+), K(+)-ATPase inhibition by cardiosteric steroids. *Sci Rep*. 2016;6:29155.
23. Gao H, Popescu R, Kopp B, Wang Z. Bufadienolides and their anti-tumor activity. *Nat Prod Rep*. 2011;28(5):953-969.
24. Li H, Cao X, Chen X, et al. Bufadienolides induce apoptosis and autophagy by inhibiting the AKT signaling pathway in melanoma A-375 cells. *Mol Med Rep*. 2019;20(3):2347-2354.
25. Kaushik U, Aeri V, Mir SR. Cucurbitacins-An insight into medicinal leads from nature. *Pharmacogn Rev*. 2015;9(17):12-18.
26. Oi T, Asanuma K, Matsumine A, et al. STAT3 inhibitor, cucurbitacin I, is a novel therapeutic agent for osteosarcoma. *Int J Oncol*. 2016; 49(6):2275-2284.
27. Chen X, Bao J, Guo J, et al. Biological activities and potential molecular targets of cucurbitacins: a focus on cancer. *Anticancer Drugs*. 2012; 23(8):777-787.
28. Dinan L. Phytoecdysteroids: biological aspects. *Phytochemistry*. 2001;57(3):325-339.
29. Chadin I, Volodin V, Whiting P, Shirshova T, Kolegova N, Dinan L. Ecdysteroid content and distribution in plants of genus *Potamogeton*. *Biochem Syst and Ecol*. 2003; 31(4): 407-415.
30. Lafont R, Dinan L. Practical uses for ecdysteroids in mammals including humans: an update. *J Insect Sci*. 2003;3:7.
31. Festucci-Buselli RA, Contim LA, Barbosa LCA, Stuart J, Otoni WC. Biosynthesis and potential functions of the ecdysteroid 20-hydroxyecdysone-a review. *Botany*. 2008;86(9): 978-987.
32. Shakhmurova GA, Mamadalieva NZ, Zhanibekov AA, Khushbaktova ZA, Syrov VN. Effect of total ecdysteroid preparation from *Silene viridiflora* on the immune state of experimental animals under normal and secondary immunodeficiency conditions. *Pharm Chem J*. 2012;46 (4), 222–224.
33. Ishola, IO, Ochieng, CO, Olayemi SO, Jimoh MO, Lawal SM. Potential of novel phytoecdysteroids isolated from *Vitex doniana* in the treatment depression: Involvement of monoaminergic systems. *Pharmacol Biochem Behav*. 2014;127, 90–100.
34. Kregiel D, Berlowska J, Witonska I, et al. Saponin-based, biological-active surfactants from plants. In application and characterization of surfactants. Edited by Reza Najjar InTech. doi: 10.5772/65591, 2017.
35. Marrelli M, Conforti F, Araniti F, et al. Effects of saponins on lipid metabolism: a review of potential health benefits in the treatment of obesity. *Molecules*. 2016;21(10),1404.
36. Du JR, Long FY, Chen C. Research progress on natural triterpenoid saponins in the chemoprevention and chemotherapy of cancer. *Enzymes*. 2014;(36):95-130.
37. Sarkar FH, Li Y, Wang Z, et al. Cellular signaling perturbation by natural products. *Cell Signal*. 2009;21(11):1541-1547.
38. Haridas V, Higuchi M, Jayatilake GS, et al. Avicins: triterpenoid saponins from *Acacia victoriae* (Benth) induce apoptosis by mitochondrial perturbation. *Proc Natl Acad Sci USA*. 2001;98(10):5821-5826.
39. Patlolla JM, Raju J, Swamy MV, et al. Beta-escin inhibits colonic aberrant crypt foci formation in rats and regulates the cell cycle growth by inducing p21(waf1/cip1) in colon cancer cells. *Mol Cancer Ther*. 2006;5(6):1459-1466.
40. Fukumura M, Ando H, Hirai Y et al. Achyranthoside H methyl ester, a novel oleanolic acid saponin derivative from *Achyranthes fauriei* roots, induces apoptosis in human breast cancer MCF-7 and MDA-MB-453 cells via a caspase activation pathway. *J Nat Med*. 2009;63(2):181-188.
41. Ooh KF, Ong HC, Wong FC, et al. High performance liquid chromatography profiling of health-promoting phytochemicals and evaluation of antioxidant, anti-lipoxygenase, iron chelating and anti-glucosidase activities of wetland macrophytes. *Pharmacogn Mag*. 2014;10(3):443-455.
42. Dey P, Kundu A, Chakraborty HJ, et al. Therapeutic value of steroidal alkaloids in cancer: Current trends and future perspectives. *Int J Cancer*. 2019;145(7):1731-1744.
43. Dirsch VM, Müller IM, Eichhorst ST, et al. Cephalostatin 1 selectively triggers the release of Smac/DIABLO and subsequent apoptosis

- that is characterized by an increased density of the mitochondrial matrix. *Cancer Res.* 2003;63(24):8869-8876.
44. Moser BR. Review of cytotoxic cephalostatins and ritterazines: isolation and synthesis. *J Nat Prod.* 2008;71(3):487-491.
 45. Lacour TG, Guo C, Bhandaru S, et al. Interphylal Product Splicing: The First Total Syntheses of Cephalostatin 1, the North Hemisphere of Ritterazine G, and the Highly Active Hybrid Analogue, Ritterostatin GN1N1. *J Am Chem Soc.* 1998; 120(4), 692-707.
 46. Torres MC, das Chagas L Pinto F, et al. Antiophidic solanidane steroidal alkaloids from *Solanum campaniforme*. *J Nat Prod.* 2011;74(10):2168-2173.
 47. Wang K, Sasaki T, Li W, et al. Two novel steroidal alkaloid glycosides from the seeds of *Lycium barbarum*. *Chem Biodivers.* 2011;8(12):2277-2284.
 48. Jan NU, Ahmad B, Ali S, et al. Steroidal alkaloids as an emerging therapeutic alternative for investigation of their immunosuppressive and hepatoprotective potential. *Front Pharmacol.* 2017;(8):114.
 49. Antony ML, Lee J, Hahm ER, et al. Growth arrest by the antitumor steroidal lactone withaferin A in human breast cancer cells is associated with down-regulation and covalent binding at cysteine 303 of β -tubulin. *J Biol Chem.* 2014;289(3):1852-1865.
 50. Zhang H, Samadi AK, Gallagher RJ, et al. Cytotoxic withanolide constituents of *Physalis longifolia*. *J Nat Prod.* 2011;74(12):2532-2544.
 51. Zhang H, Bazzill J, Gallagher RJ, et al. Antiproliferative withanolides from *Datura wrightii*. *J Nat Prod.* 2013;76(3):445-449.
 52. Grogan PT, Sleder KD, Samadi AK, et al. Cytotoxicity of withaferin A in glioblastomas involves induction of an oxidative stress-mediated heat shock response while altering Akt/mTOR and MAPK signaling pathways. *Invest New Drugs.* 2013;31(3):545-557.
 53. Srigley CT, Haile EA. Quantification of plant sterols/stanols in foods and dietary supplements containing added phytosterols. *J Food Comp Anal.* 2015;(40):163-176.
 54. Miras-Moreno B, Sabater-Jara AB, Pedreno MA, et al. Bioactivity of phytosterols and their production in plant in vitro cultures. *J Agric Food Chem.* 2016; 64(38):7049-7058.
 55. Salehi-Sahlabadi A, Varkaneh HK, Shahdadian F, et al. Effects of Phytosterols supplementation on blood glucose, glycosylated hemoglobin (HbA1c) and insulin levels in humans: a systematic review and meta-analysis of randomized controlled trials. *J Diabetes Metab. Disord.* 2020;19(1):625-632.
 56. Moreau RA, Nystrom L, Whitaker BD, et al. Phytosterols and their derivatives: Structural diversity, distribution, metabolism, analysis, and health-promoting uses. *Prog Lipid Res.* 2018;(70):35-61.
 57. Hannan MA, Sohag AAM, Dash R, et al. Phytosterols of marine algae: Insights into the potential health benefits and molecular pharmacology. *Phytochemistry.* 2020;(69):153201.
 58. Vezza T, Canet F, de Maranon AM, et al. Phytosterols: Nutritional health players in the management of obesity and its related disorders. *Antioxidants (Basel).* 2020;9(12):1266.
 59. Liao PC, Lai MH, Hsu KP, et al. Identification of β -Sitosterol as in vitro anti-inflammatory constituent in *Moringa oleifera*. *J Agric Food Chem.* 2018;66(41):10748-10759.
 60. Turnbaugh PJ, Backhed F, Fulton L, et al. Diet-induced obesity is linked to marked but reversible alterations in the mouse distal gut microbiome. *Cell Host Microbe.* 2008;3(4):213-223.
 61. Martínez I, Perdicaro DJ, Brown AW, et al. Diet-induced alterations of host cholesterol metabolism are likely to affect the gut microbiota composition in hamsters. *Appl Environ Microbiol.* 2013;79(2):516-524.
 62. Castilla V, Ramirez J, Coto CE. Plant and animal steroids a new hope to search for antiviral agents. *Curr Med Chem.* 2010;17(18):1858-1873.
 63. Denaro M, Smeriglio A, Barreca D, et al. Antiviral activity of plants and their isolated bioactive compounds: An update. *Phytother Res.* 2020;34(4):742-768.
 64. Alkhatib A, Tsang C, Tiss A, et al. Functional foods and lifestyle approaches for diabetes prevention and management. *Nutrients.* 2017;9(12):1310.
 65. Parvez MK, Alam P, Arbab AH, et al. Analysis of antioxidative and antiviral biomarkers β -amyirin, β -sitosterol, lupeol, ursolic acid in *Guiera senegalensis* leaves extract by validated HPTLC methods. *Saudi Pharm. J.* 2018;26(5):685-693.
 66. Zhou BX, Li J, Liang XL, et al. β -sitosterol ameliorates influenza A virus-induced proinflammatory response and acute lung injury in mice by disrupting the cross-talk between RIG-I and IFN/STAT signaling. *Acta Pharmacol Sin.* 2020;41(9):1178-1196.
 67. Yuan M, Jiang Z, Bi G, et al. Pattern-recognition receptors are required for NLR-mediated plant immunity. *Nature.* 2021;592(7852):105-109.
 68. Calil IP, Fontes EPB. Plant immunity against viruses: antiviral immune receptors in focus. *Ann Bot.* 2017;119(5):711-723.
 69. Wachsmann MB, Lopez EM, Ramirez JA, et al. Antiviral effect of brassinosteroids against herpes virus and arenaviruses. *Antivir Chem Chemother.* 2000;11(1):71-77.
 70. Nakashita H, Yasuda M, Nitta T, et al. Brassinosteroid functions in a broad range of disease resistance in tobacco and rice. *Plant J.* 2003; 33(5):887-898.
 71. Yang H, Gou X, He K, et al. BAK1 and BKK1 in *Arabidopsis thaliana* confer reduced susceptibility to turnip crinkle virus. *Eur J Plant Pathol.* 2010; 127(1),149-156.
 72. Zhang DW, Deng XG, Fu FQ, et al. Induction of plant virus defense response by brassinosteroids and brassinosteroid signaling in *Arabidopsis thaliana*. *Planta.* 2015; 241(4):875-885.
 73. Shamsabadipour S, Ghanadian M, Saedi H, et al. Triterpenes and steroids from *Euphorbia denticulata* Lam. with anti-herpes simplex virus activity. *Iran J Pharm Res.* 2013;12(4):759-767.
 74. Supratman U, Fujita T, Akiyama K, et al. Insecticidal compounds from *Kalanchoe daigremontiana* x *tubiflora*. *Phytochemistry.* 2001;58(2):311-314.
 75. Munakarmi S, Chand L, Shin HB, et al. Anticancer effects of *Poncirus fructus* on hepatocellular carcinoma through regulation of apoptosis, migration, and invasion. *Oncol Rep.* 2020;44(6):2537-2546.
 76. Liu Y, Yang H, Guo Q, et al. Cucurbitacin E Inhibits Huh7 Hepatoma Carcinoma Cell Proliferation and Metastasis via Suppressing MAPKs and JAK/STAT3 Pathways. *Molecules.* 2020;25(3):560.
 77. Troost B, Mulder LM, Dioso-Toro M, et al. Tomatidine, a natural steroidal alkaloid shows antiviral activity towards chikungunya virus in vitro. *Sci Rep.* 2020;10(1):6364.
 78. Wang P, Bai J, Liu X, et al. Tomatidine inhibits porcine epidemic diarrhea virus replication by targeting 3CL protease. *Vet Res.* 2020;51(1):136.
 79. Brian DA, Baric RS. Coronavirus genome structure and replication. *Curr Top Microbiol Immunol.* 2005;(287):1-30.
 80. Fehr AR, Perlman S. Coronaviruses: an overview of their replication and pathogenesis. *Methods Mol Biol.* 2015;(1282):1-23.
 81. Mahmood N, Nasir SB, Hefferon K. Plant-Based Drugs and Vaccines for COVID-19. *Vaccines (Basel).* 2020;9(1):15.
 82. Dent SD, Xia D, Wastling JM, et al. The proteome of the infectious bronchitis virus Beau-R virion. *J Gen Virol.* 2015;96(12):3499-3506.
 83. Wu F, Zhao S, Yu B, et al. A new coronavirus associated with human respiratory disease in China. *Nature.* 2020;579(7798):265-269.
 84. Tang X, Wu C, Li X, et al. On the origin and continuing evolution of SARS-CoV-2. *Natl Sci Rev.* 2020;7(6):1012-1023.
 85. Corman VM, Muth D, Niemeyer D, et al. Hosts and Sources of Endemic Human Coronaviruses. *Adv Virus Res.* 2018;(100):163-188.

86. van Paassen J, Vos JS, Hoekstra EM, et al. Corticosteroid use in COVID-19 patients: a systematic review and meta-analysis on clinical outcomes. *Crit Care*. 2020;24(1):696.
87. Fischer R, Buyel JF. Molecular farming-The slope of enlightenment. *Biotechnol. Adv*. 2020;(40):107519.
88. Capell T, Twyman RM, Armario-Najera V, et al. Potential applications of plant biotechnology against SARS-CoV-2. *Trends Plant Sci*. 2020 25(7):635-643.
89. Shang J, Ye G, Shi K, et al. Structural basis of receptor recognition by SARS-CoV-2. *Nature*. 2020; 581(7807):221-224.
90. Bhuiyan FR, Howlader S, Raihan T, et al. Plants metabolites: Possibility of natural therapeutics against the COVID-19 pandemic. *Front Med (Lausanne)*. 2020;(7):444.
91. Fernandez-Oliva A, Ortega-Gonzalez P, Risco C. Targeting host lipid flows: Exploring new antiviral and antibiotic strategies. *Cell Microbiol*. 2019;21(3):e12996.
92. Baglivo M, Baronio M, Natalini G, et al. Natural small molecules as inhibitors of coronavirus lipid-dependent attachment to host cells: a possible strategy for reducing SARS-COV-2 infectivity? *Acta Biomed*. 2020;91(1):161-164.
93. Li SY, Chen C, Zhang HQ, et al. Identification of natural compounds with antiviral activities against SARS-associated coronavirus. *Antiviral Res*. 2005;67(1):18-23
94. Cheng PW, Ng LT, Chiang LC, et al. Antiviral effects of saikosaponins on human coronavirus 229E in vitro. *Clin Exp Pharmacol Physiol*. 2006;33(7):612-616.
95. Yang CW, Lee YZ, Hsu HY, et al. Inhibition of SARS-CoV-2 by highly potent broad-spectrum anti-coronaviral tylophorine-based derivatives. *Front Pharmacol*. 2020;(11):606097.
96. Puttaswamy H, Gowtham HG, Ojha MD, et al. In silico studies evidenced the role of structurally diverse plant secondary metabolites in reducing SARS-CoV-2 pathogenesis. *Sci Rep*. 2020;10(1):20584.
97. Shoemark DK, Colenso CK, Toelzer C, et al. Molecular simulations suggest vitamins, retinoids and steroids as ligands of the free fatty acid pocket of the SARS-CoV-2 spike protein. *Angew Chem Int Ed Engl*. 2021; 60(13):7098-8010.
98. Alka H, Akhilesh A, Archana S, et al. Cucurbitacin: As a candidate against cytokine storm in severe COVID-19 infection. *Int J Res Pharm Sci*. 2020;11(1): 928-930.
99. Conti P, Ronconi G, Caraffa A, et al. Induction of pro-inflammatory cytokines (IL-1 and IL-6) and lung inflammation by Coronavirus-19 (COVI-19 or SARS-CoV-2): anti-inflammatory strategies. *J Biol Regul Homeost Agents*. 2020;34(2):327-331.
100. Qiao J, Xu LH, He J, et al. Cucurbitacin E exhibits anti-inflammatory effect in RAW 264.7 cells via suppression of NF-κB nuclear translocation. *Inflamm Res*. 2013;62(5):461-469.
101. Escandell JM, Recio MC, Manez S, et al. Cucurbitacin R reduces the inflammation and bone damage associated with adjuvant arthritis in lewis rats by suppression of tumor necrosis factor- α in T lymphocytes and macrophages. *J Pharmacol Exp Ther*. 2007;320(2):581-590.