


# **Book of abstracts**



## **PHOTONICA2021**

VIII International School and Conference on Photonics

& HEMMAGINERO workshop

23 - 27 August 2021,

Belgrade, Serbia

*Editors*

Mihailo Rabasović, Marina Lekić and Aleksandar Krmpot

Institute of Physics Belgrade, Serbia

Belgrade, 2021

ABSTRACTS OF TUTORIAL, KEYNOTE, INVITED LECTURES,  
PROGRESS REPORTS AND CONTRIBUTED PAPERS

of

VIII International School and Conference on Photonics  
**PHOTONICA2021**

23 - 27 August 2021

Belgrade Serbia

*Editors*

Mihailo Rabasović, Marina Lekić and Aleksandar Krmpot

*Publisher*

Institute of Physics Belgrade  
Pregrevica 118  
11080 Belgrade, Serbia

*Printed by*

Serbian Academy of Sciences and Arts

*Number of copies*

200

ISBN 978-86-82441-53-3

CIP - Каталогизација у публикацији - Народна библиотека Србије, Београд

535(048)

621.37/.39:535(048)

621.37/.39:535]:61(048)

66.017/.018(048)

INTERNATIONAL School and Conference on Photonic (8; 2021; Beograd)

Book of abstracts / VIII International School and Conference on Photonics PHOTONICA2021 & HEMMAGINERO workshop, 23 - 27 August 2021, Belgrade, Serbia; editors Mihailo Rabasović, Marina Lekić and Aleksandar Krmpot. - Belgrade: Institute of Physics, 2021 (Belgrade: SASA). - V, 192 str.: ilustr.; 30 cm

Tiraž 200. - Bibliografija uz većinu apstrakata. - Registar.

ISBN 978-86-82441-53-3

1. Hemmaginero Workshop (2021; Beograd)

а) Оптика -- Апстракти б) Оптички материјали -- Апстракти в) Оптоелектроника -- Апстракти г)  
Оптоелектроника -- Биомедицина -- Апстракти д) Телекомуникације -- Апстракти

COBISS.SR-ID 44290057

## Modeling of intersubband transitions in ZnO/ZnMgO Coupled Quantum Wells

A. Atić<sup>1,2</sup>, J. Radovanović<sup>2</sup>, N. Vuković<sup>2</sup>

<sup>1</sup>Vinča Institute of Nuclear Sciences, Belgrade Serbia

<sup>2</sup>School of Electrical Engineering,

University of Belgrade, Serbia

e-mail:atic@vin.bg.ac.rs

In recent years ZnO has become a popular semiconductor with many potential applications in infra-red and THz optical devices owing to a wide direct bandgap (3.4 eV) in combination with relatively high exciton binding energy (60 meV) [1]-[2]. In this work, we model the electronic structure of coupled oxide-semiconductor quantum wells by numerically solving the system of coupled Schrödinger-Poisson equations self-consistently (Fig. 1). We compare the obtained results with the recent experimental data[3] and analyze how the variation of the layers' thicknesses affects the energy states. In addition, we examine the influence of doping to assess the differences between single well and two wells' cases, for the purpose of designing more complex multi-well optical system in the future.




Fig 1. Calculated conduction band of a double well ZnO/ZnMgO structure, together with wavefunctions squared, for a structure with 0.5 nm thick barrier at room temperature (300 K) with uniform layer doping  $n \approx 10 \times 10^{18} \text{ cm}^{-3}$  [3].

### REFERENCES

- [1] M. Belmoubarik, K. Ohtani, and H. Ohno, "Intersubband transitions in ZnO multiple quantum wells," *Appl. Phys. Lett.*, 2008.
- [2] N. Le Biavan *et al.*, "Homoepitaxy of non-polar ZnO/(Zn,Mg)O multi-quantum wells: From a precise growth control to the observation of intersubband transitions," *Appl. Phys. Lett.*, 2017.
- [3] B. Meng *et al.*, "Observation of Intersubband Absorption in ZnO Coupled Quantum Wells," *Phys. Rev. Appl.*, 2019.