

University of Massachusetts Medical School

eScholarship@UMMS

Open Access Articles

Open Access Publications by UMMS Authors

2009-03-18

In vitro and in vivo models of acute alcohol exposure

Angela Dolganiuc

University of Massachusetts Medical School

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: <https://escholarship.umassmed.edu/oapubs>



Part of the [Life Sciences Commons](#), and the [Medicine and Health Sciences Commons](#)

Repository Citation

Dolganiuc A, Szabo G. (2009). In vitro and in vivo models of acute alcohol exposure. Open Access Articles. <https://doi.org/10.3748/wjg.15.1168>. Retrieved from <https://escholarship.umassmed.edu/oapubs/2218>

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Open Access Articles by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

TOPIC HIGHLIGHT

Natalia A Osna, MD, PhD, Series Editor

In vitro and *in vivo* models of acute alcohol exposure

Angela Dolganiuc, Gyongyi Szabo

Angela Dolganiuc, Gyongyi Szabo, Department of Medicine, University of Massachusetts Medical School, 364 Plantation Street, Worcester, MA 01605-2324, United States

Author contributions: Dolganiuc A and Szabo G performed the literature search, analyzed the data and wrote the paper.

Supported by Grants AA016571 (AD) and AA014372 (GS) from NIAAAA (in part)

Correspondence to: Angela Dolganiuc, MD, PhD, Department of Medicine, University of Massachusetts Medical School, 364 Plantation Street, Worcester, MA 01605,

United States. angela.dolganiuc@umassmed.edu

Telephone: +1-508-8565955 Fax: +1-508-8564770

Received: December 17, 2008 Revised: January 5, 2009

Accepted: January 12, 2009

Published online: March 14, 2009

Abstract

Alcohol abuse is a global problem due to the financial burden on society and the healthcare system. While the harmful health effects of chronic alcohol abuse are well established, more recent data suggest that acute alcohol consumption also affects human wellbeing. Thus, there is a need for research models in order to fully understand the effect of acute alcohol abuse on different body systems and organs. The present manuscript summarizes the interdisciplinary advantages and disadvantages of currently available human and non-human models of acute alcohol abuse, and identifies their suitability for biomedical research.

© 2009 The WJG Press and Baishideng. All rights reserved.

Key words: Models; Acute alcohol abuse; Human; Non-human; Progress

Peer reviewers: Giovanni Addolorato, MD, Institute of Internal medicine, L.go Gemelli 8, Rome 00168, Italy; James Neuberger, Professor, Liver Unit, Queen Elizabeth Hospital, Birmingham B15 2TH, United Kingdom

Dolganiuc A, Szabo G. *In vitro* and *in vivo* models of acute alcohol exposure. *World J Gastroenterol* 2009; 15(10): 1168-1177 Available from: URL: <http://www.wjnet.com/1007-9327/15/1168.asp> DOI: <http://dx.doi.org/10.3748/wjg.15.1168>

INTRODUCTION

Alcohol abuse is widely spread around the globe^[1-9].

Alcohol is the third leading cause of preventable death in the United States and the third leading cause of healthy years lost to death and disability in developed nations^[9]. Humans use and abuse alcohol acutely or chronically, when alcohol consumption is frequent and dependence has developed^[10]. Although significant progress was made in the area of alcohol research during the last decades, the pathogenesis of alcohol use and abuse is not fully understood. Further, most research was focused on alcoholism, which is an advanced stage of alcohol abuse, involving chronic alcohol consumption, alcohol dependence and severe health and social consequences^[1-13]. Thus, research models are emergent in order to detail what drives human desire to consume alcohol, how the body responds to alcohol, and most important, what are the beneficial and harmful effects of acute alcohol consumption on the human body.

ACUTE ALCOHOL ABUSE (AAA): HOW BIG THE PROBLEM REALLY IS?

In the USA, a “drink” is defined as an equivalent of 14 g alcohol, which equals roughly 1 shot [1.25 oz of 40% (80-proof) liquor], 1 (12 oz) beer (4.2 mL/L, Ethanol), or 1 (4 oz) glass of wine (12 mL/L, Ethanol)^[14]. In other countries, the alcohol content of a serving is measured in “units”. One unit (about 25 mL of a 40% 80-proof liquor) contains 7.9 g of pure ethanol^[8,15]. However, in many countries the “standard drink” is used to quantify alcohol intake. More importantly, the standard drink varies significantly from country to country, from 10 mL (7.9 g) of alcohol in the UK to as high as 25 mL (19.75 g) in Japan^[16]. Current use includes at least one drink in the past 30 d; binge drinking is defined as five or more drinks on the same occasion within 2 h at least once in the past 30 d; and heavy use is defined as five or more drinks on the same occasion on at least 5 different days in the past 30 d^[11-13]. The 0.08% blood alcohol level (BAL) is the legal limit for most states in the US and it is achieved with consumption of five or more drinks for an adult male and four or more drinks for an adult female^[11-13].

Traditionally medical research focuses on the mechanisms of chronic alcohol abuse; this is due to the significant financial burden that society encountered primarily from chronic alcohol abusers^[1-13]. However, more recently acute alcohol abuse has emerged as a

Table 1 The characteristics of *in vitro* and *in vivo* models of AAA

AAA model	Advantages	Disadvantages	Area of research
<i>In vitro</i>	Low cost Technically easy to perform Large number of experimental groups Pure cell populations Single cell type or multi-cell type co-culture Strictly controlled settings yielding reproducible results	Limited alcohol metabolism Limited complexity at cellular and tissue levels Limited areas of research, not suitable for behavioral and social studies.	Behavioral and biomedical
<i>In vivo</i>	Availability of physiological routes of alcohol administration Complex interactions of all bodily organs and systems, including complex metabolism Controlled settings, caloric and composition controls Indications to individual and population variability	Ethical concerns High cost Limited information about the effect on one separate cell population.	All areas of research including biomedical, behavioral and social.

social problem^[17]. The National Survey on Drug Use and Health (NSDUH) estimated that in the USA about 4.4 million persons had used alcohol for the first time in 2004, which lead to about 12000 “new recruits” per day; this was significantly greater than in 2002 (3.9 million) and 2003 (4.1 million). Most (86.9%) of the 4.4 million recent alcohol initiates were younger than 21 years of age at the time of encounter. More than one fifth (22.%) of people age 12 or older participated in binge drinking at least once in the 30 d prior to the survey in 2004^[12]. Acute alcohol intake in the form of binge drinking in 2004 was highest for the 18- to 25-year-old age group compared with other age groups, with the peak rate occurring at age 21^[1,5-7,11-13]. The statistics also show that illness and death among young adults primarily result from lifestyle choices and behaviors, including excessive alcohol use^[18,19].

AAA: BIOMEDICAL IMPACT

The known biological effects of AAA include those of the central nervous system (CNS) and non-CNS origin. Alcohol use is characterized by symptoms of CNS intoxication, impaired brain activity, poor motor coordination, and behavioral changes^[20,21]. AAA leads to impaired CNS activity due to alcohol's effect on synthesis^[22], release^[23] and signaling^[23,24] of neurotransmitters, including serotonin^[25,26], glutamate^[27], GABA^[28], endocannabinoids^[29,30] and their receptors. AAA causes damage and functional impairment of the gastrointestinal (GI) tract, including luminal GI^[31-38], liver^[39-53], and pancreas^[56-62]; it also affects the protein, carbohydrate, and fat metabolism^[58,63-66]. AAA leads to insufficient immune system responses to infections; such deficiency was observed both in organ-specific^[67-69] and systemic infections^[70-72]. Acute alcohol intoxication impairs the ability of the host to counteract hemorrhagic shock^[73], augments corticosteroid release^[74] and delays wound healing^[75-78], thus contributing to higher morbidity and mortality^[79] and prolonged recovery from trauma^[80]. The pathogenesis of AAA effects on human health is not fully understood.

MODELS OF AAA

Research of acute alcohol consumption/abuse is entirely

based on models, due to their advantage of controlled settings. Currently there are *in vitro* and *in vivo* models of AAA; their characteristics are defined in Table 1. In contrast to chronic alcohol abuse, the research of AAA has not benefited from population studies due to recall bias^[81-84].

One important feature of AAA models is the definition of biologically meaningful levels of alcohol, either *in vitro* or *in vivo*, and their relationship to blood alcohol levels (BAL) in humans. This is an important requirement of the research models of AAA, because BAL can be detected as soon as minimal amounts of alcohol are ingested^[85], however measurable affects of alcohol on physiology and/or behavior is established at 0.08% or above this level, with individual variations depending on the species, metabolic particularities, age, gender and genetic background^[86-97]. It is also important to identify that AAA models differ by their route of alcohol delivery to achieve alcohol intoxication, some of them being physiological, such as oral administration, while others being non-physiological, when ethanol is administered by parenteral routes. Nevertheless, current research shows that the BAL levels, rather than the route of alcohol administration play a major role in the establishment of the biological effects of alcohol^[97].

Thus, optimal AAA models should fulfill several criteria: (1) Define the length of alcohol exposure. *In vitro* the length of acute alcohol treatment is variable in diverse published experimental settings and range from seconds to hours; it is currently accepted that treatment with alcohol for up to 24 h is considered as an acute setting^[98-106]. *In vivo* the consumption of alcohol in one setting implies that the entire dose of alcohol is consumed at once, while a ‘binge’ is defined by NIAAA as an excessive pattern of alcohol drinking that produces BAL greater than 0.08% within a 2-h period and may, or may not, be associated with dependence^[11,12,17,18]. Thus any model using consumption of biologically active amounts of alcohol within 2 h is considered an acceptable model of AAA^[81,107-121]. (2) Establish an exposure to an accurate concentration of ethanol. For *in vitro* studies the 10-100 mmol/L ethanol range is considered physiological, with 25 mmol/L ethanol being close to 0.08% BAL achieved *in vivo* after 4-5 drink equivalents^[7,11,12,98-106]. For the *in vivo* studies an 0.08% BAL or above this level yields

signs of intoxication and it is employed in the majority of biomedical studies^[107-121]. (3) Recruit individuals who are currently not and never have been alcohol abusers for *in vivo* studies and employ alcohol-naïve primary cells or cell lines for *in vitro* studies. Alcohol use habits of the study participants are usually determined by questionnaires^[122]. Among most frequently used questionnaires are those that incorporate the AUDIT and CAGE tests^[123-125]; the study parameters are usually permissive for males who had alcohol use of fewer than nine drinks/week, females < 6 drinks/week.

IN VITRO AAA MODEL

The *in vitro* alcohol treatment model is based on supplementation of culture media with pure alcohol, usually 200-proof ethanol. Currently supplementation of cell culture with a wide variety of alcohol concentrations, ranging from 1 to 500 mmol/L, is reported in the bio-medical literature. One of the major concerns with the *in vitro* alcohol treatment using concentrations above 100 mmol/L is the direct cytotoxic effect of alcohol on cells^[40,100]. At lower concentrations (< 100 mmol/L), alcohol changes the redox status of the cells and alters intercellular junctions^[33,126], increases the membrane fluidity of cells^[127-129] and affects the composition of lipid rafts^[106,130,131], all of which may contribute to alcohol-mediated increase in transcellular and paracellular permeability^[132,133] and thus affect cell function^[106,130-134]. Alcohol also affects the expression of adhesion molecules^[135], which may be a concern when using adherent cell types due to possible cell detachment. Additional concerns arise from the possibility of modified *ex vivo* function of some primary cells, including hepatocytes, stellate cells and their precursors, due to limited *ex vivo* environment compared to *in vivo* conditions^[136-138].

From a technical point, the acute alcohol exposure of cells *in vitro* may be hampered by alcohol evaporation. To avoid the fluctuation of alcohol concentration due to evaporation, investigators used settings where ethanol was added into the culture media and the cell culture plates were maintained for the entire duration of stimulation in a microclimate chamber at 37°C with gas mixture and an alcohol atmosphere^[139]. For example, if the desired alcohol concentration in the cell culture is 25 mmol/L, a Petri dish with 2 × the alcohol amount (50 mmol/L) was placed on the bottom of the chamber to ensure the saturation of the gas in the chamber; such conditions maintain the initial alcohol concentration ± 15% over a 24 h period^[139]. However, depending on the scientific question of the study, the declining alcohol levels *in vitro* may be desired to mimic the alcohol elimination *in vivo*; in these situations the *in vitro* experiments are disadvantaged by the absence/limitation of alcohol metabolism^[76,134].

The *in vitro* AAA model offers the possibility of primary *in vitro* exposure of alcohol-naïve cells to alcohol alone or its combinations with diverse pharmacological or naturally-derived substances^[24,28,31,35,36,42,55,68,71,72,96,103],

but also the investigation of the effects of *in vivo* exposure to alcohol followed by *ex vivo* exposure to other stimulants^[110,113,115] or *vice versa*. One other main characteristic of the *in vitro* AAA model is its simplicity, often considered as an advantage or disadvantage depending on the research goal. Most of the *in vitro* research involves culture of a single cell type^[134,139-142] or co-culture of several cell types^[143]; while such an approach brings forward the differential effect of alcohol on pure cell populations, and/or their intercellular interaction; it lacks the systemic alcohol metabolism and intercellular interactions. More recently significant efforts were invested in establishment of more complex *in vitro* systems, such as culture of cells in three dimensional systems^[100], organ slices^[144] or organ explants^[145]; while such systems are informative in the setting of chronic alcohol exposure to date there is no report of their use as an AAA model.

IN VIVO AAA MODELS

The *in vivo* models of AAA are more informative compared to the *in vitro* model due to complex physiological impact of alcohol on all bodily organs and systems, but also due to the availability of systemic alcohol metabolism. Currently there are human and non-human models of AAA, and the later include use of invertebrates^[146-147] and vertebrates^[21,25,37,44,46,47,53,65,72,86,93,94,98,104,110,111]. The invertebrate models (*Drosophila melanogaster*^[146,147], *Caenorhabditis elegans*^[105]) and those using lower vertebrates (*Zebra fish Danio rerio*)^[98] are invaluable for research of the effect of alcohol on behavior, development and maintenance of memory, and on basic signaling mechanisms. These models offer the advantage of a well-defined genetic background, high-turnover rate of experiments due to short life cycle and relatively low-cost; in light of these advantages they constitute an excellent resource for research of signaling pathways and are highly desirable for their drug-screening capacity. On the downside, significant differences in the structure and function of organs and systems compared to humans limit the informative value of invertebrate and lower vertebrate models of AAA.

The vertebrate models are preferred to those using invertebrates due to closer resemblance of their bodily structure, function, and metabolism to that of humans. However, because of intrinsic differences between humans and other vertebrates, no single non-human model is perfect since none of the models can represent all features of the complex human trait, such as motivation for social occasional or binge alcohol consumption, development of alcohol dependence and establishment of the impact on health. Further, the controlled setting of research models may not be completely satisfactory for psychology and social research, since they may not fully reproduce the social component, the motivation and the spontaneity of alcohol abuse. However, research models are invaluable for the understanding of the effects of alcohol and its

mechanisms of action on hardwired bodily systems, including the brain and all other organs and systems.

HUMAN MODELS OF AAA

Human alcohol intake in the experimental setting is the best available model of AAA, because it offers the advantage of the physiological route of alcohol consumption, the possibility to investigate human pathobiology and the availability of relatively large amounts of physiological bodily fluids for research. The disadvantages of human models of AAA include ethical concerns related to potential harmful health effects due to excessive or repeated intoxication, and the theoretical possibility of development of dependence or tolerance even after a one-time drinking session. Published models of human AAA are based on consumption of alcoholic beverages containing either distilled ethanol or wine; these models are physiological, as they involve alcohol drinking, and achieve a biologically meaningful BAL^[87,92,107,110-113,115]. The majority of the reported *in vivo* models of human AAA strictly control for the amount of alcohol based on the constant volume of alcohol per kg body weight, includes placebo-treated age and gender matched controls. However, most of these studies design the consumption of the alcohol beverage during a 2 h period of time^[92,107,110-113], which based on recent NIAAA and NSDUH classification qualifies as binge drinking^[11-13,17]. Thus the major disadvantage of the human models of AAA is that they (1) do not clearly distinguish between one-time and the binge alcohol consumption pattern, and, (2) for ethical reasons, do not allow longer binge sessions which are often observed in real-life and account for the majority of the heavy alcohol intake in young adults^[5,7,11-13,17-19].

To fulfill the requirement for an AAA model, the human studies usually include nonalcoholic individuals, who did not drink any alcohol at least 24 h prior to the study. Depending on the study design, some AAA human models require that the study participants did not take any medication, while others accept individuals taking moderate doses of anti-hypertensive medication and oral contraceptives^[107,110]. The study participants are usually required to abstain from food for at least 6 h before alcohol consumption and are allowed free access to water and a light meal before or shortly after the study^[107]. The human model of AAA is currently used for research in physiology^[86,92,111,122], hematology^[107,128] and immunology^[110,113,115].

CONSUMPTION OF DISTILLED ETHANOL MODEL

In this model the study individuals drink distilled alcohol (usually 80-proof vodka) in amounts of about 0.5-0.6 g/kg body weight, which is an equivalent of about 2 mL vodka/kg body weight in a standardized total volume of liquid (300-450 mL of water or orange juice)^[92,107,110-113].

CONSUMPTION OF NON-DISTILLED ETHANOL MODEL

In this model the study individuals drink wine to an equivalent of a pre-determined amount of ethanol/kg BW (for example, Fehr *et al.*^[107] reported use of 4.36 mL of red wine/kg of body weight as an equivalent of 0.5 g ethanol/kg BW to lead to a peak BAL of about 15 mmol/L in the first 2 h), while the control individuals are exposed to the same volume of fluid by mouth (usually water) per individual in a randomized way. The major disadvantage of this model is the use of controlled volumes of liquids that are not matched by calorie intake or by composition, which is technically challenging to achieve due to restricted availability of equivalent alcohol-free compounds. To bypass the bias concern some studies employ a cross-over approach, where each subject serves as its own control and repeats the study at least 2 wk after the first experiment with either alcohol or placebo consumption according to the cross-over design^[107].

NON-HUMAN AAA MODELS

Among non-human vertebrates commonly involved in alcohol research are primates^[90,91,148], pigs^[104,120], dogs^[114,121], mice^[70,72,74,86,89,96,109,118,119,141], rats^[88,94,108,149,150] and rabbits^[132]. The rodent AAA models (mice and rats) are used most frequently due to their relatively well-defined genetic background and the availability of diverse genetic traits, including those coding for high or low alcohol consumption^[88,89,96,109]. Most non-human AAA models currently in use^[93,95] examine relative oral self-administration from a bottle containing alcohol versus one^[86,94,108] or multiple bottles^[119] containing water (preference drinking) or administration of alcohol against the will, either by physiological (by mouth using gavage)^[54,71,72] or by non-physiological (parenteral)^[67,68] routes. Voluntary consumption of alcohol may be an optimal animal model of AAA, due to physiological route and pattern of alcohol consumption. However, in the self-administration models it is not clear when or if the animals drink to pharmacologically significant levels because the drinking is episodic and often occurs over a 24-h period. Nevertheless, these models are invaluable for research of neurobiology of acute intoxication with alcohol and for establishment of mechanisms of addiction. The AAA models using administration of alcohol against-the-will bypass all the above-mentioned inconveniences of AAA models using voluntary consumption. Alcohol administered either by physiological (by mouth using gavage) or by non-physiological (parenteral) routes yields comparable physiological effects on the central nervous system and on organs/systems that are not affected directly by the route of alcohol administration, such as muscle and brain^[97]. However, administration of alcohol per os is more physiological compared to administration via parenteral routes, yields meaningful levels of BAL and shows signs of acute alcohol intoxication^[54,71,72,132,149].

Table 2 The effect of acute alcohol abuse on GI system

GI segment	Effect of acute alcohol exposure
Oral cavity	Unknown
Esophagus	Low concentrations of alcohol (up to 5%) cause alterations in ion transports and affect the barrier function Concentrations of alcohol of 10% and above cause injury of mucosa Co-carcinogenic potency
Stomach	Motor dysfunction: decrease in lower esophageal sphincter pressure and amplitude Motor dysfunction: Inhibition of gastric emptying Mucosal damage, impaired barrier function, increased epithelial permeability Pro-inflammatory reaction: decreased gastric blood flow, vascular damage, polymorphonuclear neutrophils (PMN) dependent- and independent-mucosal damage Aggravation of <i>H pylori</i> infection
Intestine	Disruption of barrier function Epithelial apoptosis Enhanced bioavailability of some alcohol-soluble drugs and impaired absorption of key nutrients Increased paracellular intestinal permeability to toxins
Liver	Hepatocytes: Amplification of Fas-mediated hepatocyte death Generation of oxidative stress Hepatic mitochondrial dysfunction Increased free iron levels Imbalanced fatty acid metabolism Inhibition of IFN- α -induced antiviral response towards hepatotropic viruses including hepatitis C virus favors hepatitis C virus replication expression Induced histone H3 acetylation leading to increased gene expression in the liver Limited hepatic protein synthesis Arrest of liver regeneration early after partial hepatectomy and suppression of hepatic stimulator substance (HSS) activity by induction of liver cell cycle arrest Kupffer cells: Suppressed LPS-mediated priming for enhanced CC-chemokine release in vitro; up-regulated expression of CC-chemokine mRNA; primed the KC for enhanced RANTES release Desensitized HIV-1 gp120-induced CC-chemokine production Downregulates HIV-1 glycoprotein 120-induced KC and RANTES production Regulates production of reactive oxygen species Modulate the tolerance to LPS Stellate cells: Imbalanced redox potential owed to increased generation of reactive oxygen species upon GSH depletion
Pancreas	Stimulates islet blood flow, amplifies insulin secretion, induces hypoglycemia Lower baseline amylase output of acinar pancreatic cells, with the difference being significantly exacerbated by cerulein stimulation Interference with release of oxidized proteins in acinar cells Predisposes the pancreas to postprandial cholinergic stimulation that triggers cellular events leading to pancreatic inflammation Impaired apical exocytosis and redirected exocytosis to less efficient basolateral plasma membrane sites Augments elevated-[Ca ²⁺]-induced trypsin activation in pancreatic acinar zymogen granules, leading to premature activation of trypsin and tissue damage.

Among disadvantages of administration of alcohol per os are technical challenges, time consumption and high cost of the procedures. In contrast, alcohol administration by parenteral routes is relatively easy to perform technically and offers controlled settings (time, amount); on the downside, they may be less suitable for research of the effects of alcohol on organs/systems that are affected directly by the routes of alcohol administration. In this context, administration of alcohol by intraperitoneal route may be less suitable for research using peritoneal macrophages, or even liver and intestines, compared to other administration routes, such as intravenous or enteral. Further, some parenteral methods of alcohol administration are preferred over others, owing to differences in the level of technical difficulty of the procedure and the effect of alcohol on different cell types. For example, alcohol administration by intravenous route is known to affect the erythrocytes when present in high concentrations^[128]. Thus alcohol administration by intravenous route is currently

limited to creating an acute alcohol exposure during treatment of alcohol withdrawal symptoms^[151], while administration by intraperitoneal injections is widely preferred in research settings.

Similar to human AAA models, the non-human *in vivo* models employ either distilled alcohol^[53,67-72,74,89,90,96,109,116,119,121,132,135,141,152] or alcohol-containing beverages, such as wine^[152,153] and beer^[46]; the control groups are usually treated with alcohol-free caloric and composition equivalents. The vertebrate AAA models are widely used in research of biomedical effects of AAA, including brain^[23-30,116], gastrointestinal^[38-44,46-48,64-66,154], vascular^[73,153], muscle^[97] and immune^[68-72,74] systems.

THE PARTICULARITIES OF AAA MODELS FOR RESEARCH IN GASTROENTEROLOGY

In contrast to the abundance of the literature about the

effects of chronic alcohol abuse on the gastrointestinal system, research of the effects of acute alcohol abuse on the gastrointestinal (GI) tract is limited to certain cell types, as outlined in Table 2.

Currently the state of scientific knowledge suggests a tight interplay between organs and systems. The GI system is dependent on blood circulation and systemic availability of metabolites, is closely governed by both the central and the autonomous nervous system^[155,156] and contains a hallmark of resident and recruited immune cells^[157,158]. Thus, it is conceivable that the direct effects of alcohol on either of these systems will indirectly affect the function of the gastrointestinal system; this area is currently largely unexplored.

From a technical point, the GI research may take advantage of both *in vitro* and *in vivo* AAA models; however some *in vivo* models, such as those using parenteral administration of alcohol by the intraperitoneal route, may be less suitable due to the non-physiological direct contact between high concentrations of alcohol and GI tissues.

Alcohol use/abuse is associated with acute life-threatening conditions, including acute alcoholic hepatitis^[45] or acute pancreatitis^[159]. The majority of these patients report acute alcohol abuse, which is often overlapping with withdrawal from or even discontinued chronic alcohol abuse, or it follows an episode of binge drinking^[45,159]. As such, it is difficult to associate these diseases with the single-occasion AAA, yet they do not fit into the classic chronic alcohol abuse picture. This category of alcohol abuse, defined as “acute-on-chronic”, is in need of modeling for GI research.

In prospective, we currently lack in-depth knowledge in regards to the effects of acute alcohol abuse on different segments of the luminal GI tract, on liver functions, and on pancreas, including its endocrine and exocrine functions. Further, we do not know if acute alcohol consumption affects the GI stem cells and/or is involved in development of GI-derived tumors.

REFERENCES

- 1 **Caetano R**, Vaeth PA, Ramisetty-Mikler S, Rodriguez LA. The Hispanic americans baseline alcohol survey: alcoholic beverage preference across Hispanic national groups. *Alcohol Clin Exp Res* 2009; **33**: 150-159
- 2 **Davydov MI**, Zaridze D G, Lazarev AF, Maksimovich DM, Igitov VI, Boroda AM, Khvastiuk MG. [Analysis of mortality in Russian population] *Vestn Ross Akad Med Nauk* 2007; **(7)**: 17-27
- 3 **Goldfinger TM**. Beyond the French paradox: the impact of moderate beverage alcohol and wine consumption in the prevention of cardiovascular disease. *Cardiol Clin* 2003; **21**: 449-457
- 4 **Kröber HL**. Psychiatric criteria of legal responsibility after the consumption of alcohol: the German situation. *Eur Addict Res* 1998; **4**: 107-112
- 5 **Hao W**, Su Z, Liu B, Zhang K, Yang H, Chen S, Biao M, Cui C. Drinking and drinking patterns and health status in the general population of five areas of China. *Alcohol Alcohol* 2004; **39**: 43-52
- 6 **Cochrane J**, Chen H, Conigrave KM, Hao W. Alcohol use in China. *Alcohol Alcohol* 2003; **38**: 537-542
- 7 **Kypri K**, Paschall MJ, Langley J, Baxter J, Cashell-Smith M, Bourdeau B. Drinking and alcohol-related harm among New Zealand university students: findings from a national Web-based survey. *Alcohol Clin Exp Res* 2009; **33**: 307-314
- 8 **National Health and Medical Research Council of Australia**. Available from: URL: <http://www.nhmrc.gov.au/publications/synopses/ds9syn.htm>
- 9 **National Institutes of Health, National Institute on Alcohol Abuse and Alcoholism**. Resource Guide-April 2005. Available from: URL: <http://www.niaaa.nih.gov>
- 10 **Chick J**, Erickson CK. Conference summary: Consensus Conference on Alcohol Dependence and the Role of Pharmacotherapy in its Treatment. *Alcohol Clin Exp Res* 1996; **20**: 391-402
- 11 **Results from the 2001 National Survey on Drug Use and Health: National Findings**. Available from: URL: <http://www.oas.samhsa.gov>
- 12 **Results from the 2004 National Survey on Drug Use and Health: National Findings**. Available from: URL: <http://www.oas.samhsa.gov/>
- 13 **National Institutes of Health, National Institute on Alcohol Abuse and Alcoholism**. Alcohol Alert. No. 37. July 1997. Available from URL: <http://pubs.niaaa.nih.gov/publications/aa37.htm>
- 14 **National Institutes of Health, National Institute on Alcohol Abuse and Alcoholism**. Available from: URL: http://pubs.niaaa.nih.gov/publications/Practitioner/PocketGuide/pocket_guide2.htm
- 15 **UK Department of Health, alcohol publications**. Available from: URL: http://www.dh.gov.uk/en/PublicHealth/Healthimprovement/Alcoholmisuse/DH_4001740
- 16 **Wikipedia**. Available from: URL: http://en.wikipedia.org/wiki/Standard_drink
- 17 **Naimi TS**, Brewer RD, Mokdad A, Denny C, Serdula MK, Marks JS. Binge drinking among US adults. *JAMA* 2003; **289**: 70-75
- 18 **Wechsler H**, Lee JE, Kuo M, Lee H. College binge drinking in the 1990s: a continuing problem. Results of the Harvard School of Public Health 1999 College Alcohol Study. *J Am Coll Health* 2000; **48**: 199-210
- 19 **Schulenberg J**, Maggs JL, Steinman KJ, Zucker RA. Development matters: Taking the long view on substance abuse etiology and intervention during adolescence. In: Monti PM, Colby SM, O'Leary TA, eds. Adolescents, alcohol, and substance abuse: Reaching teens through brief interventions. New York: Guilford Press, 2001: 19-57
- 20 **Gohlke JM**, Griffith WC, Faustman EM. Computational models of ethanol-induced neurodevelopmental toxicity across species: Implications for risk assessment. *Birth Defects Res B Dev Reprod Toxicol* 2008; **83**: 1-11
- 21 **Shapira Y**, Lam AM, Paez A, Artru AA, Laohaprasit V, Donato T. The influence of acute and chronic alcohol treatment on brain edema, cerebral infarct volume and neurological outcome following experimental head trauma in rats. *J Neurosurg Anesthesiol* 1997; **9**: 118-127
- 22 **Pietrzykowski AZ**, Friesen RM, Martin GE, Puig SI, Nowak CL, Wynne PM, Siegelmann HT, Treistman SN. Posttranscriptional regulation of BK channel splice variant stability by miR-9 underlies neuroadaptation to alcohol. *Neuron* 2008; **59**: 274-287
- 23 **Roberto M**, Treistman SN, Pietrzykowski AZ, Weiner J, Galindo R, Mameli M, Valenzuela F, Zhu PJ, Lovinger D, Zhang TA, Hendricson AH, Morrisett R, Siggins GR. Actions of acute and chronic ethanol on presynaptic terminals. *Alcohol Clin Exp Res* 2006; **30**: 222-232
- 24 **Wilkie MB**, Besheer J, Kelley SP, Kumar S, O'Buckley TK, Morrow AL, Hodge CW. Acute ethanol administration rapidly increases phosphorylation of conventional protein kinase C in specific mammalian brain regions in vivo. *Alcohol Clin Exp Res* 2007; **31**: 1259-1267
- 25 **LeMarquand D**, Pihl RO, Benkelfat C. Serotonin and alcohol intake, abuse, and dependence: clinical evidence.

- Biol Psychiatry* 1994; **36**: 326-337
- 26 **McBride WJ**, Murphy JM, Gatto GJ, Levy AD, Yoshimoto K, Lumeng L, Li TK. CNS mechanisms of alcohol self-administration. *Alcohol Alcohol Suppl* 1993; **2**: 463-467
 - 27 **Prosser RA**, Mangrum CA, Glass JD. Acute ethanol modulates glutamatergic and serotonergic phase shifts of the mouse circadian clock in vitro. *Neuroscience* 2008; **152**: 837-848
 - 28 **Wallner M**, Hanchar HJ, Olsen RW. Low-dose alcohol actions on alpha4beta3delta GABAA receptors are reversed by the behavioral alcohol antagonist Ro15-4513. *Proc Natl Acad Sci USA* 2006; **103**: 8540-8545
 - 29 **Perra S**, Pillolla G, Luchicchi A, Pistis M. Alcohol inhibits spontaneous activity of basolateral amygdala projection neurons in the rat: involvement of the endocannabinoid system. *Alcohol Clin Exp Res* 2008; **32**: 443-449
 - 30 **Basavarajappa BS**, Ninan I, Arancio O. Acute ethanol suppresses glutamatergic neurotransmission through endocannabinoids in hippocampal neurons. *J Neurochem* 2008; **107**: 1001-1013
 - 31 **Asai K**, Buurman WA, Reutelingsperger CP, Schutte B, Kaminishi M. Modular effects of estradiol on ethanol-induced apoptosis in human intestinal epithelial cells. *Scand J Gastroenterol* 2005; **40**: 326-335
 - 32 **Asai K**, Buurman WA, Reutelingsperger CP, Schutte B, Kaminishi M. Low concentrations of ethanol induce apoptosis in human intestinal cells. *Scand J Gastroenterol* 2003; **38**: 1154-1161
 - 33 **Banan A**, Fields JZ, Decker H, Zhang Y, Keshavarzian A. Nitric oxide and its metabolites mediate ethanol-induced microtubule disruption and intestinal barrier dysfunction. *J Pharmacol Exp Ther* 2000; **294**: 997-1008
 - 34 **Fisher SJ**, Lee IJ, Swaan PW, Eddington ND. Evaluation of the effect of ethanol's toxic metabolite acetaldehyde on the gastrointestinal oligopeptide transporter, PEPT1: in vitro and in vivo studies. *Alcohol Clin Exp Res* 2008; **32**: 162-170
 - 35 **Kokoska ER**, Smith GS, Deshpande Y, Rieckenberg CL, Miller TA. Adaptive cytoprotection induced by ethanol in human intestinal cells: role of prostaglandins and calcium homeostasis. *Ann Surg* 1998; **228**: 123-130
 - 36 **Landrier JF**, Malezet-Desmoulins C, Reboul E, Marie Lorec A, Josephe Amiot M, Borel P. Comparison of different vehicles to study the effect of tocopherols on gene expression in intestinal cells. *Free Radic Res* 2008; **42**: 523-530
 - 37 **Li X**, Rana SN, Schwacha MG, Chaudry IH, Choudhry MA. A novel role for IL-18 in corticosterone-mediated intestinal damage in a two-hit rodent model of alcohol intoxication and injury. *J Leukoc Biol* 2006; **80**: 367-375
 - 38 **Ma TY**, Nguyen D, Bui V, Nguyen H, Hoa N. Ethanol modulation of intestinal epithelial tight junction barrier. *Am J Physiol* 1999; **276**: G965-G974
 - 39 **Bailey SM**, Cunningham CC. Acute and chronic ethanol increases reactive oxygen species generation and decreases viability in fresh, isolated rat hepatocytes. *Hepatology* 1998; **28**: 1318-1326
 - 40 **Baker RC**, Kramer RE. Cytotoxicity of short-chain alcohols. *Annu Rev Pharmacol Toxicol* 1999; **39**: 127-150
 - 41 **Bautista AP**. Acute alcohol intoxication and endotoxemia desensitize HIV-1 gp120-induced CC-chemokine production by Kupffer cells. *Life Sci* 2001; **68**: 1939-1949
 - 42 **Bautista AP**. Acute ethanol binge followed by withdrawal regulates production of reactive oxygen species and cytokine-induced neutrophil chemoattractant and liver injury during reperfusion after hepatic ischemia. *Antioxid Redox Signal* 2002; **4**: 721-731
 - 43 **Bautista AP**, Wang E. Acute ethanol administration downregulates human immunodeficiency virus-1 glycoprotein 120-induced KC and RANTES production by murine Kupffer cells and splenocytes. *Life Sci* 2002; **71**: 371-382
 - 44 **Bukara M**, Bautista AP. Acute alcohol intoxication and gadolinium chloride attenuate endotoxin-induced release of CC chemokines in the rat. *Alcohol* 2000; **20**: 193-203
 - 45 **Ceccanti M**, Attili A, Balducci G, Attilia F, Giacomelli S, Rotondo C, Sasso GF, Xirouchakis E, Attilia ML. Acute alcoholic hepatitis. *J Clin Gastroenterol* 2006; **40**: 833-841
 - 46 **Degrace P**, Moindrot B, Mohamed I, Gresti J, Clouet P. Moderate consumption of beer reduces liver triglycerides and aortic cholesterol deposit in LDLr-/- apoB100/100 mice. *Atherosclerosis* 2006; **189**: 328-335
 - 47 **Enomoto N**, Ikejima K, Bradford B, Rivera C, Kono H, Brenner DA, Thurman RG. Alcohol causes both tolerance and sensitization of rat Kupffer cells via mechanisms dependent on endotoxin. *Gastroenterology* 1998; **115**: 443-451
 - 48 **Karinch AM**, Martin JH, Vary TC. Acute and chronic ethanol consumption differentially impact pathways limiting hepatic protein synthesis. *Am J Physiol Endocrinol Metab* 2008; **295**: E3-E9
 - 49 **Kondili VG**, Tzirogiannis KN, Androustos CD, Papadimas GK, Demonakou MD, Hereti RI, Manta GA, Kourentzi KT, Triantaphyllou MI, Panoutsopoulos GI. The hepatoprotective effect of hepatic stimulator substance (HSS) against liver regeneration arrest induced by acute ethanol intoxication. *Dig Dis Sci* 2005; **50**: 297-307
 - 50 **Nishitani Y**, Okazaki S, Imabayashi K, Katada R, Matsumoto H. Ethanol-induced JNK activation suppressed via active Akt in hepatocytes. *Nihon Arukoru Yakubutsu Igakkai Zasshi* 2008; **43**: 35-43
 - 51 **Novitskiy G**, Traore K, Wang L, Trush MA, Mezey E. Effects of ethanol and acetaldehyde on reactive oxygen species production in rat hepatic stellate cells. *Alcohol Clin Exp Res* 2006; **30**: 1429-1435
 - 52 **Park PH**, Lim RW, Shukla SD. Involvement of histone acetyltransferase (HAT) in ethanol-induced acetylation of histone H3 in hepatocytes: potential mechanism for gene expression. *Am J Physiol Gastrointest Liver Physiol* 2005; **289**: G1124-G1136
 - 53 **Wang X**, Cederbaum AI. Acute ethanol pretreatment increases FAS-mediated liver injury in mice: role of oxidative stress and CYP2E1-dependent and -independent pathways. *Free Radic Biol Med* 2007; **42**: 971-984
 - 54 **Wheeler MD**, Thurman RG. Up-regulation of CD14 in liver caused by acute ethanol involves oxidant-dependent AP-1 pathway. *J Biol Chem* 2003; **278**: 8435-8441
 - 55 **Yan SL**, Yin MC. Protective and alleviative effects from 4 cysteine-containing compounds on ethanol-induced acute liver injury through suppression of oxidation and inflammation. *J Food Sci* 2007; **72**: S511-S515
 - 56 **Cosen-Binker LI**, Lam PP, Binker MG, Gaisano HY. Alcohol-induced protein kinase Calpha phosphorylation of Munc18c in carbachol-stimulated acini causes basolateral exocytosis. *Gastroenterology* 2007; **132**: 1527-1545
 - 57 **Ding YX**, Yang K, Chin WC. Ethanol augments elevated-[Ca²⁺]_i induced trypsin activation in pancreatic acinar zymogen granules. *Biochem Biophys Res Commun* 2006; **350**: 593-597
 - 58 **Huang Z**, Sjöholm A. Ethanol acutely stimulates islet blood flow, amplifies insulin secretion, and induces hypoglycemia via nitric oxide and vagally mediated mechanisms. *Endocrinology* 2008; **149**: 232-236
 - 59 **Palmieri VO**, Grattagliano I, Palasciano G. Ethanol induces secretion of oxidized proteins by pancreatic acinar cells. *Cell Biol Toxicol* 2007; **23**: 459-464
 - 60 **Yang AL**, Vadavkar S, Singh G, Omary MB. Epidemiology of alcohol-related liver and pancreatic disease in the United States. *Arch Intern Med* 2008; **168**: 649-656
 - 61 **Andrzejewska A**, Dlugosz JW, Jurkowska G. The effect of antecedent acute ethanol ingestion on the pancreas ultrastructure in taurocholate pancreatitis in rats. *Exp Mol Pathol* 1998; **65**: 64-77
 - 62 **Dlugosz JW**, Wróblewski E, Poplawski C, Andrzejewska A, Gabryelewicz A. The effect of beta-thia-iminoprostacyclin in taurocholate acute pancreatitis in rats: the role of antecedent acute ethanol abuse. *Pancreas* 1997; **15**: 91-98
 - 63 **Olubadewo JO**, Spitzer JA. Immune response modulation

- in acutely ethanol-intoxicated, acutely diabetic male and female rats. *Alcohol* 2003; **31**: 137-147
- 64 **Ting JW**, Lauth WW. The effect of acute, chronic, and prenatal ethanol exposure on insulin sensitivity. *Pharmacol Ther* 2006; **111**: 346-373
- 65 **Tomie Furuya D**, Binsack R, Onishi ME, Monteiro Seraphim P, Fabres Machado U. Low ethanol consumption induces enhancement of insulin sensitivity in liver of normal rats. *Life Sci* 2005; **77**: 1813-1824
- 66 **Carrasco MP**, Jiménez-López JM, Segovia JL, Marco C. Effects of ethanol on the remodeling of neutral lipids and phospholipids in brain mitochondria and microsomes. *Neurochem Int* 2007; **50**: 858-865
- 67 **Happel KI**, Odden AR, Zhang P, Shellito JE, Bagby GJ, Nelson S. Acute alcohol intoxication suppresses the interleukin 23 response to *Klebsiella pneumoniae* infection. *Alcohol Clin Exp Res* 2006; **30**: 1200-1207
- 68 **Happel KI**, Rudner X, Quinton LJ, Movassaghi JL, Clark C, Odden AR, Zhang P, Bagby GJ, Nelson S, Shellito JE. Acute alcohol intoxication suppresses the pulmonary ELR-negative CXC chemokine response to lipopolysaccharide. *Alcohol* 2007; **41**: 325-333
- 69 **Zhang P**, Bagby GJ, Stoltz DA, Summer WR, Nelson S. Granulocyte colony-stimulating factor modulates the pulmonary host response to endotoxin in the absence and presence of acute ethanol intoxication. *J Infect Dis* 1999; **179**: 1441-1448
- 70 **Carson EJ**, Pruett SB. Development and characterization of a binge drinking model in mice for evaluation of the immunological effects of ethanol. *Alcohol Clin Exp Res* 1996; **20**: 132-138
- 71 **Bagby GJ**, Zhang P, Stoltz DA, Nelson S. Suppression of the granulocyte colony-stimulating factor response to *Escherichia coli* challenge by alcohol intoxication. *Alcohol Clin Exp Res* 1998; **22**: 1740-1745
- 72 **Pruett SB**, Zheng Q, Fan R, Matthews K, Schwab C. Ethanol suppresses cytokine responses induced through Toll-like receptors as well as innate resistance to *Escherichia coli* in a mouse model for binge drinking. *Alcohol* 2004; **33**: 147-155
- 73 **Greiffenstein P**, Mathis KW, Stouwe CV, Molina PE. Alcohol binge before trauma/hemorrhage impairs integrity of host defense mechanisms during recovery. *Alcohol Clin Exp Res* 2007; **31**: 704-715
- 74 **Choudhry MA**, Li X, Chaudry IH. A role for corticosterone in impaired intestinal immunity and barrier function in a rodent model of acute alcohol intoxication and burn injury. *J Neuroimmune Pharmacol* 2006; **1**: 428-434
- 75 **Radek KA**, Matthies AM, Burns AL, Heinrich SA, Kovacs EJ, DiPietro LA. Acute ethanol exposure impairs angiogenesis and the proliferative phase of wound healing. *Am J Physiol Heart Circ Physiol* 2005; **289**: H1084-H1090
- 76 **Radek KA**, Kovacs EJ, Gallo RL, DiPietro LA. Acute ethanol exposure disrupts VEGF receptor cell signaling in endothelial cells. *Am J Physiol Heart Circ Physiol* 2008; **295**: H174-H184
- 77 **Radek KA**, Kovacs EJ, DiPietro LA. Matrix proteolytic activity during wound healing: modulation by acute ethanol exposure. *Alcohol Clin Exp Res* 2007; **31**: 1045-1052
- 78 **Fitzgerald DJ**, Radek KA, Chaar M, Faunce DE, DiPietro LA, Kovacs EJ. Effects of acute ethanol exposure on the early inflammatory response after excisional injury. *Alcohol Clin Exp Res* 2007; **31**: 317-323
- 79 **Jones JD**, Barber B, Engrav L, Heimbach D. Alcohol use and burn injury. *J Burn Care Rehabil* 1991; **12**: 148-152
- 80 **Swenson JR**, Dimsdale JE, Rockwell E, Carroll W, Hansbrough J. Drug and alcohol abuse in patients with acute burn injuries. *Psychosomatics* 1991; **32**: 287-293
- 81 **Gmel G**, Daeppen JB. Recall bias for seven-day recall measurement of alcohol consumption among emergency department patients: implications for case-crossover designs. *J Stud Alcohol Drugs* 2007; **68**: 303-310
- 82 **Searles JS**, Perrine MW, Mundt JC, Helzer JE. Self-report of drinking using touch-tone telephone: extending the limits of reliable daily contact. *J Stud Alcohol* 1995; **56**: 375-382
- 83 **Embree BG**, Whitehead PC. Validity and reliability of self-reported drinking behavior: dealing with the problem of response bias. *J Stud Alcohol* 1993; **54**: 334-344
- 84 **Alanko T**, Poikolainen K. A statistical approach to an alcoholic drinking history. *Br J Addict* 1992; **87**: 755-766
- 85 **Lindberg L**, Brauer S, Wollmer P, Goldberg L, Jones AW, Olsson SG. Breath alcohol concentration determined with a new analyzer using free exhalation predicts almost precisely the arterial blood alcohol concentration. *Forensic Sci Int* 2007; **168**: 200-207
- 86 **Bachmanov AA**, Reed DR, Li X, Li S, Beauchamp GK, Tordoff MG. Voluntary ethanol consumption by mice: genome-wide analysis of quantitative trait loci and their interactions in a C57BL/6ByJ x 129P3/J F2 intercross. *Genome Res* 2002; **12**: 1257-1268
- 87 **Birley AJ**, James MR, Dickson PA, Montgomery GW, Heath AC, Whitfield JB, Martin NG. Association of the gastric alcohol dehydrogenase gene ADH7 with variation in alcohol metabolism. *Hum Mol Genet* 2008; **17**: 179-189
- 88 **Clark JW**, Fixaris MC, Belanger GV, Rosenwasser AM. Repeated light-dark phase shifts modulate voluntary ethanol intake in male and female high alcohol-drinking (HAD1) rats. *Alcohol Clin Exp Res* 2007; **31**: 1699-1706
- 89 **Grahame NJ**, Grose AM. Blood alcohol concentrations after scheduled access in high-alcohol-preferring mice. *Alcohol* 2003; **31**: 99-104
- 90 **Grant KA**, Johanson CE. Oral ethanol self-administration in free-feeding rhesus monkeys. *Alcohol Clin Exp Res* 1988; **12**: 780-784
- 91 **Grant KA**, Leng X, Green HL, Szeliga KT, Rogers LS, Gonzales SW. Drinking typography established by scheduled induction predicts chronic heavy drinking in a monkey model of ethanol self-administration. *Alcohol Clin Exp Res* 2008; **32**: 1824-1838
- 92 **Martin NG**, Perl J, Oakeshott JG, Gibson JB, Starmer GA, Wilks AV. A twin study of ethanol metabolism. *Behav Genet* 1985; **15**: 93-109
- 93 **Siegmund SV**, Haas S, Singer MV. Animal models and their results in gastrointestinal alcohol research. *Dig Dis* 2005; **23**: 181-194
- 94 **Simms JA**, Steensland P, Medina B, Abernathy KE, Chandler LJ, Wise R, Bartlett SE. Intermittent access to 20% ethanol induces high ethanol consumption in Long-Evans and Wistar rats. *Alcohol Clin Exp Res* 2008; **32**: 1816-1823
- 95 **Wolffgramm J**, Galli G, Thimm F, Heyne A. Animal models of addiction: models for therapeutic strategies? *J Neural Transm* 2000; **107**: 649-668
- 96 **Yang X**, Wang S, Rice KC, Munro CA, Wand GS. Restraint stress and ethanol consumption in two mouse strains. *Alcohol Clin Exp Res* 2008; **32**: 840-852
- 97 **Vary TC**, Frost RA, Lang CH. Acute alcohol intoxication increases atrogen-1 and MuRF1 mRNA without increasing proteolysis in skeletal muscle. *Am J Physiol Regul Integr Comp Physiol* 2008; **294**: R1777-R1789
- 98 **Gerlai R**, Lee V, Blaser R. Effects of acute and chronic ethanol exposure on the behavior of adult zebrafish (*Danio rerio*). *Pharmacol Biochem Behav* 2006; **85**: 752-761
- 99 **Lemos C**, Peters GJ, Jansen G, Martel F, Calhau C. Modulation of folate uptake in cultured human colon adenocarcinoma Caco-2 cells by dietary compounds. *Eur J Nutr* 2007; **46**: 329-336
- 100 **Li LN**, Margolis LB, Hoffman RM. Skin toxicity determined in vitro by three-dimensional, native-state histoculture. *Proc Natl Acad Sci USA* 1991; **88**: 1908-1912
- 101 **Rao RK**. Acetaldehyde-induced barrier disruption and paracellular permeability in Caco-2 cell monolayer. *Methods Mol Biol* 2008; **447**: 171-183
- 102 **Zhang T**, Li Y, Lai JP, Douglas SD, Metzger DS, O'Brien CP, Ho WZ. Alcohol potentiates hepatitis C virus replicon expression. *Hepatology* 2003; **38**: 57-65

- 103 **Tang Y**, Banan A, Forsyth CB, Fields JZ, Lau CK, Zhang LJ, Keshavarzian A. Effect of alcohol on miR-212 expression in intestinal epithelial cells and its potential role in alcoholic liver disease. *Alcohol Clin Exp Res* 2008; **32**: 355-364
- 104 **Cloutier S**, Skaer TL, Newberry RC. Consumption of alcohol by sows in a choice test. *Physiol Behav* 2006; **88**: 101-107
- 105 **Davies AG**, Pierce-Shimomura JT, Kim H, VanHoven MK, Thiele TR, Bonci A, Bargmann CI, McIntire SL. A central role of the BK potassium channel in behavioral responses to ethanol in *C. elegans*. *Cell* 2003; **115**: 655-666
- 106 **Dolganiuc A**, Bakis G, Kodys K, Mandrekar P, Szabo G. Acute ethanol treatment modulates Toll-like receptor-4 association with lipid rafts. *Alcohol Clin Exp Res* 2006; **30**: 76-85
- 107 **Fehr M**, Galliard-Grigioni KS, Reinhart WH. Influence of acute alcohol exposure on hemorheological parameters and platelet function in vivo and in vitro. *Clin Hemorheol Microcirc* 2008; **39**: 351-358
- 108 **Ji D**, Gilpin NW, Richardson HN, Rivier CL, Koob GF. Effects of naltrexone, duloxetine, and a corticotropin-releasing factor type 1 receptor antagonist on binge-like alcohol drinking in rats. *Behav Pharmacol* 2008; **19**: 1-12
- 109 **Kamdar NK**, Miller SA, Syed YM, Bhayana R, Gupta T, Rhodes JS. Acute effects of naltrexone and GBR 12909 on ethanol drinking-in-the-dark in C57BL/6J mice. *Psychopharmacology (Berl)* 2007; **192**: 207-217
- 110 **Mandrekar P**, Catalano D, White B, Szabo G. Moderate alcohol intake in humans attenuates monocyte inflammatory responses: inhibition of nuclear regulatory factor kappa B and induction of interleukin 10. *Alcohol Clin Exp Res* 2006; **30**: 135-139
- 111 **Martin NG**, Gibson JB, Oakeshott JG, Wilks AV, Starmer GA, Craig J, Perl J. A twin study of psychomotor performance during alcohol intoxication: early results. *Prog Clin Biol Res* 1981; **69**: 89-96
- 112 **Martin NG**, Oakeshott JG, Gibson JB, Starmer GA, Perl J, Wilks AV. A twin study of psychomotor and physiological responses to an acute dose of alcohol. *Behav Genet* 1985; **15**: 305-347
- 113 **Norkina O**, Dolganiuc A, Catalano D, Kodys K, Mandrekar P, Syed A, Efros M, Szabo G. Acute alcohol intake induces SOCS1 and SOCS3 and inhibits cytokine-induced STAT1 and STAT3 signaling in human monocytes. *Alcohol Clin Exp Res* 2008; **32**: 1565-1573
- 114 **Molina PE**, Jabbour K, Williams P, Abumrad NN. Effect of acute ethanol intoxication on gluco-regulation during prolonged insulin-induced hypoglycemia. *Am J Physiol* 1994; **267**: R1280-R1287
- 115 **Szabo G**, Mandrekar P, Dolganiuc A, Catalano D, Kodys K. Reduced alloreactive T-cell activation after alcohol intake is due to impaired monocyte accessory cell function and correlates with elevated IL-10, IL-13, and decreased IFN γ levels. *Alcohol Clin Exp Res* 2001; **25**: 1766-1772
- 116 **Szumliński KK**, Diab ME, Friedman R, Henze LM, Lominac KD, Bowers MS. Accumens neurochemical adaptations produced by binge-like alcohol consumption. *Psychopharmacology (Berl)* 2007; **190**: 415-431
- 117 **Sergent O**, Tomasi A, Ceccarelli D, Masini A, Nohl H, Cillard P, Cillard J, Vladimirov YA, Kozlov AV. Combination of iron overload plus ethanol and ischemia alone give rise to the same endogenous free iron pool. *Biometals* 2005; **18**: 567-575
- 118 **Soulat T**, Philippe C, Bal dit Sollier C, Brézillon C, Berge N, Teissedre PL, Callebert J, Rabot S, Drouet L. Wine constituents inhibit thrombosis but not atherogenesis in C57BL/6 apolipoprotein E-deficient mice. *Br J Nutr* 2006; **96**: 290-298
- 119 **Tordoff MG**, Bachmanov AA. Influence of the number of alcohol and water bottles on murine alcohol intake. *Alcohol Clin Exp Res* 2003; **27**: 600-606
- 120 **Villanueva J**, Chandler CJ, Shimasaki N, Tang AB, Nakamura M, Phinney SD, Halsted CH. Effects of ethanol feeding on liver, kidney and jejunal membranes of micro-pigs. *Hepatology* 1994; **19**: 1229-1240
- 121 **Zysset T**, Preisig R, Bircher J. Increased systemic availability of drugs during acute ethanol intoxication: studies with mephenytoin in the dog. *J Pharmacol Exp Ther* 1980; **213**: 173-178
- 122 **Kip MJ**, Neumann T, Jugel C, Kleinwaechter R, Weiss-Gerlach E, Guill MM, Spies CD. New strategies to detect alcohol use disorders in the preoperative assessment clinic of a German university hospital. *Anesthesiology* 2008; **109**: 171-179
- 123 **Ewing JA**. Detecting alcoholism. The CAGE questionnaire. *JAMA* 1984; **252**: 1905-1907
- 124 **Fleming MF**, Barry KL, MacDonald R. The alcohol use disorders identification test (AUDIT) in a college sample. *Int J Addict* 1991; **26**: 1173-1185
- 125 **Fleming MF**, Barry KL. A three-sample test of a masked alcohol screening questionnaire. *Alcohol Alcohol* 1991; **26**: 81-91
- 126 **Banan A**, Keshavarzian A, Zhang L, Shaikh M, Forsyth CB, Tang Y, Fields JZ. NF-kappaB activation as a key mechanism in ethanol-induced disruption of the F-actin cytoskeleton and monolayer barrier integrity in intestinal epithelium. *Alcohol* 2007; **41**: 447-460
- 127 **Moulin M**, Carpentier S, Levade T, Arrigo AP. Potential roles of membrane fluidity and ceramide in hyperthermia and alcohol stimulation of TRAIL apoptosis. *Apoptosis* 2007; **12**: 1703-1720
- 128 **Padmini E**, Sundari BT. Erythrocyte Glutathione Depletion Impairs Resistance to Haemolysis in Women Consuming Alcohol. *J Clin Biochem Nutr* 2008; **42**: 14-20
- 129 **Dickey AN**, Faller R. How alcohol chain-length and concentration modulate hydrogen bond formation in a lipid bilayer. *Biophys J* 2007; **92**: 2366-2376
- 130 **Nourissat P**, Travert M, Chevanne M, Tekpli X, Rebillard A, Le Moigne-Müller G, Rissel M, Cillard J, Dimanche-Boitrel MT, Lagadic-Gossmann D, Sergent O. Ethanol induces oxidative stress in primary rat hepatocytes through the early involvement of lipid raft clustering. *Hepatology* 2008; **47**: 59-70
- 131 **Dai Q**, Zhang J, Pruett SB. Ethanol alters cellular activation and CD14 partitioning in lipid rafts. *Biochem Biophys Res Commun* 2005; **332**: 37-42
- 132 **Bor S**, Caymaz-Bor C, Tobey NA, Abdounour-Nakhoul S, Marten E, Orlando RC. Effect of ethanol on the structure and function of rabbit esophageal epithelium. *Am J Physiol* 1998; **274**: G819-G826
- 133 **Catalioto RM**, Festa C, Triolo A, Altamura M, Maggi CA, Giuliani S. Differential effect of ethanol and hydrogen peroxide on barrier function and prostaglandin E2 release in differentiated Caco-2 cells: selective prevention by growth factors. *J Pharm Sci* 2009; **98**: 713-727
- 134 **Donohue TM**, Osna NA, Clemens DL. Recombinant Hep G2 cells that express alcohol dehydrogenase and cytochrome P450 2E1 as a model of ethanol-elicited cytotoxicity. *Int J Biochem Cell Biol* 2006; **38**: 92-101
- 135 **Sacanella E**, Estruch R. The effect of alcohol consumption on endothelial adhesion molecule expression. *Addict Biol* 2003; **8**: 371-378
- 136 **Michalopoulos GK**, Bowen WC, Zajac VF, Beer-Stolz D, Watkins S, Kostrubsky V, Strom SC. Morphogenetic events in mixed cultures of rat hepatocytes and nonparenchymal cells maintained in biological matrices in the presence of hepatocyte growth factor and epidermal growth factor. *Hepatology* 1999; **29**: 90-100
- 137 **Yang L**, Jung Y, Omenetti A, Witek RP, Choi S, Vandongen HM, Huang J, Alpini GD, Diehl AM. Fate-mapping evidence that hepatic stellate cells are epithelial progenitors in adult mouse livers. *Stem Cells* 2008; **26**: 2104-2113
- 138 **Sancho-Bru P**, Najimi M, Caruso M, Pawelyn K, Cantz T, Forbes SJ, Roskams T, Ott M, Gehling U, Sokal E, Verfaillie C,

- Muraca M. Stem and progenitor cells for liver repopulation: can we standardize the process from bench to bedside? *Gut* 2008; [Epub ahead of print]
- 139 **Szabo G**, Mandrekar P. Human monocytes, macrophages, and dendritic cells: alcohol treatment methods. *Methods Mol Biol* 2008; **447**: 113-124
- 140 **Karavitis J**, Murdoch EL, Gomez CR, Ramirez L, Kovacs EJ. Acute ethanol exposure attenuates pattern recognition receptor activated macrophage functions. *J Interferon Cytokine Res* 2008; **28**: 413-422
- 141 **Choudhry MA**, Ren X, Romero A, Kovacs EJ, Gamelli RL, Sayeed MM. Combined alcohol and burn injury differentially regulate P-38 and ERK activation in mesenteric lymph node T cell. *J Surg Res* 2004; **121**: 62-68
- 142 **Goral J**, Choudhry MA, Kovacs EJ. Acute ethanol exposure inhibits macrophage IL-6 production: role of p38 and ERK1/2 MAPK. *J Leukoc Biol* 2004; **75**: 553-559
- 143 **Dolganiuc A**, Kodys K, Kopasz A, Marshall C, Mandrekar P, Szabo G. Additive inhibition of dendritic cell allostimulatory capacity by alcohol and hepatitis C is not restored by DC maturation and involves abnormal IL-10 and IL-2 induction. *Alcohol Clin Exp Res* 2003; **27**: 1023-1031
- 144 **Klassen LW**, Thiele GM, Duryee MJ, Schaffert CS, DeVeney AL, Hunter CD, Olinga P, Tuma DJ. An in vitro method of alcoholic liver injury using precision-cut liver slices from rats. *Biochem Pharmacol* 2008; **76**: 426-436
- 145 **Pietrzykowski AZ**, Martin GE, Puig SI, Knott TK, Lemos JR, Treisman SN. Alcohol tolerance in large-conductance, calcium-activated potassium channels of CNS terminals is intrinsic and includes two components: decreased ethanol potentiation and decreased channel density. *J Neurosci* 2004; **24**: 8322-8332
- 146 **LaFerriere H**, Guarnieri DJ, Sitaraman D, Diegelmann S, Heberlein U, Zars T. Genetic dissociation of ethanol sensitivity and memory formation in *Drosophila melanogaster*. *Genetics* 2008; **178**: 1895-1902
- 147 **Morozova TV**, Anholt RR, Mackay TF. Phenotypic and transcriptional response to selection for alcohol sensitivity in *Drosophila melanogaster*. *Genome Biol* 2007; **8**: R231
- 148 **Bagby GJ**, Stoltz DA, Zhang P, Bohm RP Jr, Nelson S. Simian immunodeficiency virus, infection, alcohol, and host defense. *Alcohol Clin Exp Res* 1998; **22**: 193S-195S
- 149 **Tsukamoto H**, French SW, Benson N, Delgado G, Rao GA, Larkin EC, Largman C. Severe and progressive steatosis and focal necrosis in rat liver induced by continuous intragastric infusion of ethanol and low fat diet. *Hepatology* 1985; **5**: 224-232
- 150 **Tsukamoto H**, French SW, Reidelberger RD, Largman C. Cyclical pattern of blood alcohol levels during continuous intragastric ethanol infusion in rats. *Alcohol Clin Exp Res* 1985; **9**: 31-37
- 151 **Hodges B**, Mazur JE. Intravenous ethanol for the treatment of alcohol withdrawal syndrome in critically ill patients. *Pharmacotherapy* 2004; **24**: 1578-1585
- 152 **Arola L**, Roig R, Cascón E, Brunet MJ, Fornós N, Sabaté M, Raga X, Batista J, Salvadó MJ, Bladé C. Model for voluntary wine and alcohol consumption in rats. *Physiol Behav* 1997; **62**: 353-357
- 153 **Munday JS**, Thompson KG, James KA, Manktelow BW. The effect of moderate alcohol consumption as either red or white wine in the C57BL/6 mouse atherosclerosis model. *Coron Artery Dis* 1999; **10**: 97-102
- 154 **Kaiser JP**, Beier JJ, Zhang J, David Hoetker J, von Montfort C, Guo L, Zheng Y, Monia BP, Bhatnagar A, Arteel GE. PKCepsilon plays a causal role in acute ethanol-induced steatosis. *Arch Biochem Biophys* 2009; **482**: 104-111
- 155 **Grundy D**. Signalling the state of the digestive tract. *Auton Neurosci* 2006; **125**: 76-80
- 156 **Jones MP**, Dilley JB, Drossman D, Crowell MD. Brain-gut connections in functional GI disorders: anatomic and physiologic relationships. *Neurogastroenterol Motil* 2006; **18**: 91-103
- 157 **Newberry RD**. Intestinal lymphoid tissues: is variety an asset or a liability? *Curr Opin Gastroenterol* 2008; **24**: 121-128
- 158 **Johansson-Lindbom B**, Agace WW. Generation of gut-homing T cells and their localization to the small intestinal mucosa. *Immunol Rev* 2007; **215**: 226-242
- 159 **Sand J**, Lankisch PG, Nordback I. Alcohol consumption in patients with acute or chronic pancreatitis. *Pancreatology* 2007; **7**: 147-156

S- Editor Li LF L- Editor Alpini GD E- Editor Yin DH