

University of Massachusetts Medical School

eScholarship@UMMS

UMass Center for Clinical and Translational
Science Research Retreat

2012 UMass Center for Clinical and
Translational Science Research Retreat

May 22nd, 4:30 PM - 6:00 PM

Gout Knowledge, Beliefs and Treatment Practices in a National Sample of US Primary Care Providers

Leslie R. Harrold

University of Massachusetts Medical School

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat



Part of the [Health Services Research Commons](#), [Primary Care Commons](#), and the [Rheumatology Commons](#)

Harrold LR, Mazor KM, Negron A, Ogarek J, Firreno CL, Yood RA. (2012). Gout Knowledge, Beliefs and Treatment Practices in a National Sample of US Primary Care Providers. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts_retreat/2012/posters/31

Creative Commons License



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#).

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

GOUT KNOWLEDGE, BELIEFS AND TREATMENT PRACTICES IN A NATIONAL SAMPLE OF US PRIMARY CARE PROVIDERS

Leslie R. Harrold¹, Kathy M. Mazor², Amarie Negron¹, Jessica Ogarek², Cassie Firreno², Robert A. Yood³

¹Medicine, University of Massachusetts Medical School, ²Meyers Primary Care Institute, ³Reliant Medical Group, Worcester, MA

Contact Information:

Leslie R. Harrold, MD, MPH

Phone: (508) 856-3085

Fax: (508) 856-5024

Email: Leslie.Harrold@umassmed.edu

Objectives: We sought to examine primary care providers' gout knowledge and reported treatment patterns in comparison to current treatment recommendations.

Methods: We conducted a national survey of a random sample of primary care physicians (internists and family practitioners) to assess their treatment of acute, intercritical, and tophaceous gout in comparison to published European and American gout treatment recommendations and guidelines.

Results: There were 838 respondents (response rate of 41%) most of whom worked in private practice (63%) with >16 years experience (52%). For acute podagra, 16% reported care consistent with the recommendations including avoidance of anti-inflammatory drugs in the setting of renal insufficiency, use of colchicine at a dose of ≤ 1.8 mg a day, and no initiation of a urate-lowering drug (ULD) during an acute attack (internists 21% vs. family practitioners 11%, $p < 0.001$). For intercritical gout, 3% would provide care consistent with the recommendations including initiating a ULD at the appropriate dose in the setting of renal insufficiency, with dosing titration to a serum urate level of ≤ 6 mg/dl and providing prophylaxis with colchicine (internists 5% vs. family practitioners 3%, $p = 0.12$). For tophaceous gout, 17% reported care consistent with the recommendations including ULD use with dosing titration to a serum urate level of ≤ 6 mg/dl and prophylaxis with nonsteroidal anti-inflammatories or colchicine (internists 22% vs. 13% family practitioners, $p < 0.001$).

Conclusion: While internists were more likely than family practitioners to suggest recommended care, the majority of physicians did not report optimal treatment practices, suggesting further education is needed.