

University of Massachusetts Medical School

eScholarship@UMMS

UMass Center for Clinical and Translational
Science Research Retreat

2012 UMass Center for Clinical and
Translational Science Research Retreat

May 22nd, 4:30 PM - 6:00 PM

Induction of Colon Cancer Cell Death by Cranberry Proanthocyanidins via MAPK Pathway

Anne M. Liberty

University of Massachusetts - Dartmouth

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat



Part of the Biochemistry, Biophysics, and Structural Biology Commons, Cell and Developmental Biology Commons, Medical Biochemistry Commons, and the Medical Nutrition Commons

Liberty AM, Ferreira TP, Neto CC. (2012). Induction of Colon Cancer Cell Death by Cranberry Proanthocyanidins via MAPK Pathway. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts_retreat/2012/posters/41

Creative Commons License



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-sa/3.0/).

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

INDUCTION OF COLON CANCER CELL DEATH BY CRANBERRY PROANTHOCYANIDINS VIA MAPK PATHWAY

Anne M. Liberty¹, Tracie P. Ferreira² and Catherine C. Neto¹

UMass Cranberry Health Research Center and ¹Department of Chemistry and Biochemistry;
²Department of Bioengineering, University of Massachusetts-Dartmouth, North Dartmouth, MA

Contact persons: Catherine Neto (cneto@umassd.edu), Anne Liberty (aliberty@umassd.edu)

The potential ability of dietary cranberry to inhibit colon carcinogenesis is under investigation. Compounds isolated from locally grown cranberry fruit (*Vaccinium macrocarpon*) have been shown *in vitro* to decrease proliferation of colon cancer cells, in part by induction of apoptosis. These compounds include oligomeric polyphenols known as proanthocyanidins (PACs) containing two or more epicatechin units with different types of linkages. To further elucidate the mechanism by which PACs induce cell death, we transcriptionally profiled cells treated with PACs. HCT116 and HT29 colon cancer cells were exposed to a cranberry proanthocyanidin (PACs) fraction isolated from Early Black variety cranberry fruit, at time intervals of 6, 12, 18 and 24 hours. Total RNA was extracted from PAC-treated and untreated control cells. Transcriptional profiling was performed using an Illumina microarray bead system. Microarray results revealed that expression of several members of the mitogen activated protein kinase family (MAPK) was significantly altered in the presence of PACs, leading to decreased transcription of genes in the nucleus and decreased tumor cell growth. Quantitative (Q)-PCR was used to confirm microarray data showing gene expression changes in some key apoptotic pathways. Western blotting was used to confirm the up- regulation or down-regulation of key proteins involved in the MAPK pathway. Significant changes in p53, APAF and VEGF protein expression were seen as early as eighteen hours. Flow cytometry was employed to identify changes in the cell cycle due to exposure to PACs. HCT116 and HT29 colon cancer cells showed a significant change in granularity and a significant increase in G₂ arrest compared to control when exposed to PACs for as little as six hours. This study has provided insight into mechanisms by which cranberry PACs may inhibit colon cancer.