

University of Massachusetts Medical School

eScholarship@UMMS

---

UMass Center for Clinical and Translational  
Science Research Retreat

2012 UMass Center for Clinical and  
Translational Science Research Retreat

---

May 22nd, 4:30 PM - 6:00 PM

## Factors Associated With Ordering Laboratory Monitoring Of High-Risk Medications

Shira H. Fischer

*University of Massachusetts Medical School*

*Et al.*

Let us know how access to this document benefits you.

Follow this and additional works at: [https://escholarship.umassmed.edu/cts\\_retreat](https://escholarship.umassmed.edu/cts_retreat)



Part of the [Health and Medical Administration Commons](#), and the [Health Services Research Commons](#)

---

Fischer SH, Field TS, Peterson DJ, Reed G, Gurwitz JH, Tjia J. (2012). Factors Associated With Ordering Laboratory Monitoring Of High-Risk Medications. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from [https://escholarship.umassmed.edu/cts\\_retreat/2012/posters/25](https://escholarship.umassmed.edu/cts_retreat/2012/posters/25)

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](#).

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

## **FACTORS ASSOCIATED WITH ORDERING LABORATORY MONITORING OF HIGH-RISK MEDICATIONS**

Shira Fischer<sup>1</sup>, AB, Candidate for MD/PhD; Terry S. Field<sup>1,2</sup>, DSc; Daniel Peterson<sup>2</sup>, MS; George Reed<sup>1</sup>, PhD; Jerry H. Gurwitz<sup>1,2</sup>, MD; Jennifer Tjia<sup>1,2</sup>, MD, MSCE

<sup>1</sup>University of Massachusetts Medical School, <sup>2</sup>Meyers Primary Care Institute – a joint endeavor of the University of Massachusetts Medical School, Reliant Medical Group, and Fallon Community Health Plan

Corresponding Author: Shira Fischer, AB, Candidate for MD/PhD, University of Massachusetts Medical School, Biotech Four, 377 Plantation Street, Suite 315, Worcester, MA 01605; Phone, 617-777-4844; Email, shira.fischer@umassmed.edu

### **Abstract**

#### **Background**

Knowledge about factors associated with provider ordering of appropriate testing is limited.

#### **Objective**

To determine physician factors correlated with ordering of recommended laboratory monitoring tests for high-risk medications, accounting for patient characteristics.

#### **Methods**

Analysis of the administrative claims and electronic medical records of patients prescribed a high-risk medication requiring laboratory monitoring in a large multispecialty group practice between January 1, 2008 and December 31, 2008. The outcome is a physician order for each recommended laboratory test for each prescribed medication. Key predictor variables include physician characteristics, including age, gender, specialty training, years since completing training, and prescribing volume. We used multivariable logistic regression to identify the independent association of physician and patient characteristics with ordering of laboratory tests to monitor medications after adjustment for potential confounders, taking into account clustering of drugs within patients and patients within providers.

#### **Results**

Physician orders for laboratory testing varied across drug-test pairs and ranged from 9% (Primidone–Phenobarbital level) to 97% (Azathioprine–CBC) with 50% of drug-test pairs in the 85-91% ordered range. Failure to order a test was associated with lower provider prescribing volume for study drugs and whether the physician was a specialist (primary care providers were more likely to order tests than specialists). Patients with lower patient comorbidity burden and younger patients were less likely to

have appropriate tests ordered. Drug-test combinations with black box warnings were more likely to have tests ordered.

### **Conclusions**

Interventions targeting providers should be addressed at those subgroups with the greatest potential for improvement: providers with lower frequencies of prescribing high-risk medications, and healthier and younger patients. Drug-test combinations with black box warnings have higher ordering rates, but many medications without such warnings also have evidence of harm, thus efforts to improve testing are necessary for all medications shown to be high-risk.