

University of Massachusetts Medical School

eScholarship@UMMS

Open Access Articles

Open Access Publications by UMMS Authors

2005-04-30

Double dissociated tachycardias: isorhythmic A-V dissociation at rapid atrial and ventricular rates

David H. Spodick

University of Massachusetts Medical School

Let us know how access to this document benefits you.

Follow this and additional works at: <https://escholarship.umassmed.edu/oapubs>

 Part of the [Cardiology Commons](#)

Repository Citation

Spodick DH. (2005). Double dissociated tachycardias: isorhythmic A-V dissociation at rapid atrial and ventricular rates. Open Access Articles. Retrieved from <https://escholarship.umassmed.edu/oapubs/18>

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Open Access Articles by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Double Dissociated Tachycardias: Isorhythmic A-V Dissociation at Rapid Atrial and Ventricular Rates

David H. Spodick, MD, DSc

[click an author to search for more articles by that author](#)

A 44-year-old man with coronary artery disease was evaluated. The atrium (P waves) shows sinus rhythm (P axis approximately $+80^\circ$). The ventricles are controlled by a junctional pacemaker until after beat number 13. Sinus rhythm then takes over with conduction through the junction and no change in the heart rate. The paroxysmal junctional tachycardia is at 118 bpm and the atrial (sinus) rate is approximately the same, though probably minimally slower during the dissociation (which would permit dissociation). During dissociated beats, the P-Q intervals are short (under 12 milliseconds) or absent (P simultaneous with QRS). Also typical of A-V dissociation, the P waves never get into early or mid diastole (if it were third-degree A-V block, P waves would be anywhere). Somewhat atypical is the occurrence of paroxysmal junctional tachycardia in an adult, as its occurrence is most frequent among children. As with many A-V dissociations, this may be a toxic rhythm. The T-wave configurations in V_2 and V_3 are unusual and T V_4 and T V_5 are bifid (T-U waves). The patient also has minimal QRS voltage for left ventricular hypertrophy. The vertical P wave axis suggests emphysema, unless this is actually an ectopic atrial tachycardia, which occasionally occurs with a P axis within the accepted range for sinus rhythm.

