

University of Massachusetts Medical School  
**eScholarship@UMMS**

---

UMass Center for Clinical and Translational  
Science Research Retreat

2011 UMass Center for Clinical and  
Translational Science Research Retreat

---

May 20th, 5:00 PM - 7:00 PM


## Determining the Health Utility of Urinary Incontinence in Women

Danielle Patterson  
*University of Massachusetts Medical School*

*Et al.*

Let us know how access to this document benefits you.

Follow this and additional works at: [https://escholarship.umassmed.edu/cts\\_retreat](https://escholarship.umassmed.edu/cts_retreat)

 Part of the [Clinical Epidemiology Commons](#), [Health Services Research Commons](#), [Obstetrics and Gynecology Commons](#), and the [Urology Commons](#)

---

Patterson D, Geisler B, Morse A. (2011). Determining the Health Utility of Urinary Incontinence in Women. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from [https://escholarship.umassmed.edu/cts\\_retreat/2011/posters/10](https://escholarship.umassmed.edu/cts_retreat/2011/posters/10)

Creative Commons License




This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/). This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

# Determining the Health Utility of Urinary Incontinence in Women

Danielle Patterson MD<sup>1</sup>; Benjamin Geisler MD, MPH<sup>2</sup> and Abraham Morse MD, MBA<sup>2</sup>

UMass Memorial Medical Center/University of Massachusetts Medical School, Department of Obstetrics & Gynecology, Worcester<sup>1</sup>  
Brigham and Women's Hospital/Harvard Medical School, Department of Obstetrics & Gynecology, Boston, Massachusetts<sup>2</sup>



## Objective

The goal of this study was to define the utility of urinary incontinence in women using the Standard Gamble, the gold standard method for determining health state utilities, based on a diagnosis obtained from multichannel urodynamic testing, the gold standard in clinical diagnosis.

## Background

Health state utility values are important in many areas of medical research. The values are used in cost-utility analysis, decision analysis and health related quality of life studies. To date, studies that have estimated the utility of urinary incontinence in women have relied on values from generic health related quality of life questionnaires such as the ED-5Q and Health Utilities Index or from expert opinion. [1-5] The utility of urinary incontinence in these studies appears to be unintuitively low, at 0.71 to 0.82, with perfect health represented by 1.0. [6, 7] The utility of health states that are much more debilitating, for example cancer (0.82), is higher than urinary incontinence. [6]

These studies have relied on patient self-diagnosis of incontinence. Additionally, they have considered all types of urinary incontinence together. Intuitively, one would think that quality of life would be affected differently with different types (stress, urge, mixed) and differing severity of incontinence.

## Methods

- All adult female patients who underwent urodynamic testing at Brigham and Women's Hospital were prospectively recruited
- Diagnosis of type of incontinence was made by attending physician interpretation of the urodynamic study
- No exclusion criteria
- Patients completed three validated questionnaires
  - Sandvik Severity Index**  
A validated two question symptom specific instrument to evaluate urinary incontinence
  - EQ-5D**
    - A five-domain generic quality of life questionnaire
    - Answers are converted into a utility value
  - Visual Analog Scale**
    - Vertical line from 0 (worst imaginable health) to 100 (best imaginable health)
    - Patient rates own perception of health on line
- The Standard Gamble technique was used in a standard format to determine each patient's utility value for their health state
  - Patient is asked to choose between life in current health state and varying risks of immediate painless death
  - Gold Standard method to determine patients utility preference for their health state

## References

- Manca A, Sculpher MJ, Ward K, Hilton P. A cost-utility analysis of tension-free vaginal tape versus colposuspension for primary urodynamic stress incontinence. *BJOG* 2003;110:255-62.
- Wu JM, Visco AG, Weidner AC, Myers ER. Is burch colposuspension ever cost-effective compared with tension-free vaginal tape for stress incontinence? *AJOG* 2007;197:62e1-5.
- Subak LL, Brown JS, Kraus SR, Brubaker L, et al. The "costs" of urinary incontinence for women. *Obstet Gynecol* 2006;107:908-16.
- Subak LL, Brubaker L, Chai TC, Creasman JM, et al. High costs of urinary incontinence among women electing surgery to treat stress incontinence. *Obstet Gynecol* 2008;111:899-907.
- Culligan PJ, Myers JA, Goldberg RP, Blackwell L, et al. Elective cesarean section to prevent anal incontinence and brachial plexus injuries associated with macrosomia—a decision analysis. *Int Urogynecol J* 2005;16:19-28.
- Mittman N, Trakas K, Risebrough N, Liu BA. Utility scores for chronic conditions in a community-dwelling population. *Pharmacoeconomics* 1999;15:369-76.
- Schultz SE, Kopec JA. Impact of chronic conditions. *Health Reports* 2003;14:41-53.
- Sandvik H, Seim A, Vanvik A, Hunskaar S. A severity index for epidemiological surveys of female urinary incontinence: comparison with 48-hour pad-weighing tests. *Neurourol Urodyn* 2000;19:137-45.
- Johnson JA, Coons SJ, Ergo A, Szava-Kovats G. Valuation of EuroQOL (EQ-5Q) Health States in an Adult US Sample. *Pharmacoeconomics* 1998;13:421-3.


## Results

- This pilot study of 28 patients demonstrated a significant difference in utility value derived from the Standard Gamble and the generic health-related quality of life instruments
- There were 21 patients with stress urinary incontinence, 6 patients with urge urinary incontinence and 1 patient with mixed urinary incontinence
- Mean Sandvik score was higher in Urge Incontinence subgroup
- Mean Utility from Standard Gamble was lower in Urge Incontinence group
- Sandvik scores were moderately correlated with EQ-5D, SG and VAS utilities

Table 1. Demographic Characteristics & Sandvik Score


	Combined Group	Stress Incontinence	Urge Incontinence
Age (y)	55.5 ± 15.8	58.9 ± 12.9	42.0 ± 25.5
BMI (kg/m <sup>2</sup> )	29.3 ± 7.9	29.3 ± 8.4	29.2 ± 8.3
Menopause	70 %	50%	75%
Sandvik	8 ± 3	7 ± 3	12 ± 0

Fig 1. Urinary Incontinence Utilities



\*Wilcoxon Rank Sum test

Fig 2. Stress Incontinence Utilities



\*Wilcoxon Rank Sum test

## Conclusions

- Utility scores derived from Standard Gamble were significantly higher than those derived from generic health related quality of life instruments.
- Utility scores derived from EQ-5D and VAS were similar to those previously reported in the literature. [6,7]
- Current utility values over-estimate the degree of both of urinary incontinence.
- Researchers should consider using higher utility values for urinary incontinence in future cost utility and quality of life studies.