

University of Massachusetts Medical School

eScholarship@UMMS

UMass Center for Clinical and Translational
Science Research Retreat

2013 UMass Center for Clinical and
Translational Science Research Retreat

May 8th, 12:30 PM - 1:30 PM

Analysis of sFlt Isoforms as Biomarkers for the Development of Preeclampsia

Colby A. Souders

University of Massachusetts Medical School

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat



Part of the [Biological Factors Commons](#), [Maternal and Child Health Commons](#), [Obstetrics and Gynecology Commons](#), [Translational Medical Research Commons](#), and the [Women's Health Commons](#)

Souders CA, Boatright NK, Sedan J, Molrine D, Thomas WD. (2013). Analysis of sFlt Isoforms as Biomarkers for the Development of Preeclampsia. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts_retreat/2013/posters/8

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](#).

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Analysis of sFlt Isoforms as Biomarkers for the Development of Preeclampsia

Colby A Souders¹, Naomi K Boatright¹, Jessica Sedan¹, Deborah Molrine² and William Thomas¹

MassBiologics of the University of Massachusetts Medical School, Departments of Product Discovery¹ and Clinical Affairs², 460 Walk Hill St, Boston, MA 02126.

Corresponding Author: Colby.Souders@umassmed.edu, (617)-474-4050

Abstract:

Preeclampsia is a multi-system disorder characterized by hypertension, edema and proteinuria affecting between 5-10% of pregnancies. A subset of cases progress to severe preeclampsia with exacerbated hypertension/proteinuria and evidence of nervous system, liver and/or kidney dysfunction, in addition to fetal growth restriction. Soluble fms-like tyrosine kinase-1 (sFlt) is minimally expressed in many tissues, including the placenta, and is a circulating antagonist to vascular endothelial growth factor. With progression of pregnancy, sFlt levels significantly rise, especially in women who develop preeclampsia. Diagnostic tests to predict preeclampsia in pregnant women are limited and current tests measure total sFlt in relationship to placental growth factor with varying sensitivity and specificity. We hypothesized that a pregnancy-specific splice variant of sFlt (sFlt1-14), almost exclusively expressed by the placenta, would serve as an improved serum biomarker for the development of preeclampsia. Monoclonal antibodies (mAbs) were developed that specifically bind the two predominant isoforms of sFlt (sFlt1 and sFlt1-14) by hybridoma generation from wild type mice immunized with c-terminal peptides of the two isoforms. Western blot, ELISA and affinity analysis indicated the mAbs were specific for sFlt1 or sFlt1-14 splice variants and recognized these proteins in biological fluids (amniotic fluid or serum). A quantitative capture ELISA was developed whereby total sFlt in biological fluid is captured by a unique human mAb and specific levels of sFlt1 or sFlt1-14 are detected by their respective mouse mAb, followed by anti-murine secondary antibody development. Using recombinant sFlt1 or sFlt1-14 as standards, these endogenous proteins were quantified in commercially available third trimester human pregnant sera. Future studies will measure these isoforms in sera prospectively collected from women with known outcomes of a healthy pregnancy or preeclampsia and the ability of absolute quantitation of the isoform(s) or a ratio of the two to predict the likely onset and severity of preeclampsia will be evaluated.