

University of Massachusetts Medical School

eScholarship@UMMS

---

UMass Center for Clinical and Translational  
Science Research Retreat

2013 UMass Center for Clinical and  
Translational Science Research Retreat

---

May 8th, 12:30 PM - 1:30 PM


## Obesity-Induced Diabetes and Lower Urinary Tract Fibrosis Promote Urinary Voiding Dysfunction in a Mouse Model

Mehrnaz Gharaee-Kermani  
*University of Massachusetts Boston*

*Et al.*

Let us know how access to this document benefits you.

Follow this and additional works at: [https://escholarship.umassmed.edu/cts\\_retreat](https://escholarship.umassmed.edu/cts_retreat)

 Part of the [Cell Biology Commons](#), [Endocrine System Diseases Commons](#), [Male Urogenital Diseases Commons](#), and the [Translational Medical Research Commons](#)

---

Gharaee-Kermani M, Rodriguez-Nieves JA, Macoska JA. (2013). Obesity-Induced Diabetes and Lower Urinary Tract Fibrosis Promote Urinary Voiding Dysfunction in a Mouse Model. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from [https://escholarship.umassmed.edu/cts\\_retreat/2013/posters/52](https://escholarship.umassmed.edu/cts_retreat/2013/posters/52)

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](#). This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

# **Obesity-Induced Diabetes and Lower Urinary Tract Fibrosis Promote Urinary Voiding Dysfunction in a Mouse Model**

Mehrnaz Gharaee-Kermani, Jose A. Rodriguez-Nieves and Jill A. Macoska

Center for Personalized Cancer Therapy and Department of Biology, The University of Massachusetts, Boston,

**Background:** Progressive aging- and inflammation-associated fibrosis effectively remodels the extracellular matrix to increase prostate tissue stiffness and reduce urethral flexibility, resulting in urinary flow obstruction and Lower Urinary Tract Symptoms (LUTS). In the current study we sought to test whether senescence-accelerated mouse prone (SAMP)6 mice, which were reported to develop prostatic fibrosis, would also develop LUTS, and whether these symptoms would be exacerbated by diet-induced obesity and concurrent Type 2 Diabetes Mellitus (T2DM).

**Methods:** To accomplish this, SAMP6 and AKR/J background strain mice were fed regular mouse chow, low fat diet chow, or high fat diet chow for 8 months, then subjected to glucose tolerance tests, assessed for plasma insulin levels, evaluated for urinary voiding function, and assessed for lower urinary tract fibrosis.

**Results:** The results of these studies show that SAMP6 mice and AKR/J background strain mice develop diet-induced obesity and T2DM concurrent with urinary voiding dysfunction. Moreover, urinary voiding dysfunction was more severe in SAMP6 than AKR/J mice and was associated with pronounced prostatic and urethral tissue fibrosis.

**Conclusions:** Taken together, these studies suggest that obesity, T2DM, lower urinary tract fibrosis, and urinary voiding dysfunction are inextricably and biologically linked.

Supported by MICHR grant U034697 (MGK) and NIH/NIDDK grant 1P20DK090870-03 (JAM)