

University of Massachusetts Medical School

eScholarship@UMMS

---

UMass Center for Clinical and Translational  
Science Research Retreat

2013 UMass Center for Clinical and  
Translational Science Research Retreat

---

May 8th, 12:30 PM - 1:30 PM

## Efficacy of Intraoperative Recurrent Laryngeal Nerve Monitoring: A Single-Institutions' Experience


Rashelle N. Damle

*University of Massachusetts Medical School*

*Et al.*

Let us know how access to this document benefits you.

Follow this and additional works at: [https://escholarship.umassmed.edu/cts\\_retreat](https://escholarship.umassmed.edu/cts_retreat)

 Part of the [Health Services Research Commons](#), [Surgery Commons](#), [Surgical Procedures, Operative Commons](#), and the [Translational Medical Research Commons](#)

---

Damle RN, Dinh K, Larkin AC, Quinlan RM, Whalen GF. (2013). Efficacy of Intraoperative Recurrent Laryngeal Nerve Monitoring: A Single-Institutions' Experience. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from [https://escholarship.umassmed.edu/cts\\_retreat/2013/posters/66](https://escholarship.umassmed.edu/cts_retreat/2013/posters/66)

Creative Commons License



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#).

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

**Title:** Efficacy of Intraoperative Recurrent Laryngeal Nerve Monitoring: A Single-Institutions' Experience.

**Authors:** Damle, Rachele N<sup>1</sup>, Dinh, Kate<sup>1</sup>, Larkin Anne C<sup>2</sup>, Quinlan, Robert<sup>2</sup>, Whalen, Giles<sup>2</sup>.

**Institutional Affiliations:**

1. Center for Outcomes Research  
Department of Surgery  
UMass University Medical Center  
Worcester, MA

2. Department of Surgical  
Oncology/Endocrine Surgery  
UMass University Medical Center  
Worcester, MA

**Contact Information:**

Rachele Damle  
55 Lake Ave North  
Room S3-752  
Worcester, MA 01655  
[Rachele.Damle@umassmemorial.org](mailto:Rachele.Damle@umassmemorial.org)  
(209) 678-8279

**ABSTRACT:**

**Objective:** To evaluate the efficacy of intraoperative nerve monitoring (IONM) during thyroidectomy in preventing recurrent laryngeal nerve (RLN) injury.

**Design:** Retrospective cohort study.

**Setting:** Academic institution.

**Patients:** Consecutive sample of subjects undergoing thyroidectomy by experienced endocrine surgeons between 2006 and 2008 at a single institution.

**Intervention:** Intraoperative RLN monitoring.

**Main outcome measure:** RLN injury.

**Results:** Between 2006 and 2008, 296 subjects underwent thyroid lobectomy or total thyroidectomy by the authors. One patient was excluded because of preoperative documentation of RLN injury. IONM was used in 253 (88%) cases, with a total of 403 nerves at risk of injury. Loss of RLN signal following surgical dissection occurred in 13 cases, prompting a change in surgical plan in one case. Post-operative laryngoscopy was performed in eight patients with hoarseness, documenting vocal cord paralysis in one patient who had clear intraoperative anatomic evidence of RLN injury. In no case did loss of RLN signal after dissection lead to nerve injury in the absence of anatomical evidence of injury as detected by the surgeon.

**Conclusions:** IONM added cost and resulted in surgeon angst in cases of malfunction without a clear benefit in RLN identification and protection. Anatomic identification of the RLN should remain the gold standard in preventing RLN injury during thyroidectomy.