University of Massachusetts Medical School

eScholarship@UMMS

UMass Center for Clinical and Translational Science Research Retreat

2014 UMass Center for Clinical and Translational Science Research Retreat

May 20th, 12:30 PM

Targeted Destruction of Triple Negative Breast Cancer using Nanoparticles

Adeyinka C. Adejumo University of Massachusetts Lowell

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat

Part of the Cancer Biology Commons, Nanomedicine Commons, Neoplasms Commons, Therapeutics Commons, and the Translational Medical Research Commons

Adejumo AC, Ochin CC, Jadia R, Kanik FE. (2014). Targeted Destruction of Triple Negative Breast Cancer using Nanoparticles. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts_retreat/2014/posters/2

Creative Commons License

This work is licensed under a Creative Commons Attribution-Noncommercial-Share Alike 3.0 License. This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Targeted Destruction of Triple Negative Breast Cancer using Nanoparticles

Adeyinka C. Adejumo¹, Chinedu Charles Ochin¹, Rahul Jadia², Fulya Ekiz Kanik¹, Ikjot Singh Sohal³, Prakash Rai^{1,2,3,4}*

¹ University of Massachusetts Lowell, Biomedical Engineering and Biotechnology Program, 1 University Avenue, 01854, Lowell, MA.

² University of Massachusetts Lowell, Department of Plastics Engineering, 1 University Avenue, 01854, Lowell, MA.

³ University of Massachusetts Lowell, Department of Biological Sciences, 1 University Avenue, 01854, Lowell, MA.

⁴ University of Massachusetts Lowell, Department of Chemical Engineering, 1 University Avenue, 01854, Lowell, MA. Prakash Rai@uml.edu

Photodynamic therapy (PDT) is a combination of light and photosensitizing drug in which a photosensitizer is injected intravenously and accumulates in the tissue. This tissue is then irradiated by light at an appropriate wavelength and the drug leads to cytotoxicity with a cascade of biochemical responses which affects and inactivates the cancer cells in the tumor tissue.¹ In the cells, PDT generally induces mitochondrial damage and apoptosis which destroy the tissue and induce an antitumor activity upon illumination.²

Benzoporphyrin-derivative verteporfin (BPD) and curcumin are two photosensitizer drugs having the capability in use of PDT. The therapeutic potential of BPD and curcumin as photosensitizers is limited by their low aqueous solubility and applicability. In this study, it is aimed to increase the bioavailability of these drugs to induce antitumor activity in the tumor cells. Liposomes and PLGA-based polymeric nanoparticles (PLGA NPs) are used as the drug carrier nanoparticles which have been considered as perfect carriers for cancer therapeutics and they are remarkably used in cancer therapy.^{3,4} BPD and curcumin-loaded liposomes and PLGA NPs were synthesized and characterized in terms of size, zeta potential, drug encapsulation efficiency and morphology. These nanoparticles were targeted towards breast cancer cell line (MDA-MB-231) *in vitro* MTS cytotoxicity assays and imaging.

References

- 1. Manyak MJ, Russo A, Smith PD, Glatstein E. J Clin Oncol. 1988;6:380-91.
- 2. Xue L, Chiu S, Oleinick NL. Expl Cell Res. 2001; 263: 145-155.
- 3. Torchilin VP. Nat Rev Drug Discov 2005;4:145-160.
- 4. Acharya S, Sahoo SK. Adv Drug Deliv Rev. 2011;63: 170-183.