

University of Massachusetts Medical School

eScholarship@UMMS

UMass Center for Clinical and Translational
Science Research Retreat

2014 UMass Center for Clinical and
Translational Science Research Retreat

May 20th, 12:30 PM


Pericyte NF- κ B Activation Enhances Endothelial Cell Proliferation and Proangiogenic Cytokine Secretion

Katherine E. LaBarbera
University of Massachusetts Amherst

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat

 Part of the [Cell Biology Commons](#), [Cellular and Molecular Physiology Commons](#), [Exercise Science Commons](#), and the [Translational Medical Research Commons](#)

LaBarbera KE, Hyldahl RD, Witkowski S. (2014). Pericyte NF- κ B Activation Enhances Endothelial Cell Proliferation and Proangiogenic Cytokine Secretion. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts_retreat/2014/posters/61

Creative Commons License



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#).

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Pericyte NF- κ B activation enhances endothelial cell proliferation and proangiogenic cytokine secretion

Katherine E. LaBarbera¹, Robert D. Hyldahl², Sarah Witkowski¹

¹*Department of Kinesiology, University of Massachusetts Amherst*, ²*Department of Exercise Sciences, Brigham Young University*.

Correspondence: Cardiovascular and Molecular Physiology Laboratory, Department of Kinesiology, Totman Bldg., 30 Eastman Ln., University of Massachusetts, Amherst, MA 01003.
Email: klabarbe@kin.umass.edu

Pericytes are skeletal muscle resident, multipotent stem cells that are localized to capillaries. They respond to damage through activation of nuclear-factor kappa-B (NF- κ B), a transcription factor that regulates many cellular processes including inflammation. Research has shown that pericyte NF- κ B activation positively affects myoblast proliferation. It is unknown how pericyte NF- κ B affects signaling and proliferation of endothelial cells, an important component of muscle tissue microcirculation. **PURPOSE:** To determine the effects of altered pericyte NF- κ B activity on endothelial cell proliferation and identify inflammatory factors involved in this cell-cell signaling. **METHODS:** Human primary pericytes were transfected with vectors designed to increase or decrease NF- κ B activity (or empty vector control). Transfected pericytes were co-cultured with human microvascular endothelial cells (HMVECs) using transwell inserts. HMVEC proliferation was assessed via cell counting at 24 and 48 hr. Secreted cytokines in cell culture supernatants were screened using a Luminex multiplex assay. **RESULTS:** HMVEC proliferation was greater in the increased pericyte NF- κ B activity condition compared to the decreased NF- κ B condition at 24 and 48 hr (1.3 fold, $p=0.002$). At 24 hr, cytokine secretion was greater in the increased NF- κ B condition compared to control and decreased NF- κ B conditions for 14 cytokines, including interleukin-8 (IL-8; 6.4 fold, $p<0.01$), monocyte chemoattractant protein-1 (MCP-1; 4.4 fold, $p<0.01$), and regulated on activation, normal T cell expressed and secreted (RANTES; 10.6 fold, $p<0.01$). **CONCLUSION:** NF- κ B activation in pericytes caused increased HMVEC proliferation, which may have been mediated by proinflammatory and proangiogenic cytokines known to be under the transcriptional regulation of NF- κ B.

Supported by: ACSM Foundation Doctoral Student Research Grant