

University of Massachusetts Medical School

eScholarship@UMMS

UMass Center for Clinical and Translational
Science Research Retreat

2014 UMass Center for Clinical and
Translational Science Research Retreat

May 20th, 12:30 PM

Epstein-Barr Virus (EBV)-lytic Cross-reactive Influenza-A (IAV) Memory CD8 T-cells in EBV Sero-negative Middle-aged Adults

Rabinarayan Mishra

University of Massachusetts Medical School

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat



Part of the Immunology of Infectious Disease Commons, Immunopathology Commons, Infectious Disease Commons, Pathology Commons, Translational Medical Research Commons, Virology Commons, and the Virus Diseases Commons

Mishra R, Watkin LB, Aslan N, Gil A, Ruiz De Luzuriaga K, Selin LK. (2014). Epstein-Barr Virus (EBV)-lytic Cross-reactive Influenza-A (IAV) Memory CD8 T-cells in EBV Sero-negative Middle-aged Adults. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts_retreat/2014/posters/65

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Epstein-Barr virus (EBV)-lytic cross-reactive Influenza-A (IAV) memory CD8 T-cells in EBV sero-negative middle-aged adults

Rabinarayan Mishra¹, Levi Watkin¹, Nuray Aslan¹, Anna Gil¹, Katherine Luzuriaga², Liisa K. Selin¹

¹Department of Pathology, ²Department of Molecular Medicine, University of Massachusetts Medical School, Worcester

EBV is a common human pathogen, which infects ~90% of people and establishes a life-long chronic infection. The clinical outcomes of acute infection can range from asymptomatic to severe immunopathology such as infectious mononucleosis (IM). However, for unknown reasons 5-10% of middle-aged adults (>35 years) remain EBV-seronegative (EBV-SN) when the virus infects the vast majority of people, and is actively shed at high titers during chronic infection. Here we show that EBV-SN (ASN) HLA-A2+ middle-aged adults possess a unique IAV-M1-GIL₅₈₋₆₆ memory CD8 T-cell response that cross-reacts with EBV lytic epitopes that differs from teenage EBV-SN (TSN) (18-19 years) and EBV-seropositive (EBV-SP) adult donors. The five tested HLA-A2+ EBV-SN middle-aged adults had a significantly increased IAV-M1₅₈₋₆₆-GIL tetramer+ CD8 frequency compared to EBV-SP donors. Upon exposure to EBV antigens in vitro both IAV-M1₅₈₋₆₆GIL/EBV-BMLF1₂₈₀₋₂₈₈-GLC and IAV-M1₅₈₋₆₆-GIL/EBV-BRLF1₁₀₉₋₁₁₇-YVL, functionally cross-reactive CD8+ responses could be detected in the peripheral blood of middle-aged EBV-SN donors, while only IAV-M1/EBV-YVL cross-reactive responses were detected in some teenage EBV-SN or EBV-seropositive people. Surprisingly, these IAV-M1-GIL-specific CD8 T-cells in middle-aged EBV-SN adults expanded dramatically to EBV lytic antigens and produced cytokines at high functional avidity. They lysed EBV-infected targets and showed potential (by CD103 expression) to enter mucosal epithelial tissue where infection initiates. Additionally, these cross-reactive cells had an oligo-clonal T-cell receptor repertoire different than EBV-SP donors. Taken together these data suggest that an altered cross-reactive T cell repertoire could mediate protective immunity against viral infection. Our results imply that sero-negative adults might have the ability to resist viral infection via heterologous immunity. (NIH-AI49320).

Contact: Rabinarayan.Mishra@umassmed.edu
Ph-508-856-6314