

University of Massachusetts Medical School

eScholarship@UMMS

UMass Center for Clinical and Translational
Science Research Retreat

2014 UMass Center for Clinical and
Translational Science Research Retreat

May 20th, 12:30 PM

Cranberry Fruit and Leaf Polyphenols Inhibit Staphylococcus Bacterial Biofilms


Catherine C. Neto

University of Massachusetts - Dartmouth

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat

 Part of the [Bacterial Infections and Mycoses Commons](#), [Bacteriology Commons](#), [Biochemistry Commons](#), [Organic Chemicals Commons](#), [Plant Sciences Commons](#), [Public Health Commons](#), and the [Translational Medical Research Commons](#)

Neto CC, MacLean J, Song B, Dovell A, Kwasny S, Opperman T. (2014). Cranberry Fruit and Leaf Polyphenols Inhibit Staphylococcus Bacterial Biofilms. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts_retreat/2014/posters/86

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](#).

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Cranberry fruit and leaf polyphenols inhibit *Staphylococcus* bacterial biofilms

Catherine Neto¹, Jason MacLean¹, Biqin Song¹, Anthony Dovell¹, Steven Kwasny², Timothy Opperman²

¹UMass Cranberry Health Research Center and Department of Chemistry and Biochemistry, UMass-Dartmouth, North Dartmouth, MA

²Microbiotix, Inc., Worcester, MA 01605

Contact: Catherine Neto (cneto@umassd.edu)

Cranberry (*Vaccinium macrocarpon*) is known for urinary tract health benefits associated with reducing the adhesion of *E. coli* bacteria. This property has been linked to cranberry polyphenols known as proanthocyanidins. *Staphylococcus* bacteria are a growing public health concern due to development of resistant strains. Identification of agents that inhibit biofilm formation by these bacteria may provide a new route to reduce infection in clinical settings. Fruit and leaves of North American cranberry (*Vaccinium macrocarpon*) and cranberry juice were fractionated and screened for their ability to prevent biofilm formation by several strains of *S. aureus* and *S. epidermidis* bacteria. MALDI-TOF MS analysis of the most bioactive fractions identified the major constituents as proanthocyanidin oligomers (PACs) with A-type linkages, ranging in size from 2-12 degrees of polymerization. Further characterization by NMR is underway. The polyphenol-rich fractions from cranberry leaf, fruit and juice inhibited biofilm formation by strains of *S. aureus* and *S. epidermidis*, with MBIC as low as 3.1 µg/mL, and without significant bacteriocidal activity. Thus, compounds from cranberry fruit, plant material and juice may be useful in reducing *Staphylococcus* biofilms without promoting resistance.