

University of Massachusetts Medical School

eScholarship@UMMS

---

Implementation Science and Practice Advances  
Research Center Publications

Psychiatry

---

2013

## Patterns of Psychotherapy Attendance in Emerging and Mature Adults

Maryann Davis

*University of Massachusetts Medical School*

*Et al.*

Let us know how access to this document benefits you.

Follow this and additional works at: [https://escholarship.umassmed.edu/psych\\_cmhsr](https://escholarship.umassmed.edu/psych_cmhsr)



Part of the [Health Services Research Commons](#), [Psychiatric and Mental Health Commons](#), [Psychiatry Commons](#), [Psychiatry and Psychology Commons](#), and the [Psychoanalysis and Psychotherapy Commons](#)

---

### Repository Citation

Davis M, Fisher WH, Lidz CW, Gershenson B. (2013). Patterns of Psychotherapy Attendance in Emerging and Mature Adults. Implementation Science and Practice Advances Research Center Publications.

Retrieved from [https://escholarship.umassmed.edu/psych\\_cmhsr/620](https://escholarship.umassmed.edu/psych_cmhsr/620)

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Implementation Science and Practice Advances Research Center Publications by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

# Patterns of Psychotherapy Attendance in Emerging and Mature Adults

Maryann Davis, PhD; William Fisher, PhD; Charles Lidz, PhD; and Bernice Gershenson, MPH

## INTRODUCTION

- Approximately 760,000 emerging adults use outpatient psychotherapy in the U.S. each year (Olfson et al., 2002).
- Emerging adults are 1.6-7.9 times more likely to drop out of mental health treatment than fully mature adults (Edlund et al., 2002; Olfson et al., 2002).
- This study compared temporal patterns of attendance and non attendance between emerging and mature adults.

## METHODS

### SAMPLE

The 443 individuals aged 16-55 who initiated individual outpatient psychotherapy between September 1 and November 31, 2006 at a large community mental health center, Community Healthlink, in Worcester, MA

**Table 1.** Baseline characteristics of emerging adults (ages 16-30) and mature adults (ages 31-55) in a sample of individuals initiating individual psychotherapy at a community mental health center (N=433)

Variable	Emerging Adults (N=205)	Mature Adults (N=228)	Total (N=433)
<b>Participant Characteristics</b>			
Gender (%Male)	60.0%	56.6%	58.2%
<b>Diagnosis**</b>			
Schizophrenia & Bipolar Disorders	21.9%	32.9%	27.3%
Affective & Anxiety Disorders	43.4%	47.4%	45.5%
Other Disorders	31.2%	10.1%	20.1%
<b>Payment method*</b>			
Public: Medicaid, CHAMPUS, Medicare, State agency contract	62.9%	76.8%	70.2%
Private Insurance	22.9%	16.2%	19.4%
<b>Concurrent Services Characteristics</b>			
Medication Consult**	47.8%	65.4%	57.0%
Family/Couples Therapy **	19.5%	7.5%	13.2%
Group Therapy	5.9%	9.2%	7.6%
SA Treatment*	2.4%	7.5%	5.1%

\*\*\*p<.001, \*\*p<.01, \*p<.05

**Table 2.** Variables extracted from de-identified administrative data set

Independent Variables:			
◆ Unique client identifier	◆ Therapist ID	◆ Gender	◆ Primary clinical diagnosis
◆ Health care coverage source	◆ Type of service provided	◆ Age	
Dependent Variables:			
◆ Service dates of all outpatient treatment for 78 weeks following initiation of individual psychotherapy			

### DEPENDENT VARIABLES:

1. The total number of sessions in 18 months
2. The number of sessions per month for each of the 18 months

### INDEPENDENT VARIABLES:


1. **Age Group.** Emerging Adults =16-30 years, Mature Adults =31-55 years
2. **Concurrent treatment.** Yes/No for outpatient individual substance abuse, group mental health, family treatment, medication consult, between 1st and last psychotherapy session
3. **Background variables.**
  - a. Gender
  - b. Health care coverage (Private Insurance, Public Coverage, none)
  - c. Primary diagnoses of record
    - i. Schizophrenia and Bipolar Disorder (Schizophrenia & other psychotic disorders and all Bipolar Disorders)
    - ii. Affective and Anxiety Disorders (Major Depressive Disorder, all other mood disorders, & all Anxiety disorders)
    - iii. Other disorders, (Most commonly (78%) adjustment disorders)

### ANALYSIS:


- Age differences analyzed using analysis of covariance (using repeated measures for monthly session attendance)
- Temporal patterns were analyzed using developmental trajectory analysis (Nagin & Land, 1993)

## RESULTS

Total number of sessions attended was significantly lower in Emerging than Mature Adults 13.15+1.09 vs. 17.73+1.03 (F(1, 432)=9.22, p=.003).



**Figure 1.** Age differences in the number of sessions per month, while covarying diagnostic group, and health care payor, revealed a significant month by age group interaction (F(17,413)=1.71, p<.05).



**Figure 2.** Patterns of psychotherapy attendance in a community MH center. Three treatment persistence patterns; Low (13%), Moderate (19.4%), and High (6.2%). Two patterns of treatment desistance; Rapid (40.1%), and Slow (21.3%).

**Table 3.** Age Group (emerging vs. mature adults) by Therapy Attendance Pattern group differences in baseline characteristics and treatment variables in a sample of individuals initiating individual psychotherapy at a community mental health center (N=433)

Variables	Rapid Desisters		Slower Desisters		Persisters	
	Mature Adults (N=86)	Emerging Adults (N=87)	Mature Adults (N=37)	Emerging Adults (N=56)	Mature Adults (N=105)	Emerging Adults (N=62)
% Within Age Group ++	37.7%	42.4%	16.2%	27.3%	46.1%	30.2%
Total Sessions	3.94 (1.19)	3.28(1.16)	12.65(1.79)	14.43(1.47)	30.33(1.06)	26.64(1.38)
Est. Marg. Means (SD)*						
Primary Diagnosis***††						
Schizophrenia or Bipolar	30.6%	17.1%	46.9%	14.5%	37.3%	35.6%
Affective or Anxiety	51.4%	51.2%	53.1%	34.5%	52.4%	47.5%
Other	18.1%	31.7%	0.0%	50.9%	11.2%	16.9%
Health Care Coverage***†						
Public	72.3%	76.6%	100.0%	58.1%	85.4%	80.4%
Private	27.7%	23.4%	0.0%	41.9%	17.5%	19.6%
% Female	43.0%	35.6%	45.9%	39.3%	42.9%	46.8%
Concurrent Treatment						
Group**†	.0%	1.1%	13.5%	8.9%	15.2%	9.7%
Substance Abuse+	12.8%	4.6%	2.7%	1.8%	4.8%	0.00%
Family ** †††	2.3%	6.9%	2.7%	30.4%	13.3%	27.4%
Medication Consult***†††	30.2%	24.3%	83.8%	64.5%	87.6%	86.8%
In Emerging Adults Only						
Est. Marg. Means (SD)						
Age in years	21.85 (0.48)	20.224(0.61)	21.37(0.57)			
Proportion Minors <sup>b</sup>	.223(.048)	.410(.060)	.284(.057)			

Trajectory group differences; Age group Difference ++p<.01  
Trajectory group difference (main effect) ap<.001  
within Mature Adults ap=.053, \*p<.01, \*\*p<.001  
within Emerging Adults bp=.055, †p<.05, ††p<.01, †††p<.001



## DISCUSSION

- Emerging Adults attend fewer psychotherapy sessions than mature adults, independent of health care coverage type or diagnostic category.
- There are different patterns of treatment desistance, which imply different treatment retention efforts earlier and later in treatment.
- Emerging adults are more likely than mature adults to display desisting patterns of psychotherapy attendance.
- There are likely important interactions of legal rights, parental influence, and changing life circumstances that distinguish emerging adults who are legal adults from those that are minors.

## CONCLUSIONS

1. Emerging adults initiating psychotherapy would likely benefit from targeted efforts to retain them in psychotherapy.
2. Those targeted efforts would benefit from;
  - a. Research that identifies factors that could be the focus of interventions, that are common and different between emerging and mature adults to help guide new and adapted treatment emerging adult retention interventions
  - b. Research that identifies malleable factors within emerging adults that distinguish rapid and slower desisters to help guide treatment retention efforts early in therapy from those later in therapy
3. Identifying the factors that distinguish between desisting emerging adults who are legal minors from legal adults may help identify unique treatment retention issues faced by young adults on the threshold of adulthood that treatment retention interventions for adolescents or generic adults may not address.

## REFERENCES CITED

- Barrett, M. S., Chua, W. J., Crits-Christoph, P., Gibbons, M. B., & Thompson, D. (2008). Early withdrawal from mental health treatment: Implications for psychotherapy practice. *Psychotherapy: Theory, Research, Practice, Training, 45*(2), 247.
- Edlund, M.J., Wang, P.S., Berglund, P.A., Kats, S.J., Lin, E., & Kessler, R.C. (2002). Dropping out of mental health treatment: Patterns and predictors among epidemiological survey respondents in the United States and Ontario. *American Journal of Psychiatry 159*: 845-851.
- Olfson, M., Marcus, S., Druss, B., & Pinkus, H. (2002). National trends in the use of outpatient psychotherapy. *American Journal of Psychiatry, 159*, 1914-1920.

## ACKNOWLEDGEMENTS:

This research was funded by a grant from the National Institute of Mental Health (RC1MH088542). Grateful thanks to David Haddad for assistance accessing and understanding the administrative database, and leadership at Community HealthLink for willingness to share data.