

University of Massachusetts Medical School
eScholarship@UMMS

Open Access Articles

Open Access Publications by UMMS Authors

2015-08-01

Influenza virus infections in the tropics during the first year of life

Daniel H. Libraty
University of Massachusetts Medical School

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: <https://escholarship.umassmed.edu/oapubs>

 Part of the [Influenza Humans Commons](#), [International Public Health Commons](#), [Maternal and Child Health Commons](#), [Pediatrics Commons](#), and the [Virus Diseases Commons](#)

Repository Citation

Libraty DH, Zhang L, Caponpon M, Capeding RZ. (2015). Influenza virus infections in the tropics during the first year of life. Open Access Articles. <https://doi.org/10.1093/tropej/fmv020>. Retrieved from <https://escholarship.umassmed.edu/oapubs/2627>

Creative Commons License



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 License](https://creativecommons.org/licenses/by-nc/4.0/)

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Open Access Articles by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

RESEARCH LETTER

Influenza virus infections in the tropics during the first year of life

by Daniel H. Libraty,¹ Lei Zhang,¹ Mercydina Caponpon,² and Rosario Z. Capeding³

¹Division of Infectious Diseases, Department of Medicine, University of Massachusetts Medical School, Worcester, USA 01655

²San Pablo City Health Office, San Pablo, Laguna, Philippines 1770

³Departments of Medicine and Microbiology, Research Institute for Tropical Medicine, Metro Manila, Philippines 1770

Correspondence: Daniel H. Libraty, Rm S6-802b, Division of ID, UMMS, 55 Lake Ave N, Worcester, MA 01655, USA. Tel: 508-856-4182.

Fax: 508-856-4890. E-mail <daniel.libraty@umassmed.edu>

ABSTRACT

Pediatric influenza virus infections in the tropics, particularly during infancy, are not well described. We identified influenza virus infections among infants with non-dengue acute undifferentiated febrile illnesses in San Pablo, Laguna, Philippines, as part of an ongoing clinical study of dengue virus infections during infancy. We found that 31% of infants with non-dengue acute undifferentiated febrile illnesses in San Pablo, Laguna, Philippines, had influenza virus infections. The majority were influenza A virus infections and outpatient cases. The infant ages were 11.1 [9.8–13.0] months (median [95% confidence interval]), and the cases clustered between June and December. Influenza episodes are a common cause of non-dengue acute undifferentiated febrile illnesses in the tropics during the first year of life.

Pediatric influenza virus infections in the tropics, particularly during infancy, are not well described. We identified influenza virus infections among infants with non-dengue acute undifferentiated febrile illnesses in San Pablo, Laguna, Philippines, as part of an ongoing clinical study of dengue virus infections during infancy [1].

Influenza virus infections were searched for among infants with non-dengue acute febrile illnesses. An influenza virus infection in non-dengue febrile study infants was identified by a positive IgM serology by ELISA to influenza A or B virus (Fitzgerald Industries) in either the acute or convalescent serum sample. Sera were first treated with

protein G beads (Millipore Inc.) to remove maternal and infant origin IgG. A positive IgM serology in the protein G-treated serum samples was defined as an ELISA optical density (OD) value \geq 85% of the weak positive control OD.

We tested sera from 121 non-dengue acute undifferentiated febrile illnesses in 118 infants from San Pablo, Laguna, Philippines, for IgM to influenza A and B viruses. We identified 31 influenza A virus infections and 6 influenza B virus infections (31% influenza virus infections). The infant ages were 11.1 [9.8–13.0] months (median [95% confidence interval]), male:female ratio was 23:14 and there were $n = 14$ hospitalized cases and $n = 23$ outpatient cases

Table 1. Characteristics of infants with influenza virus infections

Number of infants	<i>N</i> = 37
Age (months) (median [95% CI])	11.1 [9.8–13.0]
Gender (male:female)	23:14
Influenza virus type	<i>N</i> = 31 Influenza A virus infections <i>N</i> = 6 Influenza B virus infections
Hospitalization status	<i>N</i> = 14 hospitalized <i>N</i> = 23 outpatient




Fig. 1. (A) Age distribution of infants with influenza virus infections, and (B) month of illness distribution of infants with influenza virus infections.

(Table 1). The ages and month of illness distributions for the infant influenza virus infections are shown in Figure 1. The influenza virus infections clustered between June and December. A positive IgM serology to influenza A or B virus was seen between 1 and 79 days after fever onset.

The clinical diagnoses at the time of presentation among the infants with influenza virus infections are shown in Table 2. The sign/symptoms noted at the time of presentation among the infants with influenza virus infections are shown in Table 3. There were $n = 15$ infants with influenza virus infections who had anthropometric measurements performed at the time of the acute febrile illness. Eight of the 15 (53%) infants had WHO length-for-age z scores < -2 .

In a previous study of respiratory illnesses among children < 5 years old in Dhaka, Bangladesh, the incidence of influenza virus infections was 102 of 1000

Table 2. Clinical diagnoses at time of presentation among infants with influenza virus infections

Clinical diagnoses at presentation	<i>N</i>
Upper respiratory infection	22
Lower respiratory infection	5
Gastroenteritis	4
Dengue	4
Hyperreactive airways	3
Oral lesions	2
Lower extremity cellulitis	1
Acute viral illness	1

child-years [2]. We found that 31% of infants with non-dengue acute undifferentiated febrile illnesses in San Pablo, Laguna, Philippines, had influenza virus infections. The majority were influenza A virus

Table 3. Sign/symptoms of infants with influenza virus infections

Sign/symptom	N (%)
Upper respiratory infection	22/37 (60)
Vomiting	13/37 (35)
Cold hands/feet	12/37 (32)
Drowsiness	5/37 (14)
Dyspnea	5/37 (14)
Rash	4/37 (11)
Febrile seizure	2/37 (5)

infections and outpatient cases. The infant ages were 11.1 [9.8–13.0] months (median [95% confidence interval]), and the cases clustered between June and December (rainy season).

In all, 73% of our study infants with influenza virus infections had sign/symptoms of a respiratory infection, and more than three-fourth had sign/symptoms of a respiratory infection or vomiting or diarrhea. Among the 15 infants with anthropometric

measurements at the time of their influenza virus infection, half were stunted. Stunted infants were overrepresented among infants with influenza compared with the general population. In conclusion, influenza episodes are a common cause of non-dengue acute undifferentiated febrile illnesses in the tropics during the first year of life.

FUNDING

National Institutes of Health/National Institute of Allergy and Infectious Diseases [R01 AI091820]. The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

REFERENCES

1. Libraty DH, Acosta LP, Tallo V, *et al.* A prospective nested case-control study of Dengue in infants: rethinking and refining the antibody-dependent enhancement dengue hemorrhagic fever model. *PLoS Med* 2009;6:e1000171.
2. Brooks WA, Goswami D, Rahman M, *et al.* Influenza is a major contributor to childhood pneumonia in a tropical developing country. *Pediatric Infect Dis J* 2010;29:216–21.