University of Massachusetts Medical School

eScholarship@UMMS

Senior Scholars Program

School of Medicine

2011-05-02

Improving Delivery of Evidence-Based Prenatal Care in a Family Medicine Clinic

Rachel Rosenberg University of Massachusetts Medical School

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/ssp

Part of the Community Health and Preventive Medicine Commons, Life Sciences Commons, Maternal and Child Health Commons, Obstetrics and Gynecology Commons, and the Preventive Medicine Commons

Repository Citation

Rosenberg, Rachel and Golding, Jeremy, "Improving Delivery of Evidence-Based Prenatal Care in a Family Medicine Clinic" (2011). University of Massachusetts Medical School. *Senior Scholars Program*. Paper 117.

https://escholarship.umassmed.edu/ssp/117

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Senior Scholars Program by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.



Improving Delivery of Evidence-Based Prenatal Care in a Family Medicine Clinic



Rachel Rosenberg MSIV, Jeremy Golding MD University of Massachusetts Medical School

BACKGROUND:

According to the Institute of Medicine, using evidencebased decision-making is one of the key principles that will enable the health care system to provide consistent, highquality medical care to all people. This can be a challenge when providing care to pregnant women, as ethical issues regarding research in this population have resulted in a relative dearth of high quality randomized control trials providing evidence for prenatal issues. The challenge of providing evidence-based prenatal care is further compounded in a busy Family Medicine teaching practice where pregnant women represent a relatively small fraction of the patients seen on a daily basis.

PURPOSE:

The purpose of this project was to develop concise, evidence-based protocols for the management of several common prenatal problems and implement them in a Family Medicine clinic in order to improve the quality of care provided to pregnant women in this practice.

METHODS:

First, 4 common prenatal problems were identified: 1) Obesity in Pregnancy, 2) Prior Preterm Labor, 3) Gestational Diabetes, 4) Chronic Hypertension in Pregnancy. For each of these problems, a comprehensive review of the literature was performed. Priority was given to guidelines from professional organizations, meta-analyses and randomized control trials. Using the strongest evidence from all of these sources, a one-page protocol was developed for each condition. The protocols then underwent a review process by the physicians at the clinic. In areas where no strong evidence existed, medico-legal considerations and consensus-derived provider preferences were incorporated into the protocols.

Figure 1: Example of evidence-based prenatal protocol Has patient had previous pregnancy in which she went into labor before 37 weeks gestation? Checklist format If no, continue with routine care If yes, she is at greatly increased risk for preterm labor during this pregnancy and facilitates use in these quidelines should be considered. Consensusbusy clinical derived provider practice Assess for additional risk factors for preterm delivery (smoking, periodontal preferences and disease, African descent, cocaine or heroin use, prior 2nd trimester medico-legal issues are If patient is a smoker, counsel her about smoking cessation; if she uses illicit incorporated into substances, refer as needed for treatment protocols Consider screening for asymptomatic BV and treating if positive Obtain consult from high-risk obstetrician: discuss whether and when to transfer care, or whether co-management is suitable. If decision is to transfer care, discuss and document explicit plan for transfer of patient care back to HFHC at 34 weeks if no other high-risk OB conditions are present Administer hydroxyprogesterone 250 mg IM weekly starting by 16 weeks When the evidence gestation and continuing until 36 weeks gestation is inconclusive, If BV screening was performed in the 1st trimester, and test result was provider discretion positive, re-test for BV after treatment has finished should be used OPTIONAL PRACTICES (supporting data is inconclusive): -Transvaginal ultrasound to measure cervical length Can be done once around 24 weeks or serially from 20 weeks Cervical length less than 25 mm is a strong predictor of preterm birth Evidence is strongest for women with multiple previous preterm labors or cervical length less than 15 mm on transvaginal ultrasound

RESULTS:

In each of the 4 problems of interest, there existed at least one recommendation that had strong evidence to support it. These recommendations included screening tools, counseling topics and pharmacologic interventions.

CONCLUSIONS & NEXT STEPS:

The first phase of this project has resulted in the development of concise, evidence-based protocols for care of patients with 4 common prenatal problems that can now be instituted in the Family Medicine clinic. These protocols incorporate the strongest evidence available, and on issues where no strong evidence is available, they take into consideration medico-legal issues and provider preferences derived from a consensus process. We hope that the availability of these protocols will result in more consistent, evidencebased prenatal care. The next steps will be to assess provider utilization of and satisfaction with the protocols, as well as gather outcomes data to see if the implementation of these protocols results in better patient outcomes.

References:

Institute of Medicine, Crossing the Quality Chasm: A New Health
System for the Twenty-first Century (Washington: National
Academy Press, 2001).