### University of Massachusetts Medical School

### eScholarship@UMMS

**UMass Center for Clinical and Translational** Science Research Retreat

2016 UMass Center for Clinical and Translational Science Research Retreat

May 20th, 12:30 PM

# Plasma PLP Concentration and Depressive Symptomatology, over time, in older Latino Adults

Sandra P. Arévalo University of Massachusetts at Lowell

Ft al.

## Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts\_retreat



Part of the Dietetics and Clinical Nutrition Commons, and the Mental Disorders Commons

Arévalo SP, Scott T, Falcón LM, Tucker KL. (2016). Plasma PLP Concentration and Depressive Symptomatology, over time, in older Latino Adults. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts\_retreat/2016/posters/12

Creative Commons License



This work is licensed under a Creative Commons Attribution-Noncommercial-Share Alike 3.0 License. This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For

more information, please contact Lisa.Palmer@umassmed.edu.

Contact: Sandra P. Arévalo, PhD; sandra\_arevalogarcia@uml.edu; 617-799-3225

#### Title of Poster:

Plasma PLP Concentration and Depressive Symptomatology, over time, in older Latino Adults

#### Authors:

Sandra P. Arévalo, PhD;<sup>a</sup> Tammy Scott, PhD, MPH;<sup>b</sup>; Luis M. Falcón, PhD;<sup>c</sup> Katherine L. Tucker, PhD<sup>a</sup> Postdoctoral Research Associate, Department of Clinical Laboratory and Nutritional Sciences, University of Massachusetts at Lowell, 3 Solomont Way, Suite 4, Lowell, MA 01854-3092

- <sup>b</sup> Assistant Professor, Psychiatry, Tufts University School of Medicine, 150 Harrison Avenue, Boston, MA 02111
- <sup>c</sup> Dean, College of Fine Arts, Humanities and Social Sciences, University of Massachusetts at Lowell, 820 Broadway Street, Lowell, MA
- <sup>a</sup> Professor, Department of Clinical Laboratory and Nutritional Sciences, University of Massachusetts at Lowell, 3 Solomont Way, Suite 4, Lowell, MA 01854-3092

### Abstract:

<u>Background</u>: Low vitamin B-6 status has been linked to depressive symptomatology. However, most studies have been cross-sectional and may not have controlled for relevant confounders. Few studies have examined this association in Latino populations at high risk for major depression.

<u>Design</u>: We used two-level hierarchical linear regression models (HLM) for continuous outcomes. Level-1 data included three measures of participant's depressive symptomatology collected at baseline, 2y follow-up and 5y follow-up. Participants constituted level-2 data. Vitamin B-6 was associated with depressive symptomatology across these time points.

<u>Objective</u>: We examined the longitudinal association of vitamin B-6 status with depressive symptomatology across 3 time points over ~ 5-7 y in a cohort of older Puerto Rican adults, a population previously identified to be at high risk for depressive symptomatology and clinical depression.

Results: Plasma pyridoxyl-5'-phosphate (PLP) concentration, a time-varying predictor, was significantly associated with depressive symptomatology. Study participants with PLP deficiency, vs. optimal PLP, had higher baseline depressive symptoms (22±14, vs. 20±13); this differential remained constant over time and persisted after controlling for age, sex, education, BMI, smoking and alcohol use, other relevant nutritional factors, perceived stress, stressful life events and allostatic load; and use of antidepressant medication. However, PLP concentration was not associated with the rate of change in depressive symptomatology over time.

<u>Conclusions</u>: Suboptimal plasma PLP is associated with higher depressive symptomatology in older Puerto Rican adults and this appears to persist over time. Our data suggest that identification and treatment of vitamin B-6 deficiency may be a useful preventive and intervention approach in this population.