University of Massachusetts Medical School

eScholarship@UMMS

UMass Center for Clinical and Translational Science Research Retreat

2016 UMass Center for Clinical and Translational Science Research Retreat

May 20th, 12:30 PM

Geometric Control of YAP-dependent Mechanotransduction: A Proposed Model

Ngozi A. Eze Worcester Polytechnic Institute

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat

Part of the Biochemistry Commons, Biomechanics and Biotransport Commons, Biophysics Commons, and the Cellular and Molecular Physiology Commons

Eze NA, Cirka HA, Billiar KL. (2016). Geometric Control of YAP-dependent Mechanotransduction: A Proposed Model. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts_retreat/2016/posters/19

Creative Commons License

This work is licensed under a Creative Commons Attribution-Noncommercial-Share Alike 3.0 License. This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu. Abstract: Geometric Control of YAP-dependent Mechanotransduction: A Proposed Model Authors: Ngozi A. Eze, Heather A. Cirka, Kristen L. Billiar Department of Biomedical Engineering, Worcester Polytechnic Institute, Worcester, MA

The Billiar lab is interested in the interplay between mechanical tension and programmed cell death (namely, apoptosis) in cells growing on micro-contact printed aggregates. The Billiar lab uses a bioinspired hydrogel to develop an *in vitro* model for mechanosensitive signaling in mammalian cells. The micro-contact printed cell aggregates experience a loss of tensional homeostasis at the center of the aggregates, which results in selective cell death at the center, but not periphery of the aggregates, followed by calcification, similar to excised diseased aortic valves. However, the subcellular mechanisms responsible for transducing the mechanical cues from the loss of tensional homeostasis to pro-apoptotic signaling have yet to be elucidated; the Billiar lab is interested in finding this link.

Mechanotransduction is the functional link between mechanical cues and the consequent subcellular biochemical response.¹ Cells sense and respond to their physical surroundings via cell-cell junctions, cell-matrix adhesions, and intracellular actin networks.¹ For example, in epithelial cells, restriction of cell growth to spatially patterned circular arrays leads to (1) increased proliferation and (2) higher tractional stresses at the periphery than at the center.²⁻³ Proliferation at the periphery of these circular cell aggregates is YAP-dependent, with nuclear localization of YAP at the periphery.⁴ Transcriptional co-activator YAP is (1) a nuclear relay of mechanical signals,⁵ (2) the main transcriptional effector of the Hippo pathway,⁶ and (3) involved in both proliferation (via TEAD promoter) and apoptosis (via p73 promoter).⁷ Cell competition is an apoptosis-dependent cell communication phenomenon based on cell fitness comparisons, and which creates "loser" cells that die via apoptosis and "winner" cells that survive.⁸⁻⁹ For example, co-culture of TEAD-activity-manipulated fibroblasts with WT induces cell competition, in which cells with higher TEAD activity "won," and cells with lower TEAD activity "lost" (underwent apoptosis).¹⁰

<u>Hypothesis</u>: Culture of fibroblasts in geometrically constrained, circular cell aggregates induces cell competition via the formation of "winner" and "loser" cell populations due to differences in tensional homeostasis experienced at the periphery vs. center of the aggregates.

Ngozi A. Eze, Ph.D. Post-Doctoral Fellow & Lab Manager Tissue Mechanics & Mechanobiology Lab Department of Biomedical Engineering Worcester Polytechnic Institute PI: Prof. Kristen L. Billiar, Ph.D. (Department Head) naeze@WPI.EDU

<u>References</u>

1. DuFort, C. C.; M. J. Paszek; V. M. Weaver. Balancing forces: architectural control of mechanotransduction. *Nat Rev Mol Cell Bio*. **2011**, *12* (5), 308-319.

2. Nelson, C. M.; R. P. Jean; J. L. Tan; W. F. Liu; N. J. Sniadecki; A. A. Spector; C. S. Chen. Emergent patterns of growth controlled by multicellular form and mechanics. **2005**, *102* (33), 11594-11599.

3. Shraiman, B. I. Mechanical feedback as a possible regulator of tissue growth. *P Natl Acad Sci USA*. **2005**, *102* (9), 3318-3323.

4. Aragona, M.; T. Panciera; A. Manfrin; S. Giulitti; F. Michielin; N. Elvassore; S. Dupont; S. Piccolo. A Mechanical Checkpoint Controls Multicellular Growth through YAP/TAZ Regulation by Actin-Processing Factors. *Cell.* **2013**, *154* (5), 1047-1059.

5. Halder, G.; S. Dupont; S. Piccolo. Transduction of mechanical and cytoskeletal cues by YAP and TAZ. **2012**, *13* (9), 591-600.

6. Codelia, V. A.; G. P. Sun; K. D. Irvine. Regulation of YAP by Mechanical Strain through Jnk and Hippo Signaling. *Curr Biol.* **2014**, *24* (17), 2012-2017.

7. Bertini, E.; T. Oka; M. Sudol; S. Strano; G. Blandino. YAP: At the crossroad between transformation and tumor suppression. *Cell Cycle*. **2009**, *8* (1), 49-57.

8. Levayer, R.; E. Moreno. Mechanisms of cell competition: Themes and variations. **2013**, 200 (6), 689-698.

9. Amoyel, M.; E. A. Bach. Cell competition: how to eliminate your neighbours. **2014**, *141* (5), 988-1000.

10. Mamada, H.; T. Sato; M. Ota; H. Sasaki. Cell competition in mouse NIH3T3 embryonic fibroblasts is controlled by the activity of Tead family proteins and Myc. **2015**, *128* (4), 790-803.