

University of Massachusetts Medical School

eScholarship@UMMS

UMass Center for Clinical and Translational
Science Research Retreat

2016 UMass Center for Clinical and
Translational Science Research Retreat

May 20th, 12:30 PM

Burden of Adverse Metabolic Factors Is Associated With Increased Left Ventricular Concentricity in Adults With Normal- Range Body Mass Index: The Framingham Heart Study

Philimon Gona
University of Massachusetts Boston

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat



Part of the [Cardiology Commons](#), [Cardiovascular Diseases Commons](#), [Cardiovascular System Commons](#), [Medical Physiology Commons](#), and the [Physiology Commons](#)

Gona P, Lee J, Salton CJ, O'Donnell CJ, Manning WJ, Chuang ML. (2016). Burden of Adverse Metabolic Factors Is Associated With Increased Left Ventricular Concentricity in Adults With Normal-Range Body Mass Index: The Framingham Heart Study. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts_retreat/2016/posters/26

Creative Commons License



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#). This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Burden of Adverse Metabolic Factors Is Associated With Increased Left Ventricular Concentricity in Adults With Normal-Range Body Mass Index: The Framingham Heart Study

*Philimon Gona, PhD MPH¹, Jane Lee, MS PhD², Carol J Salton, BA³, Christopher J O'Donnell, MD MPH^{2,5}, Warren J Manning MD³, Michael L Chuang, MD⁴

¹Department of Exercise and Health Sciences, University of Massachusetts, Boston; ²The NHLBI's Framingham Heart Study;

³Cardiovascular Division and ⁴Department of Radiology, Beth Israel Deaconess Medical Center; ⁵Harvard Medical School

Abstract

Introduction: Persons with normal-range body mass index (BMI) but adverse metabolic characteristics associated with obesity have been described as metabolically-obese normal weight (MONW). We sought to determine whether adverse metabolic profile is associated with alterations in left ventricular (LV) structure or function among adults with normal BMI.

Methods: From the 1794 Framingham Heart Study Offspring cohort adults who underwent cardiac magnetic resonance imaging (CMRI), we identified 446 free of non-skin cancer and prevalent clinical cardiovascular disease (CVD) who had $18.5 \leq \text{BMI} < 25.0$ kg/m² and complete covariates. We calculated a metabolic score (MS) where 1 point was assigned for each of: a) fasting glucose ≥ 100 mg/dL or diabetes; b) SBP ≥ 140 or DBP ≥ 90 mmHg or antihypertensive treatment; c) TG ≥ 150 or HDL_C $< 40(\text{M}) / < 50(\text{W})$ mg/dL or lipid treatment; d) HOMA-IR ≥ 2.5 ; e) waist circumference $\geq 102/88$ cm for M/W. Participants were classified as MS0 (no points), MS1 (exactly 1 point), or MS2+ (≥ 2 points). LV mass (LVM), end-diastolic volume (EDV), ejection fraction (EF), and concentricity (LVM/EDV) were measured from breathhold cine SSFP CMR scans; we calculated LVM/BSA. Analysis of covariance (ANCOVA) was used to compare MS1 and MS2+ groups to the MS0 group. CMRI variables were adjusted for sex, age, heart rate (HR) and body size (BSA); LVM/BSA was adjusted for sex, age, HR only. We also tested for linear trend across metabolic groups.

Results: LV concentricity increased with worsening metabolic status. This was driven by lower LV EDV, not increased LVM. LVM did not differ across (trend) or between MS-groups. LVEDV decreased across groups but only MS2 differed significantly from MS0. LVEF increased slightly but significantly across MS-groups.

Conclusions: In a community-dwelling cohort, among participants who were free of cancer and clinical CVD and had normal BMI, worsening metabolic profile was associated with adverse remodeling of the left ventricle, reflected by greater LV concentricity.

Phil, Gona PhD

Associate Professor of Biostatistics
College of Nursing and Health Sciences
University of Massachusetts
100 Morrissey Blvd
Boston, MA 02125-3393
Phone: 617-287-4826
FAX: 617-287-7527

phil.gona@umb.edu