#### University of Massachusetts Medical School

### eScholarship@UMMS

UMass Center for Clinical and Translational Science Research Retreat

2016 UMass Center for Clinical and Translational Science Research Retreat

May 20th, 12:30 PM

# Theoretical and Experimental Analysis of the Antioxidant and Anti-amyloid Features of Synthetic Resveratrol Mimics

William Horton University of Massachusetts Boston

Et al.

## Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts\_retreat

Part of the Chemical Actions and Uses Commons, Medicinal and Pharmaceutical Chemistry Commons, Medicinal-Pharmaceutical Chemistry Commons, and the Nervous System Diseases Commons

Horton W, Kokel A, Török F, Tran C, Török M, Török B. (2016). Theoretical and Experimental Analysis of the Antioxidant and Anti-amyloid Features of Synthetic Resveratrol Mimics. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts\_retreat/2016/posters/36

Creative Commons License

This work is licensed under a Creative Commons Attribution-Noncommercial-Share Alike 3.0 License. This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu. Theoretical and Experimental Analysis of the Antioxidant and Anti-amyloid Features of Synthetic Resveratrol Mimics

William Horton, Anne Kokel, Fanni Török, Chris Tran, Marianna Török, Béla Török

Department of Chemistry, University of Massachusetts Boston, 100 Morrissey Blvd., Boston, MA 02125

Diaryl hydrazones, possessing similar structure to the popular red wine antioxidant resveratrol, have been previously identified as multitarget compounds interfering with several processes associated with the pathogenesis of Alzheimer's disease (AD). These compounds exhibited particularly strong inhibition of the amyloid beta (A $\beta$ ) peptide self-assembly, including blocking the formation of fibrils and oligomers, species that are widely accepted to be neurotoxic. The molecules were also powerful free radical scavengers and thus have a potential to defend against oxidative stress.

In order to learn more about the mode of action of the compounds, theoretical and experimental studies have been carried out. First, the structural, energetic and electronic features of the core structure have been elucidated by density funtional theory (DFT) calculations. The DFT results identified the most likely form of the compounds, which was applied to a broad range of calculations using substituted derivates. Based on the structural information several characteristics such as logP, H-binding energy, HOMO-LUMO energies and band gap and electron densities were calculated. The compounds were subjected to three different antioxidant assays (DPPH, ABTS and ORAC). The % radical scavenging has been analyzed as a function of the above determined structural parameters in order to identify the role of the energetic and electronic features in the antioxidant activity. The analysis of the same parameters as potential markers to the anti-amyloid activity has also been carried out.

Isotope labeling via H-D exchange and *in situ* hydrazone-radicals as well as hydrazone-A $\beta$  complex formation, applying <sup>1</sup>H-NMR and HRMS, have also been used to experimentally observe the potential role of various tautomeric forms and the partially delocalized electron structure of the hydrazones.

### Contact:

William Horton <u>william.horton001@gmail.com</u> PhD graduate student at the University of Massachusetts Boston in the Chemistry Department focusing on Biological Chemistry.