#### University of Massachusetts Medical School

### eScholarship@UMMS

UMass Center for Clinical and Translational Science Research Retreat

2016 UMass Center for Clinical and Translational Science Research Retreat

May 20th, 12:30 PM

# Sedentary Behavior and Cardiovascular Disease Risk Factors among Latino Adults

Valerie J. Silfee University of Massachusetts Medical School

Et al.

## Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts\_retreat

Part of the Behavioral Disciplines and Activities Commons, Cardiovascular Diseases Commons, Clinical Epidemiology Commons, Community Health and Preventive Medicine Commons, and the Epidemiology Commons

Silfee VJ, Lemon SC, Lora V, Rosal MC. (2016). Sedentary Behavior and Cardiovascular Disease Risk Factors among Latino Adults. UMass Center for Clinical and Translational Science Research Retreat. Retrieved from https://escholarship.umassmed.edu/cts\_retreat/2016/posters/81

Creative Commons License

This work is licensed under a Creative Commons Attribution-Noncommercial-Share Alike 3.0 License. This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu. **Title: Sedentary Behavior and Cardiovascular Disease Risk Factors among Latino Adults** Valerie J. Silfee, PhD<sup>1</sup>, Stephenie C. Lemon, PhD<sup>1</sup>, Vilma Lora<sup>2</sup>, Milagros C. Rosal, PhD<sup>1</sup> <sup>1</sup>University of Massachusetts Medical School Division of Preventive and Behavioral Medicine <sup>2</sup> Director of Social Justice Initiatives,YWCA of Greater Lawrence; Health Task Force Coordinator, City of Lawrence Mayor's Health Task Force

#### Abstract

Background: Compared to other racial/ethnic subgroups in the U.S., Latinos experience increased rates of cardiovascular disease (CVD) and CVD risk factors such as hypertension, inactivity, and diabetes. Sedentary behavior has also been defined as an additional risk factor for CVD, independent of physical activity participation. However, while sedentary behavior has been associated with increased risk for CVD among primarily White samples, previous studies in Latinos have shown mixed results. Purpose: To explore the relationships between sedentary behavior and CVD risk factors, including BMI, waist circumference, blood pressure, physical activity, dyslipidemia, and diabetes, among a sample of Latino adults. Methods: Crosssectional secondary analysis of the Latino Health and Well-Being Study. Latino adults were recruited from the Greater Lawrence Family Health Center (N= 602). Surveys of sedentary behavior and physical activity were verbally administered. Anthropometric measurements included weight, height, waist circumference and blood pressure. Medical record data for diabetes and dyslipidemia were obtained. Results: This study showed that individuals in older age strata, females, and individuals with a less than high school education were more sedentary than their younger, male, and more educated counter parts. Sedentary behavior was positively associated with BMI ( $\beta$  = .164, p < .001) and waist circumference ( $\beta$  = .162, p < .001). There were no associations between sedentary behavior and blood pressure, high cholesterol, diabetes, or physical activity. **Conclusions:** There is growing evidence that sedentary behavior may have its own unique set of metabolic consequences. However, the consequences of sedentary behavior may not be uniform across subgroups. Evaluating the relationship between sedentary behavior and CVD risk is critical in identifying behaviors, like sedentariness, that contribute to the development of CVD.

Corresponding Author: Valerie J. Silfee, PhD 55 Lake Avenue North Worcester, MA 01655-0002 Phone: (508)-856-7561 Email: <u>Valerie.Silfee@umassmed.edu</u>