

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/348351138>

Prevalence of *Staphylococcus aureus* on the Hands of Health-Care Workers and the Environment of a Nigerian University Health Centre

Article · November 2020

DOI: 10.26538/tjnpr/v4i11.7

CITATION

1

READS

46

4 authors, including:



Olayemi Oluseun Akinnola

Covenant University Ota Ogun State, Nigeria

52 PUBLICATIONS 500 CITATIONS

[SEE PROFILE](#)



Bimpe Ogunleye

Covenant University Ota Ogun State, Nigeria

8 PUBLICATIONS 8 CITATIONS

[SEE PROFILE](#)



Ayodeji Samuel Ajayi

Université de Montréal

16 PUBLICATIONS 11 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Epidemiology of Infectious Diseases [View project](#)



Screening of Probiotic Characteristics of Lactic Acid Bacteria Isolated from Some Fermented Nigerian Food Products [View project](#)



Prevalence of *Staphylococcus aureus* on the Hands of Health-Care Workers and the Environment of a Nigerian University Health Centre

Olayemi O. Akinola, Bimpe O. Ogunleye, Rhoda T. Odewunmi, Ayodeji S. Ajayi

Department of Biological Sciences, Covenant University, Ota, Ogun State, Nigeria

ARTICLE INFO

Article history:

Received 28 August 2020

Revised 21 October 2020

Accepted 23 November 2020

Published online 30 November 2020

Copyright: © 2020 Akinola *et al.* This is an open-access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Hospital-acquired infections continue to be an important public health issue with unacceptable levels of morbidity and mortality. *Staphylococcus aureus* is a frequent cause of infections in the hospital. This research aimed at giving a better understanding of the prevalence of *S. aureus* in a hospital setting in order to adequately account for its potential threat. A total of 115 samples comprising 65 environmental samples obtained from trays, bed railings, doorknobs, and other surfaces of patient wards and 50 hand swabs of health-care workers were screened for *S. aureus* using established microbiological techniques. Twenty-eight isolates each represented positive cultures from environmental and hand samples, respectively. Antimicrobial susceptibility testing was determined for the *S. aureus* isolates using the Kirby-Bauer disk diffusion assay. The antibiotic resistance pattern for the hand and environmental isolates respectively gave the following results: Ciprofloxacin (32.14%, 21.43%), Trimethoprim/ Sulphamethoxazole (60.71%, 67.86%), Oxacillin (28.57%, 35.71%), Erythromycin (21.42%, 17.86%), Gentamicin (42.86%, 28.57%), Tetracycline (17.86%, 21.42%), Penicillin (42.86%, 42.85%). Twelve resistant phenotypes were identified, of which Ciprofloxacin-Gentamicin- Trimethoprim/ Sulphamethoxazole (CIP-CN-SXT) was predominant. This study shows that the hands of health-care workers and the hospital environment could serve as potential vehicles of transmission of *S. aureus* infections, thus highlighting the need for effective infection control measures.

Keywords: *Staphylococcus aureus*, Environmental samples, Hand swabs, Antimicrobial susceptibility testing.

Introduction

Nosocomial infections are a common problem in health-care settings. They are associated with increased morbidity and mortality in addition to considerable costs.¹ *Staphylococcus aureus* is an important cause of infections within the hospital, and the community and its global spread and increasing resistance to antimicrobials require numerous actions of prevention and infection control.^{2,3} *S. aureus* is transmissible between patients and occurs mostly via staff and the environment. Colonised health-care personnel have been implicated as sources of transmission in outbreaks.⁴ *S. aureus* is commonly found on surfaces in hospitals such as door handles and floors even after disinfection and can be indirectly transmitted through fomites, thereby serving as vehicles of transmission of infections.⁵ Development of any nosocomial infection, and perhaps particularly those caused by resistant organisms such as methicillin-resistant *Staphylococcus aureus*, may have medicolegal implications as many are transmitted by health-care personnel from one patient to another.⁶ Understanding the trends in the resistance pattern is important in order to have ample information for treatment and infection control.⁷ Therefore, this study was carried out to investigate the prevalence and antimicrobial susceptibility of *S. aureus* obtained from the

environment and hands of health-care workers of a Nigerian University health centre.

Materials and Methods

Collection of samples and identification of isolates

Oral informed consent was obtained from all study participants. The consent procedure as well as the ethical approval for this study was previously approved by the ethics committee of the Covenant University, Ota, Nigeria with number CU/BIOSCREC/BIO/2016/016. The study was carried out in the University health centre between October 2016 and February 2017. A total of 115 samples were investigated (65 from the environment and 50 hand samples). The environmental specimens constituted swabs obtained from trays, bed railings, doorknobs, floors, and furniture in the different patient wards, while the hand swab samples were obtained from health-care workers. The samples were transported to the Microbiology laboratory of the Department of Biological Sciences, Covenant University, Ota, for microbiological analysis. Samples were inoculated on Mannitol Salt Agar, and isolates showing characteristic yellow colour were presumptively identified as *S. aureus* by colony morphology, Gram staining, sugar fermentation, catalase production, and coagulase test. The isolates were then sub-cultured on blood agar to check for the characteristic beta haemolysis displayed by *S. aureus*.

Antimicrobial susceptibility testing

S. aureus isolates were tested for susceptibility to several antibiotics using the Kirby-Bauer disk diffusion method on Mueller-Hinton agar with a panel of 7 antibiotics according to CLSI (Clinical and Laboratory Standards Institute) guidelines, 2016. After incubation for 18 hours at 37°C, the results were recorded as sensitive (S), intermediate (I) and resistant (R). The antibiotics tested include oxacillin (5 µg), gentamicin (30 µg), ciprofloxacin (5 µg), trime-

*Corresponding author. E mail: ola.ayepola@covenantuniversity.edu.ng
Tel: +2348034785269

Citation: Akinola OO, Ogunleye BO, Odewunmi RT, Ajayi AS. Prevalence of *Staphylococcus aureus* on the Hands of Health-Care Workers and the Environment of a Nigerian University Health Centre. Trop J Nat Prod Res. 2020; 4(11):884-886. doi.org/10.26538/tjnpr/v4i11.7

Official Journal of Natural Product Research Group, Faculty of Pharmacy, University of Benin, Benin City, Nigeria.

thoprim/Sulphamethoxazole (25 µg), penicillin (10 µg) and erythromycin (15µg), tetracycline (30 µg).

Results and Discussion

From the 115 samples examined, fifty-six (48.7%) yielded positive *Staphylococcus aureus* cultures. Twenty-eight (50%) of the positive isolates were from environmental samples, and Twenty- eight (50%) were from hand samples. The isolation of *S. aureus* in the health centre reflects its ubiquitous occurrence, as previously reported.⁹ The isolation rate of 48.7% indicates a high occurrence of *S. aureus* when compared to 33.3% obtained in an earlier study from environmental and hand samples in a health-care setting in Africa.¹⁰ The occurrence of the *S. aureus* in the environmental samples (50%) may indicate poor infection control in the hospital environment, thereby making the environment to serve as a reservoir of the pathogen. This may also account for the high occurrence of the pathogen observed on the hands of health personnel, as observed in a previous study carried out in Nigeria.¹¹

The antibiotic resistance pattern as shown in Figure 1 for the hand and environmental isolates respectively is as follows: Ciprofloxacin (32.14%, 21.43%), Trimethoprim/ Sulphamethoxazole (60.71%, 67.86%), Oxacillin (28.57%, 35.71%), Erythromycin (21.42%, 17.86%), Gentamicin (42.86%, 28.57%), Tetracycline (17.86%, 21.42%), Penicillin (42.86%, 42.85%). Resistance to tetracycline and erythromycin was low in this study.

Twelve antimicrobial resistance phenotypes were identified in this study, as presented in Table 1. Of the 12 phenotypes, the most predominant was Ciprofloxacin-Gentamicin-Trimethoprim/ Sulphamethoxazole (CIP-CN-SXT), occurring in 28.6% of the isolates. In contrast, the least common phenotypes were Gentamicin-Trimethoprim/Sulphamethoxazole (CN-SXT) and Gentamicin-Penicillin-Trimethoprim/ Sulphamethoxazole (CN-P-SXT).

The present study showed that MRSA spread in health-care settings is a possibility as oxacillin (methicillin) resistance was detected in 18 (26.79%) out of the 56 *S. aureus* isolates. The high incidence of oxacillin resistance corroborates the findings of another study in Nigeria.¹¹

Approximately 60.71% of the *S. aureus* isolates were resistant to three or more classes of antibiotics. Although a higher percentage of these multidrug-resistant isolates originated from hand samples, 14.2% of the isolates were from the environment. Multidrug resistance in health-care settings has been attributed to the excessive use of antimicrobials in hospitals, which provides a major selective pressure for resistant bacteria.




Figure 1: Antimicrobial resistance pattern of *S. aureus* isolates from the hands and the environment

Table 1: Antimicrobial resistance phenotype of *S. aureus* isolates

Resistance Phenotype	Number of isolates
SXT	3(7.1%)
OX-TE-P-E	4(9.5%)
P-SXT	2(4.8%)
OX-TE-P-E-SXT	5(11.9%)
OX-P-TE	2(4.8%)
OX-P-E-SXT	2(4.8%)
CN	2(4.8%)
CIP-CN-P-SXT	4(9.5%)
CIP-CN-SXT	12(28.6%)
OX-P-SXT	4(9.5%)
CN-SXT	1(2.4%)
CN-P-SXT	1(2.4%)

CIP - Ciprofloxacin, **OX** - Oxacillin, **CN** - Gentamicin, **TE** - Tetracycline, **P** - Penicillin, **E** - Erythromycin, **SXT** - Trimethoprim / Sulphamethoxazole.

Conclusion

The findings of this study show that health-care workers, as well as the hospital environment, could serve as a reservoir of *S. aureus*. Multidrug-resistant strains are found to be common. Regular surveillance of hospital-associated *S. aureus* infections and their susceptibility to antibiotics is of utmost importance in the prevention of the spread of resistant strains in the hospital environment.

Conflict of interest

The authors declare no conflict of interest.

Authors' Declaration

The authors hereby declare that the work presented in this article is original and that any liability for claims relating to the content of this article will be borne by them.

Acknowledgments

The authors acknowledge the support of Covenant University Centre for Research, Innovation, and Discovery (CUCRID).

References

- Breathnach AS. Nosocomial infections and infection control. *Med.* 2013; 41(11):649-653.
- Castro A, Komora N, Ferreira V, Lira A, Mota M, Silva J, Teixeira P. Prevalence of *Staphylococcus aureus* from nares and hands of health care professionals in a Portuguese Hospital. *J Appl Microbiol.* 2016; 121(3):831-839.
- Ayepola OO, Olasupo NA, Egwari LO, Becker K, Schaumburg F. Molecular Characterization and Antimicrobial Susceptibility of *Staphylococcus aureus* Isolates from Clinical Infection and Asymptomatic Carriers in Southwest Nigeria. *PLoS.* 2015; 10(9):0137531.
- Price JR, Cole K, Bexley A, Kostiou V, Eyre DW, Golubchik T, Wilson DJ, Crook DW, Walker AS, Peto T, Llewelyn MJ, Paul J. Modernising Medical Microbiology informatics group. Transmission of *Staphylococcus aureus* between health-care workers, the environment, and patients

- in an intensive care unit: a longitudinal cohort study based on whole-genome sequencing. *Lancet*. 2017; 17(2): 07-214.
5. Omololu JAS. *Staphylococcus aureus* Surface Colonization of Medical Equipment and Environment, Implication in Hospital-Community Epidemiology. *J Hosp Med Manage*. 2017; 3:1.
 6. Ariom TO, Iroha IR, Moses IB, Iroha CS, Ude UI, Kalu AC. "Detection and phenotypic characterization of methicillin-resistant *Staphylococcus aureus* from clinical and community samples in Abakaliki, Ebonyi State, Nigeria." *Afr Health Sci*. 2019; 19(2):2026-2035.
 7. Ayepola OO, Egwari L, Olasehinde GI. Antibiotic resistance profile of *Staphylococcus aureus* clinical isolates from Nigeria. *Antimicrob Resist Infect Control*. 2015; 4(1):195.
 8. Clinical and Laboratory Standards Institute (CLSI). Performance standards for antimicrobial susceptibility testing; twenty-first informational supplement. M100-S21, Clinical and Laboratory Standards Institute, Wayne, PA. 2016; 31:1.
 9. Votintseva AA, Fung R, Miller RR, Knox K, Godwin H, Wyllie DH, Bowden R, Crook DW, Walker AS. Prevalence of *Staphylococcus aureus* protein A (spa) mutants in the community and hospitals in Oxfordshire. *BMC Microbiol*. 2014; 14(1):63.
 10. Abdulgader SM, Shittu AO, Nicol MP, Kaba M. Molecular epidemiology of Methicillin-resistant *Staphylococcus aureus* in Africa: a systematic review. *Front Microbiol*. 2015; 6:348.
 11. Nwokah EG, Eddeh-Adjughah O, Aleru CP. Assessment of asymptomatic methicillin-resistant *Staphylococcus aureus* carriage among health-care workers in the University of Port Harcourt Teaching Hospital, Nigeria. *SCIREA J Health*. 2017; 2(2):13-25.