

# Instruments Measuring Self-Care and Self-Management of Chronic Conditions by Community-Dwelling Older Adults: A Scoping Review

Journal of Applied Gerontology

2023, Vol. 0(0) 1–23

© The Author(s) 2023



Article reuse guidelines:

[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)

DOI: 10.1177/07334648231161929

[journals.sagepub.com/home/jag](https://journals.sagepub.com/home/jag)


Michael T. Lawless<sup>1</sup> , Matthew Tieu<sup>1,2</sup>, Raymond J. Chan<sup>1</sup>, Jeroen M. Hendriks<sup>1</sup>, and Alison Kitson<sup>1</sup>

## Abstract

Given the high prevalence of chronic conditions and multimorbidity in older adults, there is a need to better conceptualize and measure self-care and self-management to promote a person-centered approach. This scoping review aimed to identify and map instruments measuring self-care and self-management of chronic conditions by older adults. We searched six electronic databases, charted data from the studies and tools and reported the results in accordance with the PRISMA-ScR guidelines. A total of 107 articles (103 studies) containing 40 tools were included in the review. There was substantial variation in the tools in terms of their aims and scope, structure, theoretical foundations, how they were developed, and the settings in which they have been used. The quantity of tools demonstrates the importance of assessing self-care and self-management. Consideration of the purpose, scope, and theoretical foundation should guide decisions about tools suitable for use in research and clinical practice.

## Keywords

aged, self-care, self-management, chronic conditions, measurement, scoping review

### *What this paper adds*

- This review provides an overview of existing tools used to measure self-care and self-management of chronic conditions by community-dwelling older people and describes their theoretical and practical characteristics.
- This review identified a large number of tools, reinforcing the importance of measuring self-care and self-management as a core outcome measure within research and healthcare settings.

### *Application of study findings*

- This review identifies disease-specific and non-disease-specific tools that measure self-care and self-management and describes characteristics that might make them suitable for use in older adults with chronic conditions and multimorbidity.
- Certain characteristics of existing tools (e.g., length, scope, and dimensionality) can inform the choice of use in specific research and healthcare settings.
- Tools capable of differentiating and quantifying the different dimensions of self-care and self-management can support person-centered assessment and management through the delivery of tailored interventions.

**Manuscript received:** November 1, 2022; **final revision received:** February 16, 2023; **accepted:** February 18, 2023.

<sup>1</sup>Caring Futures Institute, College of Nursing and Health Sciences, Flinders University, Adelaide, SA, Australia

<sup>2</sup>College of Humanities, Arts and Social Sciences, Flinders University, Adelaide, SA, Australia

### **Corresponding Author:**

Michael T. Lawless, Caring Futures Institute, College of Nursing and Health Sciences, Flinders University, Sturt Road, Adelaide, SA 5042, Australia.

Email: [michael.lawless@flinders.edu.au](mailto:michael.lawless@flinders.edu.au)

## Introduction

By 2030, the number of people aged 60 and over is projected to reach 1.4 billion, and by 2050, this figure is expected to rise to 2.1 billion (World Health Organization [WHO], 2022a). The rapid aging of the population has raised serious public health concerns related to the increasing prevalence of chronic conditions and multimorbidity (the co-occurrence of at least two chronic conditions). Multimorbidity in older adults is complex due to the potential overlap of physical and mental health conditions, polypharmacy, and frailty (Yarnall et al., 2017). Studies have estimated that between 55% and 98% of older adults have multimorbidity (Marengoni et al., 2011), and a recent meta-analysis found that 72% of those with frailty also have multimorbidity (Vetrano et al., 2019). Older people are therefore more likely to live with multiple chronic conditions associated with lower quality of life, reduced functional ability, increased healthcare utilization, and higher mortality (National Institute for Health and Care Excellence [NICE], 2016; Palladino et al., 2016; Ryan et al., 2015; Vogeli et al., 2007). Furthermore, fragmented or conflicting care can result in a higher treatment burden (Mair & May, 2014). As mortality rates decline and the population continues to age, managing chronic conditions and multimorbidity will become increasingly challenging for health systems (Fabbri et al., 2015; Kingston et al., 2018). Current models of care for older people have been criticized for not being sufficiently proactive and responsive to individuals' diverse needs, priorities, and environments (Cesari et al., 2022; Tinetti & Fried, 2004). In addition to strengthening healthcare system factors, principles of person-centered care and "minimally disruptive medicine" are increasingly seen as critical to improving care for older adults with chronic conditions and multimorbidity (American Geriatrics Society Expert Panel on Patient-Centered Care, 2016; May et al., 2009; Muth et al., 2014; Wallace et al., 2015).

Interventions to manage chronic conditions and multimorbidity should be tailored to address known issues, such as lack of care coordination, duplication, disease and treatment burden, and significant polypharmacy (i.e., taking 10 or more medications regularly) (Skou et al., 2022). Furthermore, these interventions should consider the goals and priorities of the person receiving care, involve informal caregivers, deliver care with a focus on interprofessional expertise and collaboration, and support self-care and self-management (Dineen-Griffin et al., 2019). Self-care and self-management support interventions are widely used in many single-disease programs and have the potential to improve outcomes for older people with chronic conditions and multimorbidity in primary care and community settings (Skou et al., 2022). However, the evidence to support their effectiveness is still limited (Smith et al., 2021).

## Defining and Measuring Self-Care and Self-Management

Self-care and self-management are critical for people with chronic conditions. It is estimated that more than 99% of the daily care for chronic conditions is performed by individuals and family carers (Riegel et al., 2017). Although both terms are widely used, they are associated with various conceptualizations and definitions, contributing to a lack of clarity and consensus (Barlow et al., 2002; Grady & Gough, 2014; Matarese et al., 2018; Richard & Shea, 2011; Van de Velde et al., 2019). Despite the lack of consensus, self-management is generally understood as a component of self-care occurring in the context of a recognized health condition with a degree of healthcare provider input (Matarese et al., 2018). Specifically, self-management is defined as "the intrinsically controlled ability of an active, responsible, informed, and autonomous individual to live with the medical, role, and emotional consequences of [their] chronic condition(s) in partnership with [their] social network and healthcare provider(s)" (Van de Velde et al., 2019, p. 10). Self-care is a broader concept that refers to "the ability of individuals, families, and communities to promote health, prevent disease, maintain health, and cope with or without the support of a healthcare provider" (WHO, 2022b). Self-care, according to the Middle-Range Theory of Self-Care of Chronic Illness (Riegel et al., 2012), consists of three dimensions: treatment adherence and health-promoting practices (self-care maintenance); behavior and condition monitoring (self-care monitoring); and managing signs and symptoms when they occur (self-care management). Commonly used models include the Chronic Disease Self-Management Program (Lorig et al., 1999), the Chronic Care Model (Wagner et al., 1996), and the Flinders Chronic Condition Management Program (Battersby et al., 2007). These models endorse the notion of commonality among chronic conditions and are intended for use with diverse conditions and populations.

Qualitative research and several systematic reviews have found that self-care and self-management consist of various aspects or domains, including the different skills, attitudes, and abilities that people use to address the challenges of living with chronic conditions (Auduly et al., 2012; Boehmer et al., 2016; Liddy et al., 2014; Schulman-Green et al., 2016; Van de Velde et al., 2019). However, scholarly literature that addresses the conceptual dimensions of self-care and self-management in older adults is relatively limited. The following characteristics have been identified as defining attributes of self-care and self-management of chronic conditions and multimorbidity among older people: using financial resources to manage chronic conditions; acquiring health-related education;

making use of social supports; responding positively to health changes; continuing engagement with the health system; and active participation in chronic condition management (Garnett et al., 2018). Additionally, (Lawless et al., 2021) identified seven core theoretical constructs that are essential for older adults' self-care and self-management: temporal and spatial context; stressors; personal resources; informal social resources; formal social resources; behavioral adaptations; and quality of life outcomes. Assessing the various domains of self-care and self-management can help individuals, carers, and healthcare providers identify the specific challenges experienced by older people with chronic conditions and multimorbidity so that appropriate resources, programs, and supports can be accessed, delivered, and evaluated effectively.

Developing and implementing appropriate tools to assess self-care and self-management can inform care delivery and is necessary to demonstrate the effectiveness of programs, policies, and interventions (Nichols et al., 2020). Instruments that accurately measure self-care and self-management can allow assessment of individuals' capacity to care for themselves so that the right level of support can be provided at the right time across healthcare and community settings (Coulter et al., 2015). Developing and implementing measurement tools should occur alongside coordinated efforts to redesign the structure and financing of long-term services and supports to provide comprehensive care for older people (Fulmer et al., 2021). Although several tools have been developed to measure self-care and self-management of chronic conditions in the general adult population (e.g., Jaarsma et al., 2003; Riegel et al., 2009; Toobert et al., 2000), there is a paucity of evidence on the characteristics and validity of tools used in older adults with chronic conditions and multimorbidity. It is unclear whether existing instruments measuring self-care and self-management validated in older adults with chronic conditions are structurally and conceptually similar or different. The number of disease-specific (i.e., intended for use based on a named disease) and non-disease-specific (i.e., intended for use across diagnoses) instruments can present challenges when selecting an appropriate instrument. Furthermore, there appears to be variation in their intended purpose, structure, theoretical foundations, how they have been developed, and the settings in which they have been used.

Previous reviews have examined disease-specific and non-disease-specific instruments measuring self-care or self-management (Ausili et al., 2014; Cameron et al., 2009; Caro-Bautista et al., 2013; Han et al., 2014; Hudon et al., 2021; Lu et al., 2015; Matarese et al., 2017; Packer et al., 2018; Sidani, 2011). These reviews provided insight into disease-specific and non-disease-specific instruments for measuring self-care and self-management in adult ( $\geq 18$  years) populations. However, they were not specific to older adults with chronic conditions or multimorbidity, who often

experience greater healthcare utilization, higher treatment burden, and geriatric syndromes such as frailty. To our knowledge, there have been no previous reviews of instruments measuring self-care and self-management by older adults living with chronic conditions. Hence, the aim of this scoping review was to identify the range of tools measuring self-care and self-management of chronic conditions by community-dwelling older adults ( $\geq 60$  years).

## Methods

### Design

We conducted a scoping review following the methodological framework developed by Arksey and O'Malley (2005) and advanced by others (Daudt et al., 2013; Levac et al., 2010). The review consisted of five steps: (1) identifying the research question; (2) identifying relevant studies; (3) selecting studies based on pre-defined criteria; (4) charting the data; and (5) collating, summarizing, and reporting the results. A sixth step, consultation, is considered optional when the research team is multidisciplinary. Our research team includes a range of backgrounds (cancer, cardiovascular and older peoples' nursing, psychology, and sociology) with expertise in self-care and self-management theory, fundamental nursing care, integrated care models, survivorship, and implementation science. Scoping reviews are conducted to examine the type and range of evidence available on topic, clarify key concepts and definitions, and identify knowledge gaps in the literature to inform future research (Munn et al., 2018). A scoping review was chosen due to the large number and variability of tools in the literature. We used the Preferred Reporting Items for Systematic Reviews and Meta-analyses for Scoping Reviews (PRISMA-ScR) (Tricco et al., 2018) to guide reporting and enhance fidelity. We registered an a priori protocol with the Open Science Framework (Lawless, 2022).

### Identifying the Research Question

The objective of this scoping review was to identify and describe the range of tools available to measure self-care and self-management of chronic conditions by older adults (see [Supplementary File 1](#) for definitions of key terms). The specific review question was "what tools are available to measure self-care and self-management of chronic conditions by community-dwelling older adults?" We identified two specific objectives:

1. To identify tools that measure self-care and self-management by people (aged  $\geq 60$  years) that can be used for assessment and evaluation in clinical practice and research.
2. To map the characteristics, including their aims and scope, structure, theoretical foundations, how they have been developed, and the settings in which they have been used.

### Identifying Relevant Studies

The search aimed to locate peer-reviewed studies published between January 2002 and March 2022. This date range was chosen to ensure relevance to current self-care and self-management interventions and theory. We developed the search strategy based on recommendations for conducting scoping reviews, previous examples from the literature, and the advice of a university librarian. We searched six electronic databases: CINAHL, Scopus, ERIC, PsycINFO, MEDLINE/PubMed, ProQuest. We chose these databases for their breadth and diversity of disciplines represented. We used a combination of Medical Subject Heading (MESH) terms and keyword searches were to identify publications meeting the inclusion criteria. The search strategy used for MEDLINE/PubMed is available in [Supplementary File 2](#).

### Selecting Studies

The inclusion criteria are shown in [Supplementary File 3](#). We included studies reporting on the development, validation, or testing of tools measuring self-care or self-management in adults aged  $\geq 60$  years. This age was chosen based on the [United Nations \(2019\)](#) and the [WHO \(2022a\)](#) definition of older people as individuals aged over 60 years. Only studies with a specified theoretical foundation were included in this review based on the expectation that complex population health interventions require robust and explicit theorization for successful implementation and to function as expected in terms of change mechanisms ([Hastings et al., 2020](#); [Moore et al., 2021](#)). Articles reporting on older adults with chronic conditions that did not specify a theoretical underpinning were excluded. Included chronic conditions were identified from a list of prevalent conditions published by the Office of the Assistant Secretary for Health (OASH) in the United States Department of Health and Human Services ([Goodman et al., 2013](#)). From this list, we selected 10 prevalent chronic conditions associated with significant morbidity and mortality in older people worldwide: coronary artery disease, hypertension, heart failure, stroke, arthritis, asthma, cancer, chronic kidney disease (CKD), chronic obstructive pulmonary disease (COPD), and type 2 diabetes. These conditions were among 20 chronic conditions selected by OASH for a standard classification scheme ([Goodman et al., 2013](#)). They are prevalent among older people across low-, middle-, and high-income countries ([Ofori-Asenso et al., 2019](#); [Vancampfort et al., 2017](#)) and are frequently investigated in research on self-care and self-management interventions for adult patients ([Riegel et al., 2021](#)). From this list, we excluded studies that focused exclusively on severe mental illness or advanced dementia as these individuals often have limited capacity to participate actively in self-care or self-management in a manner consistent with current definitions. Articles reporting on instruments developed for use with older adults ( $\geq 60$  years) without the identified chronic

conditions were excluded. We excluded studies that reported exclusively on inpatient, hospital, residential aged care, or palliative care settings. No limitation was placed on the upper age, gender, ethnicity, or geographical location of participants.

Following the search, all retrieved references were imported into Endnote (Clarivate Analytics, PA, USA) and Covidence systematic review platform (Veritas Health Innovation, Melbourne, Australia) and duplicates were removed. Prior to title and abstract screening, the first four authors discussed a representative sample of studies to ensure consistency among their interpretation of the eligibility criteria. The full texts were assessed by two reviewers against the inclusion criteria using a standardized screening form. Studies that appeared to meet the inclusion criteria were retrieved in full text and a second meeting was held to verify a random selection of each reviewer's studies and to discuss any studies about which a reviewer was unsure, after which a group decision was made to exclude or include. Finally, we searched the reference list of each included article to identify additional studies. Consistent with scoping review methodology, critical appraisal of the studies was not undertaken ([Levac et al., 2010](#)).

### Charting the Data

Two reviewers independently conducted data extraction in Covidence and compared the completed tool to maintain consistency during the extraction process. We used a standardized data extraction tool ([Supplementary File 4](#)) to extract information about the study, including the setting, chronic conditions under study, and the tool used. Data extracted in relation to the tool included its aims and scope, definitions of self-care and self-management, theoretical foundation, method of development, structure, and contexts of use. Disagreements or inconsistencies were resolved through group discussion.

### Collating, Summarizing, and Reporting the Results

The extracted data were mapped and summarized using a narrative descriptive approach to expose commonality and heterogeneity among the studies ([Barnett-Page & Thomas, 2009](#)). Results are reported following the PRISMA-ScR guidelines ([Tricco et al., 2018](#)). The PRISMA-ScR checklist is available in [Supplementary File 5](#).

### Results

Database searching identified 1891 articles and searching the reference lists of included studies identified an additional 19 articles. After duplicates were removed and screening of titles and abstracts, 628 articles were assessed for eligibility and 540 were excluded based on the inclusion criteria. A total of 107 articles from 103 studies were included in the final review

(Figure 1). In total, 40 measurement instruments were included in the review (Table 1). Of the 40 tools, 23 (57.5%) were disease-specific. The most common conditions were type 2 diabetes (20.0%,  $n = 8$ ), heart failure (7.5%,  $n = 3$ ), hypertension (7.5%,  $n = 3$ ), COPD (7.5%,  $n = 3$ ), and CKD (7.5%,  $n = 3$ ). Seventeen tools (42.5%) were non-disease-specific, meaning that they were developed to be applicable irrespective of diagnosis. Eleven tools (27.5%) were used to

measure self-care or self-management in people with multimorbidity (i.e.,  $\geq 2$  chronic conditions). Supplemental Table 1 provides a summary of the included studies.

### Frequency of Use of Each Tool

The Self-Care of Heart Failure Index (SCHFI) was the most frequently used tool among the included studies (15.0%  $n = 16$ ),




Figure 1. Prisma flow chart.

Table 1. Summary of included tools.

Name of Tool and Author(s)	What the Tool Measures	Definition of Self-Care/Self-Management	Disease-Specific or Non-Disease-Specific	Theoretical Underpinning	Structure	Settings in which the Tool has Been Used
Appraisal of Self-Care Agency (ASAS) (Evers et al., 1993)	Operability of self-care agency	Definition [self-care agency]: "A person's capability or power to perform self-care operations"	Non-disease-specific	Self-care deficit nursing theory	24 items (unidimensional)	Primary care
Arthritis Self-Efficacy Questionnaire (ASES) (Lorig et al., 1989)	Self-efficacy for arthritis management	Definition [self-efficacy]: "One's belief that one can perform a specific behavior or task in the future"	Disease-specific (arthritis)	None specified	2 subscales, 11 items	Outpatient clinics
Chronic Kidney Disease Self-Efficacy Instrument (CKD-SE) (Lin et al., 2012)	Self-efficacy for chronic kidney disease management	Definition [self-efficacy]: "An individual's confidence in their ability to overcome barriers in order to perform disease-specific self-management behaviors"	Disease-specific (chronic kidney disease)	None specified	4 subscales, 25 items	Medical centers
Chronic Kidney Disease Self-Management Instrument (CKD-SM) (Lin et al., 2013)	Chronic kidney disease self-management	Definition [self-management]: "The full range of activities undertaken by a person with a chronic condition, ranging from the preventive activities undertaken by healthy people [at] home to the day-to-day tasks undertaken by an individual to manage symptoms, treatments, consequences and life-style changes associated with chronic conditions"	Disease-specific (chronic kidney disease)	None specified	4 subscales, 29 items	Medical centers
COPD Self-Efficacy Scale (CSES) (Wigal et al., 1991)	Self-efficacy for managing COPD	Definition [self-efficacy]: "Personal convictions people have regarding whether or not they feel they can successfully execute particular behaviors in order to produce certain outcomes"	Disease-specific (COPD)	Self-efficacy theory	5 subscales, 34 items	Hospital, community care centers
Diabetes Management Self-Efficacy Scale (DMSES) (Bijl et al., 1999)	Self-efficacy for diabetes management	Definition [self-efficacy]: "People's judgement of their capabilities to organize and execute courses of action required to attain designated types of performances"	Disease-specific (diabetes)	Social cognitive theory	20 items (unidimensional)	Outpatient clinics

(continued)

Table 1. (continued)

Name of Tool and Author(s)	What the Tool Measures	Definition of Self-Care/Self-Management	Disease-Specific or Non-Disease-Specific	Theoretical Underpinning	Structure	Settings in which the Tool has Been Used
Diabetes Mellitus Type 2 Self-Efficacy Scale (DMSES) adapted version. (Sol et al., 2006)	Self-efficacy for diabetes management	Implied definition [self-efficacy]: "A person's confidence to carry out behavior necessary to reach a desired goal, is an important precondition for successful self-management and behavior change"	Disease-specific (diabetes)	Social cognitive theory	9 items (unidimensional)	Outpatient clinics
Diabetes Self-Efficacy Scale (DSES) (Lorig et al., 2001a)	Self-efficacy for diabetes management	None provided	Disease-specific (diabetes)	Self-efficacy theory	8 items (unidimensional)	Primary care
Diabetes Self-Management Instrument Short Form (DSMI-20) (Lee et al., 2016)	Diabetes self-management	Definition [self-management]: "Individual(s) can take actions to: create a healthy lifestyle for themselves; meet their social, emotional and psychological needs; care for their long-term condition; and to prevent further illness or accidents"	Disease-specific (diabetes)	None specified	4 subscales, 20 items	Community
Diabetes Self-Management Questionnaire (DSMQ) (Schmitt et al., 2013)	Diabetes self-management activities	None provided	Disease-specific (diabetes)	None specified	4 subscales, 16 items	Primary care
European Heart Failure Self-care Behavior Scale (EHFScBS) (Jaarsma et al., 2003)	Heart failure self-care behaviors	Definition [self-care]: "The decision and strategies undertaken by the individual in order to maintain life, healthy functioning and wellbeing. Self-care behavior of a person can be universal (needed by every person), health-deviated (i.e., arises from health problems) and/or developmental (i.e., arises from a specific stage of life)"	Disease-specific (heart failure)	Self-care deficit nursing theory	12 items (unidimensional)	Primary care

(continued)

Table 1. (continued)

Name of Tool and Author(s)	What the Tool Measures	Definition of Self-Care/Self-Management	Disease-Specific or Non-Disease-Specific	Theoretical Underpinning	Structure	Settings in which the Tool has Been Used
Exercise of Self-Care Agency Scale (ESCA) (Kearney & Fleischer, 1979)	Self-care agency	Definition [self-care agency]: "Four subconstructs are attributed to an individual's self-care agency: (a) Active versus passive response to situations; (b) an individual's motivations; (c) the knowledge base of the individual; and (d) the person's sense of self-worth"	Non-disease-specific	Self-care deficit nursing theory	4 subscales, 43 items	Health centers
FERUS26 (Jack, 2007)	Self-management skills and resources	Definition [self-management]: "A behavioral and cognitive strategy that can help patients who suffer from chronic conditions to structure their behavior, to learn problem-solving skills and how to achieve effective disease management goals"	Non-disease-specific	None specified	6 subscales, 26 items	Primary care
Health Education Impact Questionnaire (heiQ) (Osborne et al., 2007)	Impact of patient education on self-management	Implied definition [patient education programs]: "At a minimum, such programs should improve both health knowledge and health literacy, which increase an individual's capacity to obtain, process, and understand basic health information and services needed to make appropriate health-related decisions"	Non-disease-specific	Program Logic Model	8 subscales, 42 items	Arthritis foundations (community)
Heart Failure Decision Influence Inventory (HFDII) (Hicks & Holm, 2003)	Heart failure self-management decision-making	Implied definition [self-management]: "Appropriate self-management may slow disease progression and prevent repeated and expensive hospital readmissions...but adherence to therapeutic regimens varies greatly among individuals with heart failure"	Disease-specific (heart failure)	Naturalistic decision-making	5 subscales, 56 items	Heart failure clinic

(continued)



Table 1. (continued)

Name of Tool and Author(s)	What the Tool Measures	Definition of Self-Care/Self-Management	Disease-Specific or Non-Disease-Specific	Theoretical Underpinning	Structure	Settings in which the Tool has Been Used
Hypertension Self-Care Questionnaire (HSCQ) (Eghballi-Babadi et al., 2019)	Hypertension self-care	Definition [self-care]: "The ability of individuals, families and communities to promote health, prevent diseases and maintain health for fighting disease and disability with or without the support of a physician"	Disease-specific (hypertension)	Self-care deficit nursing theory	5 subscales, 16 items	Community
Mobility and Self-Care-Patient (MOSES-Patient) (Farin et al., 2007)	Mobility, self-care, and domestic life	None provided	Non-disease-specific	World Health Organization International Classification of Functioning, Disability and Health framework	12 subscales, 58 items	Rehabilitation clinics
Partners in Health Scale (PIH) (Battersby et al., 2003)	Chronic condition self-management	Definition [self-management]: "Self-management involves the individual working in partnership with their carer(s) and health professionals so that (s)he can: 1. Know their condition and various treatment options. 2. Negotiate a plan of care; (i.e., Care Plan). 3. Engage in activities that protect and promote health. 4. Monitor and manage the symptoms and signs of the condition(s). 5. Manage the impact of the condition on physical functioning, emotions and interpersonal relationships"	Non-disease-specific	None specified	12 items (unidimensional)	Community health services, elderly care residential setting

(continued)

Table 1. (continued)

Name of Tool and Author(s)	What the Tool Measures	Definition of Self-Care/Self-Management	Disease-Specific or Non-Disease-Specific	Theoretical Underpinning	Structure	Settings in which the Tool has Been Used
Patient Activation Measure (PAM) (Hibbard et al., 2004)	Patient activation	Definition [patient activation]: "Those who are activated believe patients have important roles to play in self-managing care, collaborating with providers, and maintaining their health. They know how to manage their condition and maintain functioning and prevent health declines; and they have the skills and behavioral repertoire to manage their condition, collaborate with their health providers, maintain their health functioning, and access appropriate and high-quality care"	Non-disease-specific	None specified	22 items (unidimensional)	Primary care, Veteran community organizations
Patient Activation Measure-13 (PAM-13) (Hibbard et al., 2005)	Patient activation	None provided	Non-disease-specific	None specified	13 items (unidimensional)	Primary care, outpatient clinics, community
Chronic Illness Care (PACIC) (Glasgow et al., 2005a)	Care provision aligned with Chronic Care Model (CCM)	None provided	Non-disease-specific	Chronic Care Model	5 scales, 20 items	Primary care, community
Patient Assessment of Chronic Illness Care 5A (PACIC 5A) Questionnaire (Glasgow et al., 2005b)	Care provision aligned with Chronic Care Model (CCM)	None provided	Non-disease-specific	Chronic Care Model	5 subscales, 26 items	Primary care
Patient Assessment of Chronic Illness Care short version (PACIC-S) (Cramm & Nieboer, 2012)	Care provision aligned with Chronic Care Model (CCM)	None provided	Non-disease-specific	Chronic Care Model	5 subscales, 11 items	Diabetes management programs, primary care

(continued)

**Table 1.** (continued)

Name of Tool and Author(s)	What the Tool Measures	Definition of Self-Care/Self-Management	Disease-Specific or Non-Disease-Specific	Theoretical Underpinning	Structure	Settings in which the Tool has Been Used
Perceived Competence Scale (PCS) (Williams et al., 2004)	Perceived competence for diabetes self-management	None provided	Disease-specific (diabetes)	Self-determination theory	4 items (unidimensional)	Community
Perceived Competence Scale for Hypertension Self-Management (Dye et al., 2003)	Perceived competence for hypertension self-management	None provided	Disease-specific (hypertension)	Self-determination theory	4 items (unidimensional)	Community
Revised Heart Failure Self-Care Behavior Scale (Artinian et al., 2002)	Heart failure self-care behaviors	Definition [self-care]: "The practice of activities that individuals initiate and perform on their own behalf in the interest of maintaining life, health, continuing personal development, and well-being"	Disease-specific (heart failure)	Self-care deficit nursing theory	5 subscales, 29 items	Hospital, cardiology clinic of Veterans Affairs medical centre
Self-Care in Chronic Obstructive Pulmonary Disease Inventory (SC-COPDI) (Matarrese et al., 2020)	Self-care of COPD	Definition [self-care]: "A process of maintaining health through health-promoting and managing illness practices. Self-care is performed in both healthy and ill states, with specific behaviors captured in three core concepts: Self-care maintenance, self-care monitoring, and self-care management"	Disease-specific (COPD)	Middle-range theory of self-care of chronic illness	3 subscales, 32 items	Hospital units, outpatient clinics, primary care practices, rehabilitation centers
Self-Care of Chronic Illness Inventory (SC-CII) (Riegel et al., 2018)	Self-care of chronic illness	Implied definition [self-care]: "Self-care is performed in both healthy and ill states. When one has a chronic illness, self-care addresses the behavioral requirements to maintain stability and control symptoms"	Non-disease-specific	Middle-range theory of self-care of chronic illness	3 subscales, 20 items	Community, inpatient and outpatient settings

(continued)

Table 1. (continued)

Name of Tool and Author(s)	What the Tool Measures	Definition of Self-Care/Self-Management	Disease-Specific or Non-Disease-Specific	Theoretical Underpinning	Structure	Settings in which the Tool has Been Used
Self-Care of Coronary Heart Disease Inventory (SC-CHDI) (Vaughan Dickson et al., 2017)	Self-care of coronary heart disease	Definition [self-care]: "A naturalistic decision-making process of maintaining health through health-promoting practices and managing illness"	Disease-specific (coronary heart disease)	Middle-range theory of self-care of chronic illness	3 subscales, 22 items	Outpatient clinical services, cardiac rehabilitation programs
Self-Care of Diabetes Inventory (SCODI) (Ausili et al., 2017)	Diabetes self-care	Definition [self-care]: "A process of maintaining health through health promoting practices and managing illness"	Disease-specific (diabetes)	Middle-range theory of self-care of chronic illness	4 subscales, 40 items	Diabetes outpatient services
Self-Care of Heart Failure Index (SCHFI) (Riegel et al., 2004)	Self-care of heart failure	Definition [self-care]: "A naturalistic decision-making process involving the choice of behaviors that maintain physiologic stability (self-care maintenance) and the response to symptoms when they occur (self-care management)"	Disease-specific (heart failure)	Situation-specific theory of heart failure self-care	v.4 = 15 items, 3 subscales SCHFI v6.2 = 22 items, 3 subscales; SCHFI v7.2 = 29 items, 3 subscales	Specialist medical centers, cardiology clinics, community, cardiac rehabilitation programs, cardiac support groups, outpatient heart failure clinics, primary care, inpatient settings, ambulatory cardiovascular clinics
Self-Care of Hypertension Inventory (SC-HI) (Dickson et al., 2021)	Self-care of hypertension	Definition [self-care]: "A process of maintaining health with health-promoting practices and monitoring and managing the signs and symptoms of a chronic condition"	Disease-specific (hypertension)	Middle-range theory of self-care of chronic illness	3 subscales, 24 items	Community
Self-Care Self-Efficacy Scale (SCSES) (Yu et al., 2021)	Self-efficacy for self-care	Definition [self-care]: "A deliberate process where patients perform behaviors such as adherence to treatment, symptom recognition, and response to minimize complications and maintain health"	Non-disease-specific	Middle-range theory of self-care of chronic illness	10 items (unidimensional)	Inpatient and outpatient settings
Self-Care Self-Efficacy Scale COPD (SCES-COPD) (Matarese et al., 2020)	Self-efficacy for COPD self-care	Definition [self-care]: "A process of maintaining health through health-promoting and managing illness practices"	Disease-specific (COPD)	Middle-range theory of self-care of chronic illness	7 items (unidimensional)	Hospital units, outpatient clinics, primary care practices, rehabilitation centers

(continued)

Table 1. (continued)

Name of Tool and Author(s)	What the Tool Measures	Definition of Self-Care/Self-Management	Disease-Specific or Non-Disease-Specific	Theoretical Underpinning	Structure	Settings in which the Tool has Been Used
Self-Efficacy for Managing Chronic Disease 6-item scale (SEMCD-6) (Lorig et al., 2001b)	Self-efficacy for managing chronic disease	Definition [self-efficacy]: "Confidence in one's ability to manage different aspects of one's health functioning"	Non-disease-specific	Self-efficacy theory	6 items (unidimensional)	Primary care, community, clinical physiotherapy, senior centers, telehealth programs, chronic disease self-management programs and workshops
Self-Efficacy for Managing Chronic Disease Scales (SEMCD) (Lorig et al., 1996)	Self-efficacy for managing chronic disease	Definition [self-efficacy]: "The perceived capability to perform specific actions required to achieve concrete goals"	Non-disease-specific	Self-efficacy theory	10 subscales, 33 items	Community, primary care, general internist practices, health clinics providing services for older adults
Self-Management Ability Scale-30 (SMAS-30) (Schuurmans et al., 2005)	Self-management ability	Implied definition [self-management]: "The successful self-management of aging concerns the ways in which people are able to realize well-being and in particular, how they can sustain it, even when their resources decline"	Non-disease-specific	Successful self-management of aging theory	6 subscales, 30 items	Primary care, pulmonary rehabilitation assessment center
Self-Management Ability Scale-Short version (SMAS-S) (Gramm et al., 2012)	Self-management ability	Implied definition [self-management abilities]: "Self-management abilities (internal resources) are needed to manage external resources in such a way that physical and social well-being are maintained or restored when lost"	Non-disease-specific	Successful self-management of aging theory	6 subscales, 18 items	Primary care, community
Self-Management behavior Questionnaire (SMBQ) (Curtin et al., 2008)	Self-management behavior	Implied definition [self-management]: "Self-care is the 'action' dimension of self-management and is based on the premise that the best outcomes of health care result when patients are actively involved in their own care"	Disease-specific (chronic kidney disease)	Social cognitive theory	5 subscales, 37 items	Outpatient clinics
Summary of Diabetes Self-Care Activities (SDSCA) (Toobert et al., 2000)	Diabetes self-care behaviors	Implied definition [self-care]: "Diabetes self-care includes a range of activities...and it is now well established that these different components do not correlate highly"	Disease-specific (diabetes)	None specified	5 subscales, 11 items	Primary care, community pharmacies, outpatient clinics

followed by the Self-Efficacy to Manage Chronic Disease 6-item scale (SEMCD-6) (13.1%,  $n = 14$ ), the Patient Assessment of Chronic Illness Care (PACIC) (10.3%,  $n = 11$ ), the Self-Efficacy to Manage Chronic Disease scales (SEMCD) (9.3%,  $n = 10$ ), and the Summary of Diabetes Self-Care Activities (SDSCA) (7.5%,  $n = 8$ ). Twenty-two (55.0%) of the 40 tools were used in only one study.

### Aims and Scope of Included Tools

The aims and scope of the included tools (i.e., what the tool is intended to measure) is reported in Table 1. Thirteen of the 40 tools (32.5%) measured self-care or elements of self-care, such as self-care agency (e.g., Exercise of Self-Care Agency scale [ESCA]) and self-care behaviors (e.g., European Heart Failure Self-Care Behavior Scale [EHFScBS]). Eleven tools (27.5%) measured self-management or elements of self-management, including self-management skills and resources (e.g., FERUS26), and perceived competence for self-management (e.g., Perceived Competence Scale [PCS]). Ten tools (25.0%) measured self-efficacy related to chronic disease self-care or self-management, such as the Self-Care Self-Efficacy Scale (SCSES). The remainder of the tools measured patient activation (e.g., Patient Activation Measure-13 [PAM-13]) and care provision aligned with the Chronic Care Model (e.g., PACIC). Only three tools (7.5%) (ASAS, SMAS-30, SMAS-S) were developed to measure self-care or self-management in older adults.

Twenty-three tools (57.5%) included a clearly labeled definition of self-care, self-management, or related concepts such as self-efficacy, which provided insight into the intended focus of the tool (Table 1). Eight tools (20%) alluded to definitions or used vague language without clearly identifying the construct being measured (implied definitions). Nine tools (22.5%) did not provide a definition. Definitions ranged from specific and disease-focused (e.g., “*self-monitoring of blood glucose, eating a low-saturated-fat diet, and checking one’s feet*”; SDSCA) to broader definitions (e.g., “*a person’s capability or power to perform self-care operations*”; ASAS). Definitions provided by developers of 17 tools (42.5%) focused on behavioral/medical strategies, including actions undertaken to maintain a healthy lifestyle (with and without a chronic condition) and manage medical aspects of chronic conditions (e.g., monitoring/managing symptoms, treatment adherence). In comparison, those used by authors of 15 tools (37.5%) focused on cognitive/decision-making strategies, including the intellectual processes used for decision-making or to develop self-care and self-management skills. Definitions provided by the developers of the Partners in Health Scale (PIH) and the Health Education Impact Questionnaire (heiQ) focused on resource utilization and health navigation.

We originally intended to map and synthesize the items in the scales and subscales of the tools to assess their scope and breadth, as well as the extent to which each tool measures discrete or overlapping concepts. However, this proved

problematic because of the variability in how the tools described and operationalized self-care and self-management. For example, some items measured individual knowledge, skills, and attitudes related to maintaining health and managing chronic conditions, such as “*How confident are you that you can keep the fatigue caused by your disease from interfering with the things you want to do?*” (SEMCD-6). Other items measured behaviors performed by the person who is ill to maintain health, limit the risk of illness, and control chronic disease, such as “*How often do you monitor for medication side-effects?*” (Self-Care of Chronic Illness Inventory [SC-CII]) and “*I weigh myself every day*” (EHFScBS). In several instances, studies did not report the label of subscales or describe the scale in sufficient detail to allow a synthesis.

### Methods of Tool Development

Items included in the tools were developed based on a review of the academic literature, clinical guidelines, theoretical constructs, or existing validated instruments ( $n = 18$ ), qualitative methods, such as focus groups and interviews ( $n = 7$ ), and assessment of content validity by an expert panel ( $n = 17$ ). Items in four tools were adapted from previous tools or subscales embedded within previous tools. The method of development was not reported for eight tools. Items in less than half of the tools ( $n = 19$ ) were developed by researchers in combination with clinicians or other experts, such as specialist physicians ( $n = 8$ ), nurses ( $n = 8$ ), case managers or service coordinators ( $n = 3$ ), diabetes educators ( $n = 3$ ), and physiotherapists ( $n = 2$ ). Five tools had some level of patient/consumer involvement in item development or refinement, such as assessing items for relevance, comprehensibility, and comprehensiveness (e.g., SC-COPDI).

### Theoretical Foundation

Theoretical foundation was defined as any reference to, or application of, a model, theory, or framework to inform or underpin the development of the measurement tool. Most of the tools (75.0%,  $n = 30$ ) were based on a named theory, model, or framework, as stated by the original authors of the tool. The most common theories, models, and frameworks underpinning the tools were the Middle-range Theory of Self-Care of Chronic Illness (17.5%,  $n = 7$ ), Self-Efficacy Theory/Social Cognitive Theory (17.5%,  $n = 7$ ), Self-Care Deficit Nursing Theory (12.5%,  $n = 5$ ), and the Chronic Care Model (7.5%,  $n = 3$ ). 10 tools did not explicitly specify a theory, model, or conceptual framework.

### Contexts of Use

The 103 studies included data collected from the USA ( $n = 36$ ), Italy ( $n = 10$ ), Australia ( $n = 10$ ), Canada ( $n = 9$ ), Germany ( $n = 9$ ), the Netherlands ( $n = 8$ ), China ( $n = 7$ ), and the UK ( $n = 6$ ). Countries with five or less studies included

Switzerland ( $n = 3$ ), Iran ( $n = 2$ ), South Korea ( $n = 2$ ), Brazil ( $n = 2$ ), Mexico ( $n = 2$ ), Finland ( $n = 1$ ), France ( $n = 1$ ), Singapore ( $n = 1$ ), Spain ( $n = 1$ ), and Thailand ( $n = 1$ ). Two studies included data obtained from more than one country. Of the 40 tools, 12 (30.0%) were used in more than one country. Of the 40 tools, most were used in outpatient clinics ( $n = 20$ ), community ( $n = 17$ ), primary care ( $n = 17$ ), rehabilitation ( $n = 6$ ), and hospital or inpatient settings ( $n = 6$ ). Eighteen tools were used in more than one setting.

### Structure of the Tools

The length of the tools ranged from four items (PCS) to 58 items (MOSES-Patient). The number of subscales or domains in each tool ranged from one to 12. Thirteen tools (32.5%) were unidimensional, meaning that they measured a single underlying construct or dimension (e.g., self-efficacy). Many of the tools had multiple variations or iterations. A “variation” was defined as a revision or modification to the structure or administration of a tool such that the number of items (i.e., number of items administered), language, scales/subscales, or scoring modality differed from the published version. Short forms of existing tools reported in a peer-reviewed publication were considered a separate tool rather than a variation. In addition, minor changes in wording, such as to reflect an accompanying person (e.g., family member) completing the tool, or the administration of an individual subscale from a tool, were not considered indicative of variation. Of the 40 tools, six had more than one variation.

### Discussion

Our review identified 40 tools and found that most were developed to measure a specific construct or assess self-care or self-management of a specific condition. Only three tools were developed specifically for adults aged over 60 years with chronic conditions. Although several tools have been validated in samples of older adults, based on this review, none of the identified tools can be fully recommend for use to measure self-care or self-management among older people with chronic conditions and multimorbidity. Like earlier reviews (e.g., Hudon et al., 2021; Packer et al., 2018), we found considerable variation in the definitions of self-care and self-management used, reflecting ongoing conceptual inconsistency and fragmentation within the literature. Although many authors stated that their tool measured self-care or self-management, closer inspection revealed that many tools measured related constructs such as self-efficacy or patient activation, replicating previous studies. This lack of conceptual precision raises questions about whether tools claiming to measure self-care and self-management are measuring consistently defined constructs. The different ways in which self-care and self-management are conceptualized and operationalized might partly explain the mixed evidence of effectiveness of interventions in chronic illness (Lee et al.,

2022). The terms self-care and self-management continue to be used interchangeably, despite efforts over the last decade to delineate the concepts (Grady & Gough, 2014; Matarese et al., 2018; Richard & Shea, 2011; Van de Velde et al., 2019). Instruments developed for specific conditions often use the terminology that is standard within that scientific community. This reflects the reality of clinical practice, which is often compartmentalized into silos. For example, self-management is predominantly used in diabetes, COPD, coronary artery disease, arthritis, and asthma, whereas self-care is the term predominantly used in heart failure. Riegel et al. (2021) note that the inconsistency might be traced to early adoption of the self-management theories of Bandura (1997) and Lorig et al. (1999) versus early adoption of the self-care theory of Orem (1985). Many authors have since written about the similarities and differences between concepts (e.g., Matarese et al., 2018). Reaching agreement on the meaning and use of the terms may contribute to bridging disciplinary silos currently precluding effective knowledge exchange (Jaarsma et al., 2020).

Although an increasing proportion of older patients experience multimorbidity and complex needs, we found that disease-specific instruments were more common than non-disease-specific instruments, a finding reported in previous studies (Packer et al., 2018). Thirteen of the included tools were unidimensional, while the remainder mostly measured specific dimensions of self-care and self-management, such as symptom monitoring or treatment adherence. Several reviews demonstrate that self-care and self-management comprise multiple interacting domains, including the intrinsic and extrinsic resources that people use to address the daily challenges of living with chronic conditions (Auduly et al., 2012; Boehmer et al., 2016; Lawless et al., 2021; Van de Velde et al., 2019). Although unidimensional instruments might be beneficial due to their brevity and provide clinically useful information, they might have limited ability to identify specific care needs, as well as intrinsic and extrinsic resources, to guide care tailored to patients' individual circumstances. Other instruments provide a single composite score despite analysis providing evidence of underlying multidimensionality. Whilst single-score instruments might be valid, reliable, and brief, they have limited utility in identifying individual patients' self-care and self-management support requirements or in personalized care planning (Coulter et al., 2015; Packer et al., 2020). Instruments capable of assessing the various domains of self-care and self-management are needed to move beyond “one size fits all” approaches that are responsive to people's diverse and changing needs. Research into the active ingredients of interventions and advancing a person-centered approach also depends on the ability to distinguish and measure separate domains of self-care and self-management.

Developing accurate and comprehensive measures to assess self-care and self-management by individuals and support

across health services has the potential to promote integrated and personalized care and support for older people with multimorbidity (Keddy et al., 2021; Nichols et al., 2020; Shepherd et al., 2022; WHO, 2015). Personalized care planning, defined as a series of discussions between a patient and a health professional to clarify goals, options, and preferences and develop an agreed plan of action, embodies the core principles of person-centeredness and shared decision-making (Coulter et al., 2015). The aim of personalized care planning is to support individuals and carers to self-manage their health and wellbeing, typically using a combination of behavior change techniques (e.g., goal setting, action planning, health coaching, motivational interviewing) to achieve collaborative outcomes (Ahmed et al., 2021). Personalized care planning provides an opportunity to advance a proactive and person-centered approach that expands the traditional medical focus of care to a more comprehensive, socially oriented approach. Personalized care plans should contain various components including a package of multi-component interventions, management of underlying chronic diseases and geriatric syndromes, social care and support, and support for self-care and self-management. To deliver person-centered and feasible treatment programs, healthcare providers therefore need to appreciate the limits and possibilities of people's capacity to enact self-care and self-management across different domains (e.g., physical, personal, emotional, social, financial, environmental) (Boehmer et al., 2018). Although self-care and self-management are components of personalized care planning, it remains unclear whether existing self-care and self-management tools are suitable as part of integrated person-centered assessment and pathways for older people.

### *Strengths and Limitations*

This review used the rigorous methodology developed by Arksey and O'Malley (2005) and advanced by Levac et al. (2010). Searching of reference lists allowed us to identify additional articles and instruments. A limitation of this review was the difficulty in locating and classifying relevant instruments resulting from the lack of accepted conventions for naming tools and the various repositories for outcome measures. Some tools with unique names were found to be variations or duplicates of another tool with a different name, which had undergone revisions or modifications (e.g., to apply to a specific chronic condition). Moreover, as noted by Packer et al. (2018), unless a measure has been commercialized, often it is hard to determine whether versions modified for other languages or cultures exist, or which is the most recent version of the tool. These difficulties were mitigated by referring to the original articles describing the development and/or validation of the tool and cross-checking tables and independently completed data extraction forms. It is possible that some tools were not included in the review because we focused on studies reporting on older adults with prevalent

chronic conditions and excluded studies that reported exclusively on asymptomatic, acute, and psychiatric diagnoses; we excluded studies that reported exclusively on inpatient, hospital, residential aged care, or palliative care settings; and we only included tools where items could be located and viewed. However, the large number of studies and tools identified through database searching allowed us to describe the extent and range of the existing literature. The variability of the items in the tools, as explained above, meant that we were unable to synthesize the items in a meaningful and precise way; this should be a key aim of future research.

### *Future Directions*

Building on the current review, further research is needed to assess the content and quality of the identified instruments to clarify which measures are appropriate for the target population of older people with chronic conditions and multimorbidity. Like previous studies (e.g., Hudon et al., 2021; Packer et al., 2018), the definitions and theoretical foundations used to develop the instruments included in this review were varied, indicating that the concepts of self-care and self-management require further clarification and refinement. Although self-care and self-management can be defined in various ways across chronic conditions and cultural groups, the use of consistent and coherent definitions can provide a shared language for communication across services and enable more robust measurement. Clearer terminology may also facilitate the implementation of self-care and self-management support in person-centered, integrated care for older adults with chronic conditions and multimorbidity. A shift towards embedding self-care and self-management support across health services requires the development of comprehensive, validated measures to demonstrate the effectiveness of policies, programs, and interventions. However, many existing measures are not designed to evaluate the effectiveness of complex self-care and self-management interventions with multiple interacting components. Theory-based approaches, such as the Medical Research Council framework, provide comprehensive and systematic methods for developing and evaluating complex interventions (Moore et al., 2015; O'Cathain et al., 2019). In addition, complex systems thinking could be used to gain insight into the complexity of chronic disease self-care and self-management and identify promising leverage points for interventions (Baugh Littlejohns & Wilson, 2019). These methods could be combined with a partnership approach, in which end users participate actively throughout the development process to understand the perspectives and wider context of the potential target population, including healthcare providers, carers, and older people (Slattery et al., 2020). A co-design process could be used generate novel self-care and self-



management instruments designed for application in healthcare settings that incorporate the perspectives and values of older people.

### Practice Implications

This review contributes to decision-making in clinical practice in terms of selecting appropriate instruments to measure self-care and self-management among older adults with chronic conditions and multimorbidity. The characteristics of existing tools (e.g., length, scope, and dimensionality) might inform the selection of a specific tool for use in clinical care. Ideally, the tool should assess the person's individual needs and circumstances, as well as their values and preferences, aligned with recommendations on supporting self-care and self-management. Using dedicated tools in clinical practice would potentially contribute to improved delivery of care tailored to the needs of the person. Although comprehensive, multidimensional tools can enable measurement of different aspects of self-care and self-management, they can increase the response burden, thereby making them unfeasible for use in routine clinical practice. Conversely, instruments containing fewer items would reduce the response time required to administer them in a clinical context. However, they might lack adequate specificity in pinpointing individual support needs, as well as guiding tailored interventions and appropriate referral pathways. For use in routine practice, instruments should aim for brevity and must be feasible to administer yet provide enough specific information to support individualized patient care. A brief screening instrument might prove more practical to implement within busy primary care settings to identify individuals with reduced capacity to self-manage who would benefit from accessing additional resources and supports. Screening is a common practice in a variety of medical settings, such as mental health screening (Lee et al., 2018) and social support screening (Schultz et al., 2022). Screening tools can also be used to identify risk factors such as frailty using simple measures such as assessment of gait speed, timed-up-and-go tests, or the PRISMA-7 questionnaire (Ambagtsheer et al., 2020; NICE, 2016). Health and social care workers could carry out screening of older adults' capacity to self-manage their chronic condition(s) in primary care and community settings to inform an individualized approach to assessment and management. Ideally, a positive result would then be followed up by a longer, more comprehensive assessment in which the person's capacities and care needs are assessed in greater depth and a personalized care plan is created. Such assessments might incorporate existing metrics related to disease-related factors (e.g., disease severity), socioeconomic factors (e.g., social support), multimorbidity (e.g., cognition, co-morbid conditions), and environmental factors (e.g., access to care, built environment) (Jaarsma et al., 2020).

### Conclusion

We conducted this review to map the range of tools that measure self-care and self-management of chronic conditions by community-dwelling older adults and describe their characteristics. Identifying instruments that are suitable for use in research and clinical practice can guide healthcare providers in their selection. Robust and rigorous assessment and monitoring would allow verification of the effectiveness of any efforts undertaken by healthcare providers to support self-care and self-management. They would also support the implementation of person-centered care by refocusing care and support planning on what matters to older people and their carers, identifying unmet needs and areas of strength, and tailoring support accordingly. However, the number of tools, including duplication and variability within literature, highlights gaps between policy discourse and the rigor with which self-care and self-management of chronic conditions are currently conceptualized and measured. The variability in the literature makes it difficult to determine the potential utility of existing tools for measuring self-care and self-management in older people with chronic conditions and multimorbidity. Selecting a tool for use in a specific context depends on the rigor of the tool, its feasibility (e.g., length), and the applicability of the tool to that context. Some tools identified in this review appear suitable for use with older people. However, further research is needed to assess the quality of existing tools to assist researchers, policymakers, and clinicians in assessing self-care and self-management to promote tailored care for older adults with complex care needs.

### Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: RC receives salary support from Australian National Health and Medical Research Council (APP1194051). JH is supported by a Future Leader Fellowship provided by the Australian Heart Foundation.

### ORCID iD

Michael T. Lawless  <https://orcid.org/0000-0002-2536-6442>

### Supplemental Material

Supplemental material for this article is available online.

### References

- Ahmed, S., Heaven, A., Lawton, R., Rawlings, G., Sloan, C., & Clegg, A. (2021). Behaviour change techniques in personalised care planning for older people: A systematic review. *The British*

*Journal of General Practice: The Journal of the Royal College of General Practitioners*, 71(703), e121–e127. <https://doi.org/10.3399/bjgp20X714017>

- Ambagtsheer, R. C., Archibald, M. M., Lawless, M., Kitson, A., & Beilby, J. (2020). Feasibility and acceptability of commonly used screening instruments to identify frailty among community-dwelling older people: A mixed methods study. *BMC Geriatrics*, 20(1), 152–211. <https://doi.org/10.1186/s12877-020-01551-6>.
- American Geriatrics Society Expert Panel on Person-Centered Care, Brummel Smith, K., Butler, D., Frieder, M., Gibbs, N., Henry, M., & Vladeck, B. C. (2016). Person centered care: A definition and essential elements. *Journal of the American Geriatrics Society*, 64(1), 15–18. <https://doi.org/10.1111/jgs.13866>
- Arksey, H., & O'Malley, L. (2005). Scoping studies: Towards a methodological framework. *International Journal of Social Research Methodology*, 8(1), 19–32. <https://doi.org/10.1080/1364557032000119616>
- Artinian, N. T., Magnan, M., Sloan, M., & Lange, M., P. (2002). Self-care behaviors among patients with heart failure. *Heart & Lung*, 31(3), 161–172. <https://doi.org/10.1067/mhl.2002.123672>.
- Audulv, A., Asplund, K., & Norbergh, K. G. (2012). The integration of chronic illness self-management. *Qualitative Health Research*, 22(3), 332–345. <https://doi.org/10.1177/1049732311430497>
- Ausili, D., Barbaranelli, C., Rossi, E., Rebora, P., Fabrizi, D., Coghi, C., Luciani, M., Vellone, E., Di Mauro, S., & Riegel, B. (2017). Development and psychometric testing of a theory-based tool to measure self-care in diabetes patients: The self-care of diabetes inventory. *BMC Endocrine Disorders*, 17(1), 66–72. <https://doi.org/10.1186/s12902-017-0218-y>
- Ausili, D., Masotto, M., Dall'Ora, C., Salvini, L., & Di Mauro, S. (2014). A literature review on self-care of chronic illness: Definition, assessment and related outcomes. *Professioni Infermieristiche*, 67(3), 180–189. <https://doi.org/10.7429/pi.2014.673180>
- Bandura, A. (1997). *Self-efficacy: The exercise of control*. Freeman.
- Barlow, J., Wright, C., Sheasby, J., Turner, A. P., & Hainsworth, J. (2002). Self-management approaches for people with chronic conditions: A review. *Patient Education and Counseling*, 48(2), 177–187. [https://doi.org/10.1016/s0738-3991\(02\)00032-0](https://doi.org/10.1016/s0738-3991(02)00032-0)
- Battersby, M. W., Ask, A., M Reece, M., J Markwick, M., & P Collins, J. (2003). The Partners in Health scale: The development and psychometric properties of a generic assessment scale for chronic condition self-management. *Australian Journal of Primary Health*, 9(3), 41–52. <https://doi.org/10.1071/PY03022>
- Barnett-Page, E., & Thomas, J. (2009). Methods for the synthesis of qualitative research: A critical review. *BMC Medical Research Methodology*, 9(1), 59. <https://doi.org/10.1186/1471-2288-9-59>
- Battersby, M., Harvey, P., Mills, P. D., Kalucy, E., Pols, R. G., Frith, P. A., McDonald, P., Esterman, A., Tsourtos, G., Donato, R., Pearce, R., & McGowan, C. (2007). SA HealthPlus: A controlled trial of a statewide application of a generic model of chronic illness care. *The Milbank Quarterly*, 85(1), 37–67. <https://doi.org/10.1111/j.1468-0009.2007.00476.x>
- Baugh Littlejohns, L., & Wilson, A. (2019). Strengthening complex systems for chronic disease prevention: A systematic review. *BMC Public Health*, 19(1), 729–813. <https://doi.org/10.1186/s12889-019-7021-9>
- Bijl, J. V. D., Poelgeest Eeltink, A. V., & Shortridge Baggett, L. (1999). The psychometric properties of the diabetes management self efficacy scale for patients with type 2 diabetes mellitus. *Journal of Advanced Nursing*, 30(2), 352–359. <https://doi.org/10.1046/j.1365-2648.1999.01077.x>
- Boehmer, K. R., Gionfriddo, M. R., Rodriguez-Gutierrez, R., Dabrh, A. M. A., Leppin, A. L., Hargraves, I., May, C. R., Shippee, N. D., Castaneda-Guarderas, A., Palacios, C. Z., Bora, P., Erwin, P., Montori, V. M., & Montori, V. M. (2016). Patient capacity and constraints in the experience of chronic disease: A qualitative systematic review and thematic synthesis. *BMC Family Practice*, 17(1), 127–223. <https://doi.org/10.1186/s12875-016-0525-9>
- Boehmer, K. R., Kyriacou, M., Behnken, E., Branda, M., & Montori, V. M. (2018). Patient capacity for self-care in the medical record of patients with chronic conditions: A mixed-methods retrospective study. *BMC Family Practice*, 19(1), 164–167. <https://doi.org/10.1186/s12875-018-0852-0>
- Cameron, J., Worrall-Carter, L., Driscoll, A., & Stewart, S. (2009). Measuring self-care in chronic heart failure: A review of the psychometric properties of clinical instruments. *The Journal of Cardiovascular Nursing*, 24(6), E10–E22. <https://doi.org/10.1097/JCN.0b013e3181b5660f>
- Caro-Bautista, J., Martín-Santos, F. J. J., & Morales-Asencio, J. M. (2014). Systematic review of the psychometric properties and theoretical grounding of instruments evaluating self-care in people with type 2 Diabetes Mellitus. *Journal of Advanced Nursing*, 70(6), 1209–1227. <https://doi.org/10.1111/jan.12298>
- Cesari, M., Sumi, Y., Han, Z. A., Perracini, M., Jang, H., Briggs, A., Amuthavalli Thiyagarajan, J., Sadana, R., & Banerjee, A. (2022). Implementing care for healthy ageing. *BMJ Global Health*, 7(2), e007778. <https://doi.org/10.1136/bmjgh-2021-007778>
- Coulter, A., Entwistle, V. A., Eccles, A., Ryan, S., Shepperd, S., & Perera, R. (2015). Personalised care planning for adults with chronic or long term health conditions. *The Cochrane Database of Systematic Reviews*, 2015(3), CD010523. <https://doi.org/10.1002/14651858.CD010523.pub2>
- Cramm, J. M., & Nieboer, A. P. (2012). Factorial validation of the patient assessment of chronic illness care (PACIC) and PACIC short version (PACIC-S) among cardiovascular disease patients in The Netherlands. *Health and Quality of Life Outcomes*, 10(1), 104–107. <https://doi.org/10.1186/1477-7525-10-104>
- Cramm, J. M., Strating, M. M. H., De Vreede, P. L., Steverink, N., & Nieboer, A. P. (2012). Validation of the self-management ability scale (SMAS) and development and validation of a shorter scale (SMAS-S) among older patients shortly after

- hospitalisation. *Health and Quality of Life Outcomes*, 10(1), 9–17. <https://doi.org/10.1186/1477-7525-10-9>
- Curtin, R. B., Walters, B. A. J., Schatell, D., Pennell, P., Wise, M., & Klicko, K. (2008). Self-efficacy and self-management behaviors in patients with chronic kidney disease. *Advances in Chronic Kidney Disease*, 15(2), 191–205. <https://doi.org/10.1053/j.ackd.2008.01.006>
- Daudt, H. M., Van Mossel, C., & Scott, S. J. (2013). Enhancing the scoping study methodology: A large, inter-professional team's experience with Arksey and O'Malley's framework. *BMC Medical Research Methodology*, 13(1), 48–49. <https://doi.org/10.1186/1471-2288-13-48>
- Dickson, V. V., Fletcher, J., & Riegel, B. (2021). Psychometric testing of the self-care of hypertension inventory version 3.0. *The Journal of Cardiovascular Nursing*, 36(5), 411–419. <https://doi.org/10.1097/JCN.0000000000000827>
- Dineen-Griffin, S., Garcia-Cardenas, V., Williams, K., & Benrimoj, S. I. (2019). Helping patients help themselves: A systematic review of self-management support strategies in primary health care practice. *PloS One*, 14(8), e0220116. <https://doi.org/10.1371/journal.pone.0220116>
- Dye, C. J., Haley-Zitlin, V., & Willoughby, D. (2003). Insights from older adults with type 2 diabetes: Making dietary and exercise changes. *The Diabetes Educator*, 29(1), 116–127. <https://doi.org/10.1177/014572170302900116>
- Eghbali-Babadi, M., Feizi, A., Khosravi, A., Nouri, F., Taheri, M., & Sarrafzadegan, N. (2019). Development and evaluation of the psychometric properties of a hypertension self-care questionnaire. *ARYA Atherosclerosis*, 15(5), 241–249. <https://doi.org/10.22122/arya.v15i5.1835>
- Evers, G. C. M., Isenberg, M. A., Philipsen, H., Senten, M., & Brouns, G. (1993). Validity testing of the Dutch translation of the appraisal of the self-care agency ASA-scale. *International Journal of Nursing Studies*, 30(4), 331–342. [https://doi.org/10.1016/0020-7489\(93\)90105-4](https://doi.org/10.1016/0020-7489(93)90105-4)
- Fabbi, E., Zoli, M., Gonzalez-Freire, M., Salive, M. E., Studenski, S. A., & Ferrucci, L. (2015). Aging and multimorbidity: New tasks, priorities, and frontiers for integrated gerontological and clinical research. *Journal of the American Medical Directors Association*, 16(8), 640–647. <https://doi.org/10.1016/j.jamda.2015.03.013>
- Farin, E., Fleitz, A., & Frey, C. (2007). Psychometric properties of an International Classification of Functioning, Disability and Health (ICF)-oriented, adaptive questionnaire for the assessment of mobility, self-care and domestic life. *Journal of Rehabilitation Medicine*, 39(7), 537–546. <https://doi.org/10.2340/16501977-0083>
- Fulmer, T., Reuben, D. B., Auerbach, J., Fick, D. M., Galambos, C., & Johnson, K. S. (2021). Actualizing better health and health care for older adults. *Health Affairs*, 40(2), 219–225. <https://doi.org/10.1377/hlthaff.2020.01470>
- Garnett, A., Ploeg, J., Markle-Reid, M., & Strachan, P. H. (2018). Self-management of multiple chronic conditions by community-dwelling older adults: A concept analysis. *SAGE Open Nursing*, 4(January), 1–16. <https://doi.org/10.1177/2377960817752471>
- Glasgow, R. E., Wagner, E. H., Schaefer, J., Mahoney, L. D., Reid, R. J., & Greene, S. M. (2005a). Development and validation of the patient assessment of chronic illness care (PACIC). *Medical Care*, 43(5), 436–444. <https://doi.org/10.1097/01.mlr.0000160375.47920.8c>
- Glasgow, R. E., Whitesides, H., Nelson, C. C., & King, D. K. (2005b). Use of the Patient Assessment of Chronic Illness Care (PACIC) with diabetic patients: Relationship to patient characteristics, receipt of care, and self-management. *Diabetes Care*, 28(11), 2655–2661. <https://doi.org/10.2337/diacare.28.11.2655>
- Goodman, R. A., Posner, S. F., Huang, E. S., Parekh, A. K., & Koh, H. K. (2013). Defining and measuring chronic conditions: Imperatives for research, policy, program, and practice. *Preventing Chronic Disease*, 10(April), E66. <https://doi.org/10.5888/pcd10.120239>
- Grady, P. A., & Gough, L. L. (2014). Self-management: A comprehensive approach to management of chronic conditions. *American Journal of Public Health*, 104(8), e25–e31. <https://doi.org/10.2105/AJPH.2014.302041>
- Han, H. R., Song, H. J., Nguyen, T., & Kim, M. T. (2014). Measuring self-care in patients with hypertension: A systematic review of literature. *The Journal of Cardiovascular Nursing*, 29(1), 55–67. <https://doi.org/10.1097/JCN.0b013e3182775fd1>
- Hastings, J., Michie, S., & Johnston, M. (2020). Theory and ontology in behavioural science. *Nature Human Behaviour*, 4(3), 226. <https://doi.org/10.1038/s41562-020-0826-9>
- Hibbard, J. H., Mahoney, E. R., Stockard, J., & Tusler, M. (2005). Development and testing of a short form of the patient activation measure. *Health Services Research*, 40(6 Pt 1), 1918–1930. <https://doi.org/10.1111/j.1475-6773.2005.00438.x>
- Hibbard, J. H., Stockard, J., Mahoney, E. R., & Tusler, M. (2004). Development of the Patient Activation Measure (PAM): Conceptualizing and measuring activation in patients and consumers. *Health Services Research*, 39(4 Pt 1), 1005–1026. <https://doi.org/10.1111/j.1475-6773.2004.00269.x>
- Hicks, F. D., & Holm, K. (2003). Self-management decision influences in heart failure: A preliminary investigation. *Clinical Nursing Research*, 12(1), 69–84. <https://doi.org/10.1177/1054773803238741>
- Hudon, E., Hudon, C., Lambert, M., Bisson, M., & Chouinard, M. C. (2021). Generic self-reported questionnaires measuring self-management: A scoping review. *Clinical Nursing Research*, 30(6), 855–865. <https://doi.org/10.1177/1054773820974149>
- Jaarsma, T., Strömberg, A., Dunbar, S. B., Fitzsimons, D., Lee, C., Middleton, S., Vellone, E., Freedland, K. E., & Riegel, B. (2020). Self-care research: How to grow the evidence base? *International Journal of Nursing Studies*, 105, 103555. <https://doi.org/10.1016/j.ijnurstu.2020.103555>
- Jaarsma, T., Stromberg, A., Martensson, J., & Dracup, K. (2003). Development and testing of the European heart failure self care

- behaviour scale. *European Journal of Heart Failure*, 5(3), 363–370. [https://doi.org/10.1016/S1388-9842\(02\)00253-2](https://doi.org/10.1016/S1388-9842(02)00253-2)
- Jack, M. (2007). Ferus – questionnaire for measuring resources and self-management skills. *Hogrefe*.
- Kearney, B. Y., & Fleischer, B. J. (1979). Development of an instrument to measure exercise of self care agency. *Research in Nursing and Health*, 2(1), 25–34. <https://doi.org/10.1002/nur.4770020105>
- Keddy, A. C., Packer, T. L., Auduly, A., Sutherland, L., Sampalli, T., Edwards, L., & Kephart, G. (2021). The Team Assessment of Self-Management Support (TASMS): A new approach to uncovering how teams support people with chronic conditions. *Healthcare Management Forum*, 34(1), 43–48. <https://doi.org/10.1177/0840470420942262>
- Kingston, A., Robinson, L., Booth, H., Knapp, M., & Jagger, C., Modem project (2018). Projections of multi-morbidity in the older population in England to 2035: Estimates from the Population Ageing And Care Simulation (PACSim) model. *Age and Ageing*, 47(3), 374–380. <https://doi.org/10.1093/ageing/afx201>
- Lawless, M. (2022). *Instruments measuring self-care and self-management of chronic conditions by community-dwelling older adults: A scoping review protocol*. Center for Open Science. <https://doi.org/10.17605/OSF.IO/UBJQ5>
- Lawless, M. T., Tieu, M., Feo, R., & Kitson, A. L. (2021). Theories of self-care and self-management of long-term conditions by community-dwelling older adults: A systematic review and meta-ethnography. *Social Science and Medicine*, 287, 114393. <https://doi.org/10.1016/j.socscimed.2021.114393>
- Lee, C. L., Lin, C. C., & Anderson, R. (2016). Psychometric evaluation of the diabetes self-management instrument short form (DSMI-20). *Applied Nursing Research*, 29, 83–88. <https://doi.org/10.1016/j.apnr.2015.04.013>
- Lee, C. S., Westland, H., Faulkner, K. M., Iovino, P., Thompson, J. H., Sexton, J., Farry, E., Jaarsma, T., & Riegel, B. (2022). The effectiveness of self-care interventions in chronic illness: A meta-analysis of randomized controlled trials. *International Journal of Nursing Studies*, 134, 104322. <https://doi.org/10.1016/j.ijnurstu.2022.104322>
- Lee, L., Patel, T., Hillier, L. M., Locklin, J., Milligan, J., Pefanis, J., Costa, A., Lee, J., Slonim, K., Giangregorio, L., Hunter, S., Keller, H., & Boscart, V. (2018). Frailty screening and case-finding for complex chronic conditions in older adults in primary care. *Geriatrics*, 3(3), 39. <https://doi.org/10.3390/geriatrics3030039>
- Levac, D., Colquhoun, H., & O'Brien, K. K. (2010). Scoping studies: Advancing the methodology. *Implementation Science*, 5(1), 69. <https://doi.org/10.1186/1748-5908-5-69>
- Liddy, C., Blazkho, V., & Mill, K. (2014). Challenges of self-management when living with multiple chronic conditions: Systematic review of the qualitative literature. *Canadian family physician Medecin de famille canadien*, 60(12), 1123–1133.
- Lin, C. C., Wu, C. C., Anderson, R. M., Chang, C. S., Chang, S. C., Hwang, S. J., & Chen, H. C. (2012). The chronic kidney disease self-efficacy (CKD-SE) instrument: Development and psychometric evaluation. *Nephrology, Dialysis, Transplantation: Official Publication of the European Dialysis and Transplant Association - European Renal Association*, 27(10), 3828–3834. <https://doi.org/10.1093/ndt/gfr788>
- Lin, C. C., Wu, C. C., Wu, L. M., Chen, H. M., & Chang, S. C. (2013). Psychometric evaluation of a new instrument to measure disease self management of the early-stage chronic kidney disease patients. *Journal of Clinical Nursing*, 22(7–8), 1073–1079. <https://doi.org/10.1111/j.1365-2702.2011.04048.x>
- Lorig, K., Chastain, R. L., Ung, E., Shoor, S., & Holman, H. R. (1989). Development and evaluation of a scale to measure perceived self-efficacy in people with arthritis. *Arthritis and Rheumatism*, 32(1), 37–44. <https://doi.org/10.1002/anr.1780320107>
- Lorig, K., Stewart, A., Ritter, P., Gonzalez, V., Lynch, J., & Laurent, D. (1996). *Outcome measures for health education and other health care interventions*. Sage.
- Lorig, K. R., Ritter, P., Stewart, A. L., Sobel, D. S., Brown, B. W., Bandura, A., Gonzalez, V. M., Laurent, D. D., & Holman, H. R. (2001a). Chronic disease self-management program: 2-year health status and health care utilization outcomes. *Medical Care*, 39(11), 1217–1223. <https://doi.org/10.1097/00005650-200111000-00008>
- Lorig, K. R., Sobel, D. S., Ritter, P. L., Laurent, D., & Hobbs, M. (2001b). Effect of a self-management program on patients with chronic disease. *Effective Clinical Practice: ECP*, 4(6), 256–262.
- Lorig, K. R., Sobel, D. S., Stewart, A. L., Brown, B. W., Jr., Bandura, A., Ritter, P., Gonzalez, V. M., Laurent, D. D., & Holman, H. R. (1999). Evidence suggesting that a chronic disease self-management program can improve health status while reducing hospitalization: A randomized trial. *Medical Care*, 37(1), 5–14. <https://doi.org/10.1097/00005650-199901000-00003>
- Lu, Y., Xu, J., Zhao, W., & Han, H. R. (2016). Measuring self-care in persons with type 2 diabetes: A systematic review. *Evaluation and The Health Professions*, 39(2), 131–184. <https://doi.org/10.1177/0163278715588927>
- Mair, F. S., & May, C. R. (2014). Thinking about the burden of treatment. *BMJ*, 349, g6680. <https://doi.org/10.1136/bmj.g6680>
- Marengoni, A., Angleman, S., Melis, R., Mangialasche, F., Karp, A., Garmen, A., Meinow, B., & Fratiglioni, L. (2011). Aging with multimorbidity: A systematic review of the literature. *Ageing Research Reviews*, 10(4), 430–439. <https://doi.org/10.1016/j.arr.2011.03.003>
- Matarese, M., Clari, M., De Marinis, M. G., Barbaranelli, C., Ivziku, D., Piredda, M., & Riegel, B. (2020). The self-care in chronic obstructive pulmonary disease inventory: Development and psychometric evaluation. *Evaluation and The Health Professions*, 43(1), 50–62. <https://doi.org/10.1177/0163278719856660>
- Matarese, M., Lommi, M., & De Marinis, M. G. (2017). Systematic review of measurement properties of self reported instruments for evaluating self care in adults. *Journal of Advanced Nursing*, 73(6), 1272–1287. <https://doi.org/10.1111/jan.13204>

- Matarese, M., Lommi, M., De Marinis, M. G., & Riegel, B. (2018). A systematic review and integration of concept analyses of self care and related concepts. *Journal of Nursing Scholarship: An Official Publication of Sigma Theta Tau International Honor Society of Nursing*, 50(3), 296–305. <https://doi.org/10.1111/jnu.12385>
- May, C., Montori, V. M., & Mair, F. S. (2009). We need minimally disruptive medicine. *BMJ*, 339(August), b2803. <https://doi.org/10.1136/bmj.b2803>
- Moore, G., Campbell, M., Copeland, L., Craig, P., Movsisyan, A., Hoddinott, P., Littlecott, H., O’Cathain, A., Pfadenhauer, L., Rehfuess, E., Segrott, J., Hawe, P., Kee, F., Couturiaux, D., Hallingberg, B., & Evans, R. (2021). Adapting interventions to new contexts—the ADAPT guidance. *BMJ*, 374, n1679. <https://doi.org/10.1136/bmj.n1679>
- Moore, G. F., Audrey, S., Barker, M., Bond, L., Bonell, C., Hardeman, W., Moore, L., O’Cathain, A., Tinati, T., Wight, D., & Baird, J. (2015). Process evaluation of complex interventions: Medical Research Council guidance. *BMJ*, 350, h1258. <https://doi.org/10.1136/bmj.h1258>
- Munn, Z., Peters, M. D. J., Stern, C., Tufanaru, C., McArthur, A., & Aromataris, E. (2018). Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Medical Research Methodology*, 18(1), 143. <https://doi.org/10.1186/s12874-018-0611-x>
- Muth, C., van den Akker, M., Blom, J. W., Mallen, C. D., Rochon, J., Schellevis, F. G., Becker, A., Beyer, M., Gensichen, J., Kirchner, H., Perera, R., Prados-Torres, A., Scherer, M., Thiem, U., van den Bussche, H., Glasziou, P. P., & Glasziou, P. P. (2014). The Ariadne principles: How to handle multimorbidity in primary care consultations. *BMC Medicine*, 12(1), 223–311. <https://doi.org/10.1186/s12916-014-0223-1>
- National Institutes for Health and Care Excellence. (2016). *Multimorbidity: Clinical assessment and management*. <https://www.nice.org.uk/guidance/ng56>
- Nichols, T., Calder, R., Morgan, M., Lawn, S., Beauchamp, A., Trezona, A., Byambasuren, O., Bowman, J., Duggan, M., Clinton-McHarg, T., Willis, K., Kearns, R., Harris-Roxas, B., Wardle, J., Litt, J., Menzies, D., Dawda, P., Benrimoj, S., Dineen-Griffin, S., & Klepac, B. (2020). *Self-care for health: A national policy blueprint*. Policy paper 2020-01 [Policy report] <https://www.vu.edu.au/sites/default/files/mitchell-institute-self-care-for-health-a-national-policy-blueprint.pdf>
- O’Cathain, A., Croot, L., Sworn, K., Duncan, E., Rousseau, N., Turner, K., Yardley, L., & Hoddinott, P. (2019). Taxonomy of approaches to developing interventions to improve health: A systematic methods overview. *Pilot and Feasibility Studies*, 5(1), 41. <https://doi.org/10.1186/s40814-019-0425-6>
- Ofori-Asenso, R., Chin, K. L., Curtis, A. J., Zomer, E., Zoungas, S., & Liew, D. (2019). Recent patterns of multimorbidity among older adults in high-income countries. *Population Health Management*, 22(2), 127–137. <https://doi.org/10.1089/pop.2018.0069>
- Orem, D. E. (1985). A concept of self-care for the rehabilitation client. *Rehabilitation Nursing: The Official Journal of the Association of Rehabilitation Nurses*, 10(3), 33–36. <https://doi.org/10.1002/j.2048-7940.1985.tb00428.x>
- Osborne, R. H., Elsworth, G. R., & Whitfield, K. (2007). The Health Education Impact Questionnaire (heiQ): An outcomes and evaluation measure for patient education and self-management interventions for people with chronic conditions. *Patient Education and Counseling*, 66(2), 192–201. <https://doi.org/10.1016/j.pec.2006.12.002>
- Packer, T., Kephart, G., Auduly, Å., Keddy, A., Warner, G., Peacock, K., & Sampalli, T. (2020). Protocol for development, calibration and validation of the patient-reported inventory of self-management of chronic conditions (PRISM-CC). *BMJ Open*, 10(9), e036776. <https://doi.org/10.1136/bmjopen-2020-036776>
- Packer, T. L., Fracini, A., Auduly, Å., Alizadeh, N., van Gaal, B. G. I., Warner, G., & Kephart, G. (2018). What we know about the purpose, theoretical foundation, scope and dimensionality of existing self-management measurement tools: A scoping review. *Patient Education and Counseling*, 101(4), 579–595. <https://doi.org/10.1016/j.pec.2017.10.014>
- Palladino, R., Tayu Lee, J., Ashworth, M., Triassi, M., & Millett, C. (2016). Associations between multimorbidity, healthcare utilisation and health status: Evidence from 16 European countries. *Age and Ageing*, 45(3), 431–435. <https://doi.org/10.1093/ageing/afw044>
- Richard, A. A., & Shea, K. (2011). Delineation of self-care and associated concepts. *Journal of Nursing Scholarship: An Official Publication of Sigma Theta Tau International Honor Society of Nursing*, 43(3), 255–264. <https://doi.org/10.1111/j.1547-5069.2011.01404.x>
- Riegel, B., Barbaranelli, C., Sethares, K. A., Daus, M., Moser, D. K., Miller, J. L., Haedtke, C. A., Feinberg, J. L., Lee, S., Stromberg, A., & Jaarsma, T. (2018). Development and initial testing of the self-care of chronic illness inventory. *Journal of Advanced Nursing*, 74(10), 2465–2476. <https://doi.org/10.1111/jan.13775>
- Riegel, B., Carlson, B., Moser, D. K., Sebern, M., Hicks, F. D., & Roland, V. (2004). Psychometric testing of the self-care of heart failure index. *Journal of Cardiac Failure*, 10(4), 350–360. <https://doi.org/10.1016/j.cardfail.2003.12.001>
- Riegel, B., Jaarsma, T., & Strömberg, A. (2012). A middle-range theory of self-care of chronic illness. *Advances in Nursing Science*, 35(3), 194–204. <https://doi.org/10.1097/ANS.0b013e318261b1ba>
- Riegel, B., Lee, C. S., Dickson, V. V., & Carlson, B. (2009). An update on the self-care of heart failure index. *The Journal of Cardiovascular Nursing*, 24(6), 485–497. <https://doi.org/10.1097/JCN.0b013e3181b4baa0>
- Riegel, B., Moser, D. K., Buck, H. G., Dickson, V. V., Dunbar, S. B., Lee, C. S., Lennie, T. A., Lindenfeld, J., Mitchell, J. E., Treat-Jacobson, D. J., & Webber, D. E., American Heart Association Council on Cardiovascular and Stroke Nursing; Council on Peripheral Vascular Disease; and Council on Quality of Care and Outcomes Research and American Heart Association Council on Cardiovascular and Stroke Nursing; Council on

- Peripheral Vascular Disease; and Council on Quality of Care and Outcomes Research. (2017). self care for the prevention and management of cardiovascular disease and stroke: A scientific statement for healthcare professionals from the American heart association. *Journal of the American Heart Association*, 6(9), e006997. <https://doi.org/10.1161/JAHA.117.006997>
- Riegel, B., Westland, H., Iovino, P., Barelds, I., Bruins Slot, J., Stawnychy, M. A., Osokpo, O., Tarbi, E., Trappenburg, J. C. A., Vellone, E., Strömberg, A., & Jaarsma, T. (2021). Characteristics of self-care interventions for patients with a chronic condition: A scoping review. *International Journal of Nursing Studies*, 116(April), 103713. <https://doi.org/10.1016/j.ijnurstu.2020.103713>.
- Ryan, A., Wallace, E., O'Hara, P., & Smith, S. M. (2015). Multimorbidity and functional decline in community-dwelling adults: A systematic review. *Health and Quality of Life Outcomes*, 13(1), 168–213. <https://doi.org/10.1186/s12955-015-0355-9>
- Schmitt, A., Gahr, A., Hermanns, N., Kulzer, B., Huber, J., & Haak, T. (2013). The Diabetes Self-Management Questionnaire (DSMQ): Development and evaluation of an instrument to assess diabetes self-care activities associated with glycaemic control. *Health and Quality of Life Outcomes*, 11(1), 138–214. <https://doi.org/10.1186/1477-7525-11-138>
- Schulman Green, D., Jaser, S. S., Park, C., & Whittmore, R. (2016). A metasynthesis of factors affecting self management of chronic illness. *Journal of Advanced Nursing*, 72(7), 1469–1489. <https://doi.org/10.1111/jan.12902>
- Schultz, B. E., Corbett, C. F., Hughes, R. G., & Bell, N. (2022). Scoping review: Social support impacts hospital readmission rates. *Journal of Clinical Nursing*, 31(19–20), 2691–2705. <https://doi.org/10.1111/jocn.16143>
- Schuermans, H., Steverink, N., Frieswijk, N., Buunk, B. P., Slaets, J. P. J., & Lindenberg, S. (2005). How to measure self-management abilities in older people by self-report. The development of the SMAS-30. *Quality of Life Research: An International Journal of Quality of Life Aspects of Treatment, Care and Rehabilitation*, 14(10), 2215–2228. <https://doi.org/10.1007/s11136-005-8166-9>
- Shepherd, J., Gurney, S., & Patel, H. P. (2022). Shared decision making and personalised care support planning: Pillars of integrated care for older people. *Clinics in Integrated Care*, 12(June), 100097. <https://doi.org/10.1016/j.intcar.2022.100097>.
- Sidani, S. (2011). Self-care. In D. Doran (Ed.), *Nursing outcomes: The state of the science* (2nd ed.). Jones and Bartlett Learning.
- Skou, S. T., Mair, F. S., Fortin, M., Guthrie, B., Nunes, B. P., Miranda, J. J., Boyd, C. M., Pati, S., Mtenga, S., Smith, S. M., & Smith, S. M. (2022). Multimorbidity nature reviews. *Disease Primers*, 8(1), 48. <https://doi.org/10.1038/s41572-022-00376-4>
- Slattery, P., Saeri, A. K., & Bragge, P. (2020). Research co-design in health: A rapid overview of reviews. *Health Research Policy and Systems*, 18(1), 17. <https://doi.org/10.1186/s12961-020-0528-9>
- Smith, S. M., Wallace, E., Clyne, B., Boland, F., & Fortin, M. (2021). Interventions for improving outcomes in patients with multimorbidity in primary care and community setting: A systematic review. *Systematic Reviews*, 10(1), 271–323. <https://doi.org/10.1186/s13643-021-01817-z>
- Sol, B. G. M., Van der Graaf, Y., van der Bijl, J. J., Goessens, N. B. G., & Visseren, F. L. J. (2006). Self-efficacy in patients with clinical manifestations of vascular diseases. *Patient Education and Counseling*, 61(3), 443–448. <https://doi.org/10.1016/j.pec.2005.05.011>
- Tinetti, M. E., & Fried, T. (2004). The end of the disease era. *The American Journal of Medicine*, 116(3), 179–185. <https://doi.org/10.1016/j.amjmed.2003.09.031>
- Toobert, D. J., Hampson, S. E., & Glasgow, R. E. (2000). The summary of diabetes self-care activities measure: Results from 7 studies and a revised scale. *Diabetes Care*, 23(7), 943–950. <https://doi.org/10.2337/diacare.23.7.943>
- Tricco, A. C., Lillie, E., Zarin, W., O'Brien, K. K., Colquhoun, H., Levac, D., Moher, D., Peters, M. D. J., Horsley, T., Weeks, L., Hempel, S., Akl, E. A., Chang, C., McGowan, J., Stewart, L., Hartling, L., Aldcroft, A., Wilson, M. G., Garrity, C., & Straus, S. E. (2018). Prisma extension for scoping reviews (PRISMA-ScR): Checklist and explanation. *Annals of Internal Medicine*, 169(7), 467–473. <https://doi.org/10.7326/M18-0850>
- United Nations. (2019). *World population ageing 2019 [UN Report]*. <https://www.un.org/en/development/desa/population/publications/pdf/ageing/WorldPopulationAgeing2019-Highlights.pdf>
- Vancampfort, D., Koyanagi, A., Ward, P. B., Rosenbaum, S., Schuch, F. B., Mugisha, J., Richards, J., Firth, J., & Stubbs, B. (2017). Chronic physical conditions, multimorbidity and physical activity across 46 low- and middle-income countries. *The International Journal of Behavioral Nutrition and Physical Activity*, 14(1), 6. <https://doi.org/10.1186/s12966-017-0463-5>
- Van de Velde, D., De Zutter, F., Satink, T., Costa, U., Janquart, S., Senn, D., & De Vriendt, P. (2019). Delineating the concept of self-management in chronic conditions: A concept analysis. *BMJ Open*, 9(7), e027775. <https://doi.org/10.1136/bmjopen-2018-027775>
- Vaughan Dickson, V., Lee, C. S., Yehle, K. S., Mola, A., Faulkner, K. M., & Riegel, B. (2017). Psychometric testing of the self care of coronary heart disease inventory (SC-CHDI). *Research in Nursing and Health*, 40(1), 15–22. <https://doi.org/10.1002/nur.21755>
- Vetrano, D. L., Palmer, K., Marengoni, A., Marzetti, E., Lattanzio, F., Roller-Wirnsberger, R., Lopez Samaniego, L., Rodriguez-Manas, L., Bernabei, R., & Onder, G., Joint Action Advantage WP4 Group. (2019). Frailty and multimorbidity: A systematic review and meta-analysis. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 74(5), 659–666. <https://doi.org/10.1093/gerona/gly110>
- Vogeli, C., Shields, A. E., Lee, T. A., Gibson, T. B., Marder, W. D., Weiss, K. B., & Blumenthal, D. (2007). Multiple chronic conditions: Prevalence, health consequences, and implications for quality, care management, and costs. *Journal of General*

- Internal Medicine*, 22(Suppl 3), 391–395. <https://doi.org/10.1007/s11606-007-0322-1>
- Wagner, E. H., Austin, B. T., & Korff, M. V. (1996). Organizing care for patients with chronic illness. *The Milbank Quarterly*, 74(4), 511–544. <https://doi.org/10.2307/3350391>
- Wallace, E., Salisbury, C., Guthrie, B., Lewis, C., Fahey, T., & Smith, S. M. (2015). Managing patients with multimorbidity in primary care. *BMJ*, 350, h176. <https://doi.org/10.1136/bmj.h176>
- Wigal, J. K., Creer, T. L., & Kotses, H. (1991). The COPD self-efficacy scale. *Chest*, 99(5), 1193–1196. <https://doi.org/10.1378/chest.99.5.1193>
- Williams, G. C., McGregor, H. A., Zeldman, A., Freedman, Z. R., & Deci, E. L. (2004). Testing a self-determination theory process model for promoting glycemic control through diabetes self-management. *Health Psychology: Official Journal of the Division of Health Psychology, American Psychological Association*, 23(1), 58–66. <https://doi.org/10.1037/0278-6133.23.1.58>
- World Health Organization. (2015). *World report on ageing and health*. [WHO report] <https://apps.who.int/iris/handle/10665/186463>
- World Health Organization. (2022a). *Ageing and health*. <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>
- World Health Organization. (2022b). *WHO guideline on self-care interventions for health and well-being revision [WHO report]*. <https://www.who.int/publications/i/item/9789240052192>
- Yarnall, A. J., Sayer, A. A., Clegg, A., Rockwood, K., Parker, S., & Hindle, J. V. (2017). New horizons in multimorbidity in older adults. *Age and Ageing*, 46(6), 882–888. <https://doi.org/10.1093/ageing/afx150>
- Yu, D. S. F., De Maria, M., Barbaranelli, C., Vellone, E., Matarese, M., Ausili, D., Rejane, R. S. E., Osokpo, O. H., & Riegel, B. (2021). Cross-cultural applicability of the self-care self-efficacy scale in a multi-national study. *Journal of Advanced Nursing*, 77(2), 681–692. <https://doi.org/10.1111/jan.14617>