



Fungal phylogeny and plant functional traits structure plant–rhizosphere fungi networks in a subtropical forest

Zhu, Chuchao; Wang, Zihui; Luo, Wenqi; Feng, Jiayi; Chen, Yongfa; He, Dong; Ellwood, M. D. Farnon; Chu, Chengjin; Li, Yuanzhi

Oikos

DOI:

<https://doi.org/10.1111/oik.08992>

Published: 01/08/2022

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Zhu, C., Wang, Z., Luo, W., Feng, J., Chen, Y., He, D., Ellwood, M. D. F., Chu, C., & Li, Y. (2022). Fungal phylogeny and plant functional traits structure plant–rhizosphere fungi networks in a subtropical forest. *Oikos*, 2022(8), e08992. <https://doi.org/10.1111/oik.08992>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Fungal phylogeny and plant functional traits structure plant- rhizosphere fungi networks in a**
2 **subtropical forest**

3

4 Chuchao Zhu ^{1,2}, Zihui Wang ², Wenqi Luo ², Jiayi Feng ², Yongfa Chen ², Dong He ², M. D.

5 Farnon Ellwood ³, Chengjin Chu ² and Yuanzhi Li ²

6

7

8 ¹ *Department of Bioengineering, Zhuhai campus of Zunyi Medical University, Zhuhai, 519041,*

9 *China*

10 ² *State Key Laboratory of Biocontrol, School of Ecology and School of Life Sciences, Sun Yat-sen*

11 *University, 510275 Guangzhou, China*

12 ³ *School of Natural Sciences, Bangor University, Bangor, Gwynedd, LL57 2DG, United Kingdom*

13

14

15 *Correspondence*

16 *Yuanzhi Li*

17 *Email: liyuanzhi@mail.sysu.edu.cn, Tel: (+86) 2084111541*

18

19

20 **Abstract**

21 Although rhizosphere fungi are essential for plant survival and ecosystem functioning, little is
22 known about the processes that structure plant–fungal association networks. In this study, we
23 constructed association networks between 43 plant species and two groups of root-associated fungi
24 (mycorrhizal and pathogenic fungi; MF and PF, respectively) in a diverse subtropical forest. We
25 then evaluated the modularity of plant–MF and plant–PF networks and linked them to the
26 functional traits and phylogenies of both plants and fungi. We observed strong modularity in both
27 plant–MF and plant–PF networks. Phylogenetically related fungi tended to emerge in the same
28 modules. MF from distinct modules associated with plants with different specific root length and
29 specific root area in plant–MF networks. PF from distinct modules associated with plants with
30 different dark respiration rate and light compensation point in plant–PF networks. Plant affiliation
31 to modules was explained by both plant traits and phylogeny (22% for plant–MF and 37% for
32 plant–PF networks). In contrast, fungal affiliation to modules was explained by fungal phylogeny
33 (16% for plant–MF and 29% for plant–PF networks). Our results elucidate the link between
34 modularity in plant–root fungal networks and the functional traits and phylogeny of the plants and
35 fungi. Our study highlights the importance of traits and phylogeny in governing root fungal
36 community assembly from network perspective.

37 **Keywords:** network modules, mycorrhizal fungi, root microorganisms, pathogenic fungi,
38 photosynthetic and root traits, network assembly

39

40 **Introduction**

41 Plants and fungi can associate in mutualistic and antagonistic ways, both of which are important for
42 community assembly and ecosystem functioning (Connell 1971, Bennett and Klironomos 2018,
43 2019, Chen et al. 2019). Pathogenic fungi (PF) help maintain species diversity by reducing the
44 recruitment and survival of dominant species (Bagchi et al., 2014; Chen et al., 2019), whereas
45 mutualistic ectomycorrhizal fungi (EM) and arbuscular mycorrhizal fungi (AM) can maintain
46 species diversity through either positive or negative plant–soil feedbacks (Bennett et al. 2017, Toju
47 et al. 2018). In turn, plants may also influence the composition and diversity of soil fungal
48 communities (Dassen et al. 2017, Wang et al. 2019, Schmid et al. 2021)—for example, by changing
49 carbon availability and/or using physical or chemical defences against different types of fungal
50 symbionts (Högberg and Högberg 2002). Therefore, elucidating the structure and influencing
51 factors of plant–mycorrhizal fungal (MF) and plant–pathogenic fungal (PF) networks may provide
52 more insights into the mechanisms governing species coexistence and community assembly.

53 Network modularity is an important topological feature of networks that describes the
54 organisation of a network into different groups (modules), where species within groups associate
55 more intensely than species between groups (Olesen et al. 2007). A modular structure may reflect
56 the existence of strong preferences or selective processes that shape trophic associations between
57 fungi and their hosts (Chagnon et al. 2018). Modularity maintains species diversity and community
58 stability by confining the cascading effects of species extinction or environmental perturbation
59 within a module and preventing ripple effects from spreading to other modules (Olesen et al. 2007).
60 Investigating the modular structure can thus reveal the underlying non-random ecological processes

61 of community assembly (e.g., local adaptation and resource competition) (Valverde et al. 2020) and
62 reveal the topological vulnerability of ecological networks to disturbances (e.g., simulated species
63 extinction cascades) (Montoya et al. 2006).

64 Network modularity is often reported in macro-organisms, including food webs (Montoya et
65 al. 2015) and mutualistic pollinator and seed-dispersal networks (Donatti et al. 2011, Morente-
66 López et al. 2018). However, there is no consensus regarding the modular structure of belowground
67 plant–root fungal networks. Mixed evidence for modularity (significantly and non-significantly
68 strong modularity compared to 1000 randomised networks) was reported in mutualistic plant–MF
69 networks (Montesinos-Navarro et al. 2012, Bahram et al. 2014, Toju et al. 2014). This inconsistency
70 may be due to low sampling efforts for root samples, resulting in a limited number of root fungal
71 groups (Bahram et al. 2014). Nevertheless, strong modularity is frequently reported in antagonistic
72 tree-parasitic fungus and plant–pathogen networks (considering root PF as one component) (Vacher
73 et al. 2008, Bufford et al. 2020). However, no previous studies have independently illustrated a
74 modular structure in plant–root PF networks. Therefore, sufficient samples from diverse
75 communities are necessary to explore modularity in plant–MF and plant–root PF networks.

76 To reveal the mechanisms underlying a modular structure, a growing number of studies
77 account for module division and species composition in a module (Donatti et al. 2011, Torrecillas et
78 al. 2014, Robinson et al. 2015). For instance, the modules of aboveground mutualistic pollinator
79 and seed-dispersal networks have been reported to show convergence in functional traits and toward
80 syndromes, respectively (Donatti et al. 2011, Robinson et al. 2015). However, to our knowledge,
81 only three plant traits (plant specific root length, specific leaf area, and leaf dry matter content) have

82 been used to explore module assembly in plant–MF networks (Chagnon et al. 2015). Of these traits,
83 only leaf dry matter content (representing a plant’s investment in leaf structural tissues) has been
84 reported to show convergence in modules and significantly affect the association of AM host plants
85 with modules (Chagnon et al. 2015). This finding highlights the importance of plant ecological
86 strategies in module composition and assembly (Torrecillas et al. 2014).

87 Ecological strategies are represented by the traits of both plant and fungal components, and are
88 important in explaining fungal community assembly. Plant traits—including root morphology
89 (related to plant nutrient acquisition efficiency) and the photosynthesis and respiration rates of
90 leaves (related to carbohydrate accumulation)—are key predictors of root fungal communities
91 (Koorem et al. 2017, Sepp et al. 2019, Davison et al. 2020). For instance, plants with high specific
92 root area tend to be colonised by MF, whereas those with low specific root area tend to be colonised
93 by PF (Wang et al. 2019). Fungal traits—such as fruit body size (Abrego et al. 2017) and hyphal
94 exploration type (Olchowik et al. 2021)—are also critical in determining the colonisation success
95 and community assemblages of root fungal communities. Despite the importance of plant and
96 fungal traits in community assembly, there is a lack of evidence about their effects on module
97 assembly. Incorporating plant and fungal traits to explain module composition can provide more
98 insights into the mechanism of assembly of plant–root fungal networks.

99 As a proxy for their respective functional traits, plant and fungal phylogenies have also been
100 used to explore the mechanisms of non-random organisation of network modularity (Chagnon et al.
101 2013). Previous studies have shown that modules are clustered by host phylogeny in plant–PF
102 networks (Bufford et al. 2020), and by fungal phylogeny in plant–AM networks (Chagnon et al.

103 2015). Whether modularity in the form of fungal phylogenetic clustering is detectable in root PF
104 networks remains unknown. This is important to elucidate, as it is possible that when competing for
105 plant root resources, root PF and mutualistic MF exhibit different patterns of phylogenetic
106 conservatism in host use.

107 To better understand the structure and assembly of plant–root fungal networks, it is essential to
108 simultaneously explore plant–MF and plant–PF networks in diverse communities and link them to
109 the phylogeny and functional traits of both host plants and root fungi. In this study, we compiled
110 two datasets of plant–MF and plant–PF networks comprising of 43 plant species, 883 MF, and 113
111 PF (Wang et al. 2019). We measured 17 functional traits (including root and photosynthesis traits)
112 of the 43 host plant species in a 50-ha plot in a subtropical forest. After synthesising these data, we
113 tested the following hypotheses: (i) there is strong modularity in the plant–MF and plant–PF
114 networks in a highly diverse subtropical forest community; (ii) as in plant–seed dispersal and plant–
115 pollinator networks, module composition is expected to be constrained by functional traits and
116 phylogeny of plants and fungi; and (iii) the relative importance of functional traits and phylogeny
117 may be different in predicting module composition in these networks.

118 **Materials and methods**

119 **Study site**

120 This study was conducted in a 50-ha plot in a subtropical forest in Heishiding Nature Reserve,
121 Southern China (23°25′–23°29′ N, 111°49′–111°55′ E). The mean annual temperature of this area
122 is 19.7 °C, and the annual precipitation is 1,750 mm. The total area of this nature reserve is 4,200

123 ha, including a 2,202 ha core area and a 1,660 ha experimental area. We established the 50-ha forest
124 plot in 2012, and identified all trees with a diameter at breast height (DBH) >1 cm. In total, this plot
125 included approximately 2,69,000 stems of 213 woody plant species (Wang et al. 2019).

126 **Root sampling and molecular identification**

127 We compiled the dataset of fungal communities from 512 root samples of the 43 plant species (no
128 less than 5 sampled individuals for each plant species) in the Heishiding plot. These plant species
129 were selected based on their taxonomic placement and abundance. Specifically, we selected three
130 *Litsea* spp. and three *Lithocarpus* spp. (the two most abundant genera), along with species from
131 other genera (including some genera that were in the same families as *Litsea* and *Lithocarpus*). For
132 each plant species, we randomly selected 5–15 individuals for fine root sampling. The unequal
133 sampling efforts for each species tree resulted from: (i) the rarity of some tree species that limited
134 the collection of sufficient root samples and (ii) failure in extracting high-quality DNA. At least
135 three root fragments (each approximately 2 cm in length) around an individual tree were traced
136 from different directions and then pooled to create a single sample. The fine root samples were
137 immediately cooled on ice in the field and stored at -20 °C in a refrigerator until processing. More
138 sampling details can be found in a previously published paper (Wang et al. 2019).

139 Of 100 fine-root samples (randomly selected from all root samples), the tree species of 97 root
140 samples traced in the field were correctly confirmed by *rbcLa* sequences obtained from a Sanger
141 sequencing platform. Thus, we considered the tracing method as an accurate strategy to capture the
142 taxonomic (species level) information of sampled fine roots. Root-associated fungi were identified
143 by the internal transcribed spacer (ITS) region of fungal rDNA. After removing chimeric sequences,

144 we obtained 11,000,000 high-quality reads of the ITS region of fungal rDNA. The operational
145 taxonomic units (OTUs) of root fungi were discriminated using a threshold of 97% sequence
146 identity. Each sequence was assigned to a taxonomic label based on the UNITE database using the
147 Ribosomal Database Project (RDP) classifier (Wang, Garrity, Tiedje, & Cole, 2007). Each fungal
148 genus was then assigned into functional categories. We identified EM fungi by blasting our fungal
149 genera against the fungal genera in a database of EM taxa and lineages (Tedersoo and Smith 2013).
150 We assigned all OTUs in Glomeromycota to AM fungi (Schüßler 2002). Because we could only
151 identify 21 OTUs of AM, EM and AM were pooled to represent the MF guild. Identifying fungal
152 plant pathogens is challenging, because identification can only take place after the plants are
153 diseased. Therefore, pathogenic genera were initially identified using the FUNGuild database
154 (Nguyen et al. 2016). We then consulted the literature and retained only potential pathogens (OTUs)
155 that had been identified to the species level and are known to be pathogenic to woody plants.

156 To evaluate network modularity, we constructed a plant–PF association network including 113
157 fungal plant pathogens and 43 plant species, as well as a plant–MF association network including 883
158 mycorrhizal fungi (862 EM and 21 AM) and 43 plant species. To account for the sampling inequality,
159 each cell in each network matrix was filled with the mean abundance (sequenced reads) of each fungal
160 OTU (species) on each sampled tree, and the numbers were rounded to the nearest integer. Abundance
161 of fungal OTUs on each sampled tree was calculated after subsampling each sample to 3000 sequence
162 reads to eliminate the effects of sample size (Wang et al. 2019).

163 **Functional traits**

164 We recorded and analysed the data of 17 functional traits of the 43 host plants, including 2 leaf

165 morphological traits, 3 leaf chemical traits, 8 photosynthetic traits, and 4 root traits (Table S1).
166 Details about how these traits were sampled and measured can be found in previous studies (Feng et
167 al. 2018, He et al. 2018, Wang et al. 2019, Luo et al. 2020). The photosynthetic traits of 7 plant
168 species and the root traits of 11 plant species were unavailable, and these species were removed
169 from relevant analyses. We collected the data of 4 functional traits of fungi (growth form, fruitbody
170 type, hymenium type, and hyphal exploration type) from the FungalTraits database (Pöhlme et al.
171 2020). There was no variation in the four fungal traits for AM fungi (21 OTUs). For the other fungi
172 (PF, MF, and EM), the fungal traits available for statistical analysis are listed in Table S2.

173 **Statistical analysis**

174 ***Reconstructing phylogeny of plant and fungal species***

175 We used four plant DNA barcodes (*rbcLa*, *matK*, *trnL*, and ITS2) to reconstruct the phylogenetic
176 relationships between all local plant species using the RAxML software (Stamatakis 2014). The
177 sequences were aligned in the Clustal Omega software using the default Gonnet transition matrix,
178 with 6 bits of gap opening penalty and 1 bit of gap extension penalty. The best maximum likelihood
179 phylogeny for plants (Fig. S1) was inferred using the GTR + GAMMA evolutionary model and
180 1000 fast bootstrap replicates in the RAxML software (Stamatakis 2014).

181 We used the taxonomic rank information of local root MF and PF (996 OTUs) and one
182 outgroup (*Olpidium brassicae*) to reconstruct the phylogeny of root-associated fungi using a Perl
183 script (`taxonomy_to_tree.pl` script of Tedersoo et al. 2018). By incorporating a fungal taxonomic
184 backbone, we converted the hierarchical classification of our focal fungal taxa to a Newick-
185 formatted phylogeny (Fig. S2). In the reconstructed fungal phylogeny (Fig. S2), we assigned a

186 branch length of 60 between each of the taxonomic ranks (e.g. species, genus, family).

187 ***Detecting the modular structure of plant-root fungal association networks***

188 We first estimated modularity (M)—ranging from 0 (low modularity) to 1 (high modularity)—in
189 abundance-weighted plant–MF and plant–PF networks. A LPAwb+ algorithm was used to search the
190 module divisions that maximised weighted modularity (Beckett 2016), such that maximal
191 modularity was obtained when no better division into modules could be detected (Beckett 2016). To
192 stabilise this computation, we re-ran the algorithm 50 times and reported the most modular result
193 (i.e., the maximum value of modularity and module divisions). As modularity vary with network
194 size and connectivity, we used the z-score of network modularity $z = (M - \bar{M}_{random})/\bar{M}_{random}$
195 (Olesen et al. 2007) to quantify the degree of network modularity among networks. We generated
196 1000 randomised null networks with constant marginal totals and connectance using the “swap”
197 method (Artzy-randrup and Stone 2005, Dormann et al. 2009). From these randomised networks,
198 we calculated \bar{M}_{random} as the average modularity. The above calculations were conducted in the R
199 package *bipartite* (Dormann et al. 2009).

200 ***Detecting the constraints of phylogeny and functional traits on network modularity***

201 To understand the relationship between phylogeny and network modularity, we evaluated the
202 correlation between plant (and fungus) co-occurrence in modules and their phylogenetic proximity.
203 To evaluate the significance of the correlation, we compared the correlation coefficient calculated
204 from the matrix of observed species-module data to those calculated from 999 randomised matrices.
205 This comparison was performed using the “comm.phylo.cor” function in the R package *picante*

206 (Kembel et al. 2010). The co-occurrence of a pair of species within modules was measured using
207 Schoener's index of co-occurrence (Hardy 2008). To test whether some plant traits (Table S1)
208 differed significantly between modules in the observed network, we conducted one-way Type II
209 ANOVA and kruskal–Wallis rank sum test. Some trait variables were log-transformed to meet the
210 assumption of homogeneity of variance and normality of error distribution (Table S3). To detect the
211 impact of fungal traits on the fungal composition of modules, we performed Fisher's exact tests for
212 a series of matrices (module-fungal trait) containing the richness or abundance data of fungal
213 species belonging to each trait category in modules. Five fungal traits (growth form, fruit body type,
214 and hymenium type, hyphal exploration type and mycorrhiza type) were used to explain the fungal
215 composition of modules. Although the morphological traits for some EM fungal groups (e.g.,
216 *Cenococcum*) were incomplete in the fungal trait database (Põlme et al. 2020), we assumed that
217 these incomplete traits are sufficient for explaining network assembly. All these analyses were
218 conducted in R version 3.5.1 (R Core Team 2015).

219 ***Analysing the drivers of module composition***

220 To explain the module membership of fungi and their host species from their traits and phylogeny,
221 we used random forest models implemented in the R package *randomForest* (Liaw and Wiener
222 2002). To explain the module membership of the plant host species, we constructed a full random
223 forest model with 17 plant traits and the first 10 eigenvectors of plant phylogeny (Fig. S1) as
224 explanatory variables. To explain the module membership of root MF/PF/EM/AM fungi, we used 5
225 fungal traits (fruit body type, hymenium type, growth form, hyphal exploration type, and
226 mycorrhiza type, or the available trait combination for each group of fungi; see Table S2) and the

227 first 10 eigenvectors of the MF/PF/EM/AM phylogeny as explanatory variables (see Fig. S2).
228 Phylogenetic eigenvectors were calculated from principle coordinate analysis (PCoA) of the
229 phylogenetic distance matrix of plants/fungi (Diniz-Filho et al. 1998). Finally, we used reduced
230 random forest models to explain the modular memberships for plants and fungi, with selected
231 explanatory variables based on the smallest out-of-bag error (Evans and Murphy 2019). Model
232 accuracy was defined as the amount of variation explained by each reduced random forest model in
233 allocating plants and fungi to the observed modules. Finally, we partitioned the variance explained
234 by the reduced random forest models into unique and shared components of traits and phylogeny.

235 **Results**

236 **Structural properties of plant–MF and plant–PF association networks**

237 We observed significantly higher modularity (relative to those in randomised networks) in the
238 observed plant–PF ($z = 7.82$, $M = 0.431$, confidence interval [CI] of null models [0.043, 0.062]) and
239 plant–MF ($z = 6.95$, $M = 0.487$, CI of null models [0.056, 0.067]) association networks. The plant–
240 MF association network was divided into 16 distinct modules (Fig. 1a), with EM fungi in all
241 modules and AM fungi in 9 modules (Fig. 1a). The plant–PF association network was divided into 9
242 distinct modules (Fig. 1b).

243 **The constraints of phylogenetic history and functional traits on modularity**

244 Excluding AM fungi, closely related fungi in other functional groups were more likely to co-occur
245 in modules of the plant–root fungal association networks (Table 1). However, phylogenetic
246 relatedness between plant species did not significantly affect plant distribution across modules

247 (Table 1). In plant-MF network (Table S4), the mean nearest taxon distances (MNTD) were
248 significantly lower than null expectation in 3 modules of MF (modules 7, 9 and 16) and in 2
249 modules of plants (modules 9 and 11). In plant-PF network (Table S5), MNTD were significantly
250 lower than null expectation in only one module of PF (module 4) and plants (module 1),
251 respectively.

252 Certain functional traits of host plants varied between modules in the plant–PF and plant–MF
253 association networks (Fig. 2 and Table S3). Specifically, leaf dry matter content, root tissue density,
254 specific root length (SRL), and specific root area (SRA) differed significantly between modules in
255 the plant–MF association network (Fig. 2 and Table S3). Moreover, leaf dark respiration rate (Rd)
256 and light compensation point (LCP) differed significantly between modules in the plant–PF
257 association network (Fig. 2 and Table S3). The richness and abundance of MF across modules were
258 significantly affected by fruit body type, hymenium type, hyphal exploration type, and mycorrhizal
259 type (Table S2). The abundance (but not richness) of PF across modules was significantly affected
260 by growth form, fruit body type, and hymenium type (Table S2).

261 **Drivers of module composition**

262 In assigning fungal host plants to modules in the two observed networks, the reduced random forest
263 models provided correct prediction rates of 22–38% (Fig. 3a, c, e, g). Compared to plant
264 phylogenies, plant traits were more accurate in predicting the module memberships of plants hosts
265 for MF and AM and EM (Fig. 3a, e, g). Compared to the independent effects of plant traits and
266 phylogeny, their interactions were more accurate in predicting the module memberships of PF host
267 plants (Fig. 3c). In assigning root fungi to network modules, only fungal phylogeny was retained

268 (and fungal traits were removed) during model selection. As a result, the reduced models had
269 accuracy rates of 16–38% in assigning root fungi to the observed modules (Fig. 3b, d, f, h).

270 **Discussion**

271 **Non-random modularity in plant–root fungal association networks**

272 We found that plant–MF and plant–PF networks in a subtropical forest tended to exhibit significant
273 modularity (Fig. 1). This is similar to the patterns observed in some plant–fungus association
274 networks in harsh alpine and subalpine habitats (Toju et al. 2016) and semi-natural grasslands (Sepp
275 et al. 2019). In the forest plot, plant–fungus networks may show a modular structure due to the
276 similar host habitat preferences and spatial distribution among phylogenetically closely related
277 fungi. The fungal composition of each module of the plant–MF network was dominated by EM
278 instead of AM (Fig. 1a and Table S6). This implies that the strong modular structure of the plant–
279 MF fungal network may be primarily determined by relatively higher host specialization in EM than
280 AM (Van Der Heijden et al. 2015). Similarly, stronger host specificity of antagonistic PF (Wang et
281 al. 2019) may result in modularity in the plant–PF network. Moreover, network size (Olesen et al.
282 2007), asymmetry in species number (Pöhlme et al. 2018), and connectivity (Thébault and Fontaine
283 2010) may also affect the detection of modularity. Therefore, further studies focusing on the
284 ecological and evolutionary factors shaping network patterns are warranted to better predict the
285 modular pattern of antagonistic and mutualistic fungal networks.

286 **Phylogenetic clustering of pathogenic and mycorrhizal fungi within modules**

287 We found that fungal species that were phylogenetically closely related tended to emerge in the

288 same modules of plant–root fungal networks (plant–MF and plant–PF networks; Tables 1, S4, S5),
289 which is consistent with the findings of a previous study (Chagnon et al. 2015). These results
290 suggest that the modular organisation of plant and mycorrhizal fungal networks generally reflects
291 the main split in the fungal phylogeny. Phylogenetically related plants were also found in the
292 networks of plants with antagonistic fungi (including leaf and root decay fungi and parasitic fungi)
293 (Vacher et al. 2008, Bufford et al. 2020). Consistent with a previous study (Chagnon et al. 2013), a
294 lack of phylogenetic relatedness among plant hosts associating with root fungi in the modules
295 (Table 1) may be ascribed to low plant species richness in each module and insufficient sampling
296 efforts. We found a significant effect of fungal phylogeny, but not of host phylogeny, on plant–root
297 fungal networks (Table 1), which is not in agreement with the findings of previous studies (Davison
298 et al. 2020). This inconsistency in the effects of evolutionary history may be due to differences in
299 the functional groups of fungi and environmental conditions. Due to the relatively low number of
300 plant species that associated with fungal symbionts, our results may also be constrained by
301 statistical power. Taken together, our results suggest that the evolutionary history of plants and fungi
302 only partially explains the modular patterns observed in our plant–fungus networks.

303 **Module-level trait convergence in plant–MF and plant–PF association networks**

304 To resist attacks from root pathogens and improve the acquisition efficiency of water and nutrients
305 from mycorrhizal fungi, host plants need to allocate vast amounts of material and energy resources
306 to root fungi. Thus, the resource acquisition and allocation strategies of host plants likely regulate
307 fungal selectivity (Sachs et al. 2004) and affect the shaping and evolution of plant–fungal
308 symbioses. Here, we show that plant and fungal traits varied across the modules in plant–fungal

309 networks, potentially driving the plant-fungal associations. The specific plant traits that structure
310 network modules differed between plant–MF and plant–PF networks (Fig. 2 and Table S3).

311 In the plant–MF network in our subtropical forest, MF fungi from distinct modules varied in
312 fungal exploration traits (Table S2) and tended to associate with plants with different root traits
313 (Figs. 2, S3 and S4, Table S3). For example, *Scleroderma* spp. fungi (EM) with long-distance
314 exploration dominated in module 8 (Table S6), and *Russula* spp. fungi (EM) with contact hyphal
315 exploration dominated in module 11 (Table S6). This suggests that module membership in these
316 fungi may be driven by the high and low efficiency (respectively) of nutrient acquisition by the fine
317 roots of host plants (indicated by SRL, SRA, and specific root tips; Figs. 2 and S3, Table S3). Thus,
318 fine roots with low (or high) investment in foraging tend to shape modules containing EM fungi
319 with low (or high) investment in foraging. This is opposite to the general expectation of
320 complementary relationship between root and mycorrhizal fungi but supporting a matching strategy
321 where longer and thinner roots associated with more mycorrhizal hyphae biomass (Chen et al.,
322 2016). These results support the notion that the module assembly of plant–MF association networks
323 can mainly be ascribed to ecological strategies based on plant (Chagnon et al. 2015) and fungal
324 traits.

325 Similarly, PF from distinct modules differed in functional traits (hymenium type, growth form,
326 and fruitbody type, Table S2) and tended to associate with plants with differing photosynthetic traits
327 (Rd and LCP; see Fig. 2 and Table S3). The two most abundant genera of PF (*Cylindrocarpon* and
328 *Mycoleptodiscus*) dominated in module 4 (Table S7), which may be driven by their preference for
329 host plants with high shade-tolerance (lowest LCP and Rd; Fig. 2c, d). The other two PF genera

330 (*Colletotrichum* and *Pestalotiopsis*) were dominant in module 1 (Table S7), which may be due to
331 selection of light-dependence plants (highest LCP and Rd; Fig. 2c, d). These results suggest that
332 plant photosynthesis traits, important indicators of plant allocation on growth and defense, can
333 dictate the host selectivity on PF (García-Guzmán and Heil 2014) , while root traits, as the basis of
334 plant acquisition for water and nutrients, affect the structure of network modules for MF.

335 **The contribution of functional traits and phylogeny to a modular structure**

336 Our results show that plant traits were more important than plant phylogeny in predicting module
337 composition of MF host plants while plant phylogeny was more important for PF host plants (Fig.
338 3). This implies that PF were sharing more coevolutionary history with plant host than MF, while
339 MF were more likely selected by the ecological traits of plants. The low accuracy of fungal traits in
340 predicting module fungal composition (Fig. 3) may be due to the low variation in the fungal traits
341 database (e.g., fungal morphology did not vary within a genus) (Pölmé et al. 2020). Other
342 processes—such as environmental filtering (e.g., soil types and soil properties) (Torrecillas et al.
343 2014, Arraiano-Castilho et al. 2020) and stochastic effects—may jointly drive plant–fungal network
344 assembly by affecting the co-occurrence of plants and fungi. However, the accuracy of predicting
345 module composition was not substantially improved by incorporating functional traits, phylogeny,
346 and soil environmental variables in the model (Fig. S5). Soil environmental variables (obtained
347 from 625 soil cores sampled randomly in a forest plot) (Luo et al. 2021) as well as plant functional
348 traits (at the species level) may reduce the accuracy in predicting module composition (Fig. S5).
349 This suggests that functional traits at the individual level (including intraspecific trait variability)
350 and soil environmental variables around each plant stem should also be used to investigate modular

351 assembly. Incorporating these data may also shed light on the dynamics of root–fungal networks
352 along environmental gradients.

353 **Conclusions**

354 Analysis of plant–root fungal symbiont network is crucial in finding ecological variables, which
355 regulates host selectivity of fungi and organizes root fungal symbiont community structures. We
356 illustrate that plant nutrient acquisition efficiency indicated by root traits may regulate host
357 selectivity of MF, while plant carbohydrate accumulation and shade tolerance indicated by leaf
358 photosynthesis traits mainly regulate host selectivity of PF. Such host selectivity may be also
359 constrained by phylogenetically conservative fungal traits. These results suggest that trait and
360 phylogeny based host selectivity enhance to shape a series of network modules (groups). Overall,
361 our study contributes to elucidating the ecological and evolutionary factors shaping modular
362 structure of plant–fungal association network and reveal the mechanisms governing species
363 coexistence and community assembly.

364 **Acknowledgement** – We thank Wen Zhou for providing DNA barcoding sequences for plants.

365 **Funding** – This work was supported by the National Natural Science Foundation of China
366 (32101281 to CZ, 31925027 to CC and 31901106 to YL) and the Basic Research Project of
367 Guizhou Provincial Science and Technology Department Fund of China (grant no. QKHJC
368 ZK[2021] 096).

369 **Data availability statement**

370 Data are available from the Dryad Digital Repository: <<https://doi.org/10.5061/dryad.kwh70rz5z>>
371 (Zhu et al. 2022).

372 **References**

- 373 Abrego, N. et al. 2017. Measuring and predicting the influence of traits on the assembly processes
374 of wood-inhabiting fungi. - *J. Ecol.* 105: 1070–1081.
- 375 Arraiano-Castilho, R. et al. 2020. Plant-fungal interactions in hybrid zones: Ectomycorrhizal
376 communities of willows (*Salix*) in an alpine glacier forefield. - *Fungal Ecol.* 45: 100936.
- 377 Artzy-randrup, Y. and Stone, L. 2005. Generating uniformly distributed random networks. - *Phys.*
378 *Rev. E* 72: 056708.
- 379 Bagchi, R. et al. 2014. Pathogens and insect herbivores drive rainforest plant diversity and
380 composition. - *Nature* 506: 85–88.
- 381 Bahram, M. et al. 2014. Network perspectives of ectomycorrhizal associations. - *Fungal Ecol.* 7:
382 70–77.
- 383 Beckett, S. J. 2016. Improved community detection in weighted bipartite networks. - *R. Soc. Open*
384 *Sci.* 3: 140536.
- 385 Bennett, J. A. and Klironomos, J. 2018. Climate, but not trait, effects on plant–soil feedback depend
386 on mycorrhizal type in temperate forests. - *Ecosphere* 9: e02132.
- 387 Bennett, J. A. and Klironomos, J. 2019. Mechanisms of plant–soil feedback: interactions among
388 biotic and abiotic drivers. - *New Phytol.* 222: 91–96.
- 389 Bennett, J. A. et al. 2017. Plant-soil feedbacks and mycorrhizal type influence temperate forest
390 population dynamics. - *Science.* 355: 181–184.
- 391 Bufford, J. L. et al. 2020. Novel interactions between alien pathogens and native plants increase
392 plant–pathogen network connectance and decrease specialization. - *J. Ecol.* 108: 750–760.

- 393 Chagnon, P.-L. et al. 2013. A trait-based framework to understand life history of mycorrhizal fungi.
394 - *Trends Plant Sci.* 18: 484–491.
- 395 Chagnon, P.-L. et al. 2015. Trait-based partner selection drives mycorrhizal network assembly. -
396 *Oikos* 124: 1609–1616.
- 397 Chagnon, P. L. et al. 2018. Strong specificity and network modularity at a very fine phylogenetic
398 scale in the lichen genus *Peltigera*. - *Oecologia* 187: 767–782.
- 399 Chen, W. et al. 2016. Root morphology and mycorrhizal symbioses together shape nutrient foraging
400 strategies of temperate trees. - *Proc. Natl. Acad. Sci. U. S. A.* 113: 8741–8746.
- 401 Chen, L. et al. 2019. Differential soil fungus accumulation and density dependence of trees in a
402 subtropical forest. - *Science*. 366: 124–128.
- 403 Connell, J. H. 1971. On the role of natural enemies in preventing competitive exclusion in some
404 marine animals and in rain forest trees. - In: den Boer P. J. and Gradwell G. R. (ed), *Dynamics*
405 *of Populations*. Centre for Agricultural Publishing and Documentation, Wageningen, pp. 298–
406 312.
- 407 Dassen, S. et al. 2017. Differential responses of soil bacteria, fungi, archaea and protists to plant
408 species richness and plant functional group identity. - *Mol. Ecol.* 26: 4085–4098.
- 409 Davison, J. et al. 2020. Plant functional groups associate with distinct arbuscular mycorrhizal fungal
410 communities. - *New Phytol.* 226: 1117–1128.
- 411 Diniz-Filho, J. A. F. et al. 1998. An eigenvector method for estimating phylogenetic inertia. -
412 *Evolution*. 52: 1247–1262.
- 413 Donatti, C. I. et al. 2011. Analysis of a hyper-diverse seed dispersal network: modularity and

414 underlying mechanisms. - *Ecol. Lett.* 14: 773–781.

415 Dormann, C. F. et al. 2009. Indices, graphs and null models : analyzing bipartite ecological
416 networks. - *Open Ecol. J.* 2: 7–24.

417 Evans, J. S. and Murphy, M. A. 2019. rfUtilities: random forests model selection and performance
418 evaluation. R package version 2.1-5. - <https://github.com/jeffrejevans/rfUtilities>

419 Feng, J. et al. 2018. Comparing shade tolerance measures of woody forest species. - *Peer J* 6:
420 e5736.

421 García-Guzmán, G. and Heil, M. 2014. Life histories of hosts and pathogens predict patterns in
422 tropical fungal plant diseases. - *New Phytol.* 201: 1106–1120.

423 Hardy, O. J. 2008. Testing the spatial phylogenetic structure of local communities: statistical
424 performances of different null models and test statistics on a locally neutral community. - *J.*
425 *Ecol.* 96: 914–926.

426 He, D. et al. 2018. Intra- and interspecific trait variations reveal functional relationships between
427 specific leaf area and soil niche within a subtropical forest. - *Ann. Bot.* 121: 1173–1182.

428 Högberg, M. N. and Högberg, P. 2002. Extramatrical ectomycorrhizal mycelium contributes one-
429 third of microbial biomass and produces, together with associated roots, half the dissolved
430 organic carbon in a forest soil. - *New Phytol.* 154: 791–795.

431 Kembel, S. W. et al. 2010. Picante: R tools for integrating phylogenies and ecology. -
432 *Bioinformatics* 26: 1463–1464.

433 Koorem, K. et al. 2017. Arbuscular mycorrhizal fungal communities in forest plant roots are
434 simultaneously shaped by host characteristics and canopy-mediated light availability. - *Plant*

435 Soil 410: 259–271.

436 Liaw, A. and Wiener, M. 2002. Classification and regression by randomForest. - R News 2: 18–22.

437 Luo, W. et al. 2020. Limiting similarity shapes the functional and phylogenetic structure of root
438 neighborhoods in a subtropical forest. - New Phytol. 229: 1078–1090.

439 Luo, W. et al. 2021. Limited evidence of vertical fine-root segregation in a subtropical forest. - New
440 Phytol. 231: 2308–2318.

441 Montesinos-Navarro, A. et al. 2012. The network structure of plant-arbuscular mycorrhizal fungi. -
442 New Phytol. 194: 536–547.

443 Montoya, J. M. et al. 2006. Ecological networks and their fragility. - Nature 442: 259–264.

444 Montoya, D. et al. 2015. Functional group diversity increases with modularity in complex food
445 webs. - Nat. Commun. 6: 7379.

446 Morente-López, J. et al. 2018. Phenology drives species interactions and modularity in a plant -
447 flower visitor network. - Sci. Rep. 8: 9386.

448 Nguyen, N. H. et al. 2016. FUNGuild: an open annotation tool for parsing fungal community
449 datasets by ecological guild. - Fungal Ecol. 20: 241–248.

450 Olchowik, J. et al. 2021. Post-fire dynamics of ectomycorrhizal fungal communities in a Scots pine
451 (*Pinus sylvestris* L.) forest of Poland. - PeerJ 9: e12076.

452 Olesen, J. M. et al. 2007. The modularity of pollination networks. - Proc. Natl. Acad. Sci. 104:
453 19891–19896.

454 Pölme, S. et al. 2018. Host preference and network properties in biotrophic plant–fungal
455 associations. - New Phytol. 217: 1230–1239.

456 Pölme, S. et al. 2020. FungalTraits: a user-friendly traits database of fungi and fungus-like
457 stramenopiles. - *Fungal Divers.* 105: 1–16.

458 R Core Team 2015. R: A language and environment for statistical computing. - R Found. Stat.
459 Comput. Vienna, Austria.

460 Robinson, K. M. et al. 2015. Relative impacts of environmental variation and evolutionary history
461 on the nestedness and modularity of tree–herbivore networks. - *Ecol. Evol.* 5: 2898–2915.

462 Sachs, J. L. et al. 2004. The evolution of cooperation. - *Chicago Journals* 79: 135–160.

463 Schmid, M. W. et al. 2021. Effects of plant community history, soil legacy and plant diversity on
464 soil microbial communities. - *J. Ecol.* 109: 3007–3023.

465 Schüßler, A. 2002. Molecular phylogeny, taxonomy, and evolution of *Geosiphon pyriformis* and
466 arbuscular mycorrhizal fungi. - *Plant Soil* 244: 75–83.

467 Sepp, S. K. et al. 2019. Non-random association patterns in a plant–mycorrhizal fungal network
468 reveal host–symbiont specificity. - *Mol. Ecol.* 28: 365–378.

469 Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
470 phylogenies. - *Bioinformatics* 30: 1312–1313.

471 Tedersoo, L. and Smith, M. E. 2013. Lineages of ectomycorrhizal fungi revisited: Foraging
472 strategies and novel lineages revealed by sequences from belowground. - *Fungal Biol. Rev.* 27:
473 83–99.

474 Tedersoo, L. et al. 2018. High-level classification of the Fungi and a tool for evolutionary
475 ecological analyses. - *Fungal Divers.* 90: 135–159.

476 Thébault, E. and Fontaine, C. 2010. Stability of ecological communities and the architecture of

477 mutualistic and trophic networks. - *Science*. 329: 853–856.

478 Toju, H. et al. 2014. Assembly of complex plant-fungus networks. - *Nat. Commun.* 5: 5273.

479 Toju, H. et al. 2016. Ericaceous plant–fungus network in a harsh alpine–subalpine environment. -
480 *Mol. Ecol.* 25: 3242–3257.

481 Toju, H. et al. 2018. Network hubs in root-associated fungal metacommunities. - *Microbiome* 6:
482 116.

483 Torrecillas, E. et al. 2014. Modularity reveals the tendency of arbuscular mycorrhizal fungi to
484 interact differently with generalist and specialist plant species in gypsum soils. - *Appl.*
485 *Environ. Microbiol.* 80: 5457–5466.

486 Vacher, C. et al. 2008. Architecture of an antagonistic tree/fungus network: the asymmetric
487 influence of past evolutionary history. - *PLoS One* 3: e1740.

488 Valverde, S. et al. 2020. Coexistence of nestedness and modularity in host–pathogen infection
489 networks. - *Nat. Ecol. Evol.* 4: 568–577.

490 Van Der Heijden, M. G. A. et al. 2015. Mycorrhizal ecology and evolution: the past, the present,
491 and the future. - *New Phytol.* 205: 1406–1423.

492 Wang, Q. et al. 2007. Naive Bayesian classifier for rapid assignment of rRNA sequences into the
493 new bacterial taxonomy. - *Appl. Environ. Microbiol.* 73: 5261–5267.

494 Wang, Z. et al. 2019. Effects of host phylogeny, habitat and spatial proximity on host specificity
495 and diversity of pathogenic and mycorrhizal fungi in a subtropical forest. - *New Phytol.* 223:
496 462–474.

497

498 **Figure legends**

499 **Figure 1.** Modularity of plant–mycorrhizal fungi (plant–MF, a) and plant–pathogenic fungi (plant–
500 PF, b) networks. The plant–MF network contains 16 modules, where plants, ectomycorrhizal fungi
501 (EM), and arbuscular mycorrhizal fungi (AM) are indicated by red triangles, blue circles, and cyan
502 rectangles, respectively. The plant–PF association network contains 9 modules, where plants and PF
503 are indicated by red triangles and blue circles, respectively. The z-score of modularity has been
504 calculated using the modularity indices of the observed plant–fungal networks and the 1000
505 randomly organised networks.

506 **Figure 2.** Differences in plant photosynthetic and root functional traits between modules in plant–
507 mycorrhizal fungal (plant–MF) (a, b) and plant– pathogenic fungal (plant–PF) (c, d) association
508 networks. Bar plots only show partial plant variables at species level that are significantly different
509 among modules, as identified by analysis of variance (ANOVA, see Table S3). These variables
510 include log-transformed specific root length, log-transformed specific root area, log-transformed
511 dark respiration rate and light compensation point. Significance levels are as follows: * ($p < 0.05$),
512 ** ($p < 0.01$), and *** ($p < 0.001$). For modules including > 2 species with available traits, we
513 calculate the standard error of the traits with the error bar shown on the bar. For modules including
514 only one species with available trait, only the mean of the traits is shown. Number of plant species
515 with available functional traits in each module is denoted on the top of a bar.

516 **Figure 3.** Venn diagrams illustrating the variation explained by species traits and phylogeny in
517 assigning root fungi and their host plants to modules. In the association networks of plant–
518 mycorrhizal fungi (MF) and plant–pathogenic fungi (PF), the variation in assigning host plants of

519 fungal MF and PF to modules is partitioned into plant traits and phylogeny (a, c), whereas the
520 variation in assigning host plants for ectomycorrhizal fungi (EM) and arbuscular mycorrhizal fungi
521 (AM) to modules is uniquely explained by plant traits (e, g). The variation in assigning fungi to
522 modules is uniquely explained by the fungal phylogeny for all fungi (b, d, f, h).
523


524 **Table**

525 **Table 1.** The relationships between phylogenetic relatedness between pairs of host plants (or fungi)
 526 and their co-occurrence in a same module in plant-pathogenic fungal (PF) and plant-mycorrhizal
 527 fungal (MF) association networks. Significance effects ($p < 0.05$) are shown in bold type.


Networks	Module composition	No. of species	<i>r</i>	<i>p</i> value⁵²⁸
Plant-PF network	PF hosts	43	-0.034	0.309 ⁵²⁹
	PF	113	-0.044	<0.001
Plant-MF network	MF hosts	43	0.014	0.681 ⁵³⁰
	MF	883	-0.073	<0.001
	EM hosts	43	0.014	0.681 ⁵³¹
	EM	881	-0.075	<0.001 ⁵³²
	AM hosts	43	0.105	0.148
	AM	21	-0.061	0.381

533 **Figure 1.**

(a) plant-MF network




(b) plant-PF network



534

535




537

538


(a) Module MF hosts in plant-MF network




(b) Module MF in plant-MF network




(c) Module PF hosts in plant-PF network




(d) Module PF in plant-PF network




(e) Module EM hosts in plant-MF network



(f) Module EM in plant-MF network



(g) Module AM hosts in plant-MF network



(h) Module AM in plant-MF network

