

ON CURVATURE PROPERTIES OF CERTAIN GENERALIZED ROBERTSON-WALKER SPACETIMES

Dedicated to the memory of Professor Dr. Georgii Ionovich Kruchkovich

By

Ryszard DESZCZ and Marek KUCHARSKI

1. Introduction

The warped product $\bar{M} \times_F N$, of a 1-dimensional manifold (\bar{M}, \bar{g}) , $\bar{g}_{11} = -1$, with a warping function F and a 3-dimensional Riemannian manifold (N, \bar{g}) is said to be a generalized Robertson-Walker spacetime (cf. [2], [32]). In particular, when the manifold (N, \bar{g}) is a Riemannian space of constant curvature, the warped product $\bar{M} \times_F N$ is called a Robertson-Walker spacetime. In [11] it was shown that at every point of a generalized Robertson-Walker spacetime $\bar{M} \times_F N$ the following condition is satisfied:

(*)₁ the tensors $R \cdot R - Q(S, R)$ and $Q(g, C)$ are linearly dependent.

This condition is equivalent to the relation

$$R \cdot R - Q(S, R) = L_1 Q(g, C) \quad (1)$$

on the set \mathcal{U}_C consisting of all points of the manifold $\bar{M} \times_F N$ at which its Weyl tensor C is non-zero, where L_1 is a certain function on \mathcal{U}_C . For precise definitions of the symbols used, we refer to the Sections 2 and 3. (*)₁ is a curvature condition of pseudosymmetry type. In this paper we will investigate generalized Robertson-Walker spacetimes realizing a condition of pseudosymmetry type introduced in [25]. Namely, semi-Riemannian manifolds (M, g) , $n \geq 4$, fulfilling at every point of M the following condition

(*) the tensors $R \cdot C$ and $Q(S, C)$ are linearly dependent.

were considered in [25]. This condition is equivalent to the relation

$$R \cdot C = LQ(S, C) \quad (2)$$

on the set $\mathcal{U} = \{x \in M \mid Q(S, C) \neq 0 \text{ at } x\}$, where L is a certain function on \mathcal{U} . We note that every semisymmetric manifold ($R \cdot R = 0$) as well as every Weyl-semisymmetric manifold ($R \cdot C = 0$) realizes (*) trivially (see [25]). There exist also non semisymmetric and non Weyl-semisymmetric manifolds realizing (*) ([25]). We mention that warped products realizing curvature conditions of pseudosymmetry type were studied in: [7], [8], [9], [11], [13], [14], [15], [16], [17], [19], [20], [21], [24], [26], [28] and [29].

In Section 2 we present a review of the family of curvature conditions of pseudosymmetry type. In the next section we give results on warped products which we apply in the last two sections. In Section 4 we find necessary and sufficient conditions for a warped product to be a manifold satisfying (2). Finally, in Section 5 we present our main results.

Let (M, g) be a semi-Riemannian manifold satisfying (*). We denote by \mathcal{U}_L the set of all points of the set $\mathcal{U} \subset M$ at which the function L is non-zero. It is clear that the tensors $R \cdot C$ and $Q(S, C)$ are non-zero at every point of the set \mathcal{U}_L . Moreover, let (M, g) be a 4-dimensional warped product $\bar{M} \times_F N$, $\dim \bar{M} = 1$. We denote by \mathcal{U}_F the subset of \mathcal{U}_L consisting of all non-critical points of F . Our main result states (see Theorem 5.1) that if the 4-dimensional warped product $\bar{M} \times_F N$, $\dim \bar{M} = 1$, satisfies (*) and the set \mathcal{U}_F is a dense subset of \mathcal{U}_L then the open submanifold U_L of the manifold $\bar{M} \times_F N$ is a pseudosymmetric warped product of the 1-dimensional manifold, with the function F , defined by $F(x^1) = a \exp(bx^1)$, $a = \text{const.} > 0$, $b = \text{const.} \neq 0$, and a 3-dimensional semi-Riemannian manifold such that its Ricci tensor is of rank one and its scalar curvature vanishes identically. From this statement it follows immediately (see Corollary 5.1) that if a generalized Robertson-Walker spacetime $\bar{M} \times_F N$ realizes above assumptions then at every point of $\bar{M} \times_F N$ at least one of the tensors $R \cdot C$ or $Q(S, C)$ must vanish. Finally, using this fact we prove (see Theorem 5.2) that every Robertson-Walker spacetime satisfying (*) is a pseudosymmetric manifold.

2. Curvature Conditions of Pseudosymmetry Type

Let (M, g) be a connected n -dimensional, semi-Riemannian manifold of class C^∞ and let ∇ be its Levi-Civita connection. We define on M the endomorphisms $X \wedge Y$, $\mathcal{R}(X, Y)$ and $\mathcal{C}(X, Y)$ by

$$(X \wedge Y)Z = g(Y, Z)X - g(X, Z)Y, \quad \mathcal{R}(X, Y)Z = [\nabla_X, \nabla_Y]Z - \nabla_{[X, Y]}Z,$$

$$\mathcal{C}(X, Y) = \mathcal{R}(X, Y) - \frac{1}{n-2} \left(X \wedge \mathcal{S}Y + \mathcal{S}X \wedge Y - \frac{\kappa}{n-1} X \wedge Y \right),$$

respectively, where $X, Y, Z \in \Xi(M)$, $\Xi(M)$ being the Lie algebra of vector fields of M . The Ricci operator \mathcal{S} is defined by $S(X, Y) = g(X, \mathcal{S}Y)$, where S is the Ricci tensor and κ the scalar curvature of (M, g) , respectively. Next, we define the tensors U, G , the Riemann-Christoffel curvature tensor R and the Weyl conformal tensor C of (M, g) , by

$$\begin{aligned} U(X_1, X_2, X_3, X_4) &= g(X_1, X_4)S(X_2, X_3) + g(X_2, X_3)S(X_1, X_4) \\ &\quad - g(X_1, X_3)S(X_2, X_4) - g(X_2, X_4)S(X_1, X_3), \\ G(X_1, X_2, X_3, X_4) &= g((X_1 \wedge X_2)X_3, X_4), \\ R(X_1, X_2, X_3, X_4) &= g(\mathcal{R}(X_1, X_2)X_3, X_4), \\ C(X_1, X_2, X_3, X_4) &= g(\mathcal{C}(X_1, X_2)X_3, X_4), \end{aligned}$$

respectively. Now we can present the Weyl tensor C in the following form

$$C = R - \frac{1}{n-2}U + \frac{\kappa}{(n-2)(n-1)}G. \quad (3)$$

For a $(0, k)$ -tensor field $T, k \geq 1$, we define the $(0, k+2)$ -tensors $R \cdot T$ and $Q(g, T)$ by

$$\begin{aligned} (R \cdot T)(X_1, \dots, X_k; X, Y) &= (\mathcal{R}(X, Y) \cdot T)(X_1, \dots, X_k) \\ &= -T(\mathcal{R}(X, Y)X_1, X_2, \dots, X_k) \\ &\quad - \dots - T(X_1, \dots, X_{k-1}, \mathcal{R}(X, Y)X_k), \\ Q(g, T)(X_1, \dots, X_k; X, Y) &= ((X \wedge Y) \cdot T)(X_1, \dots, X_k) \\ &= -T((X \wedge Y)X_1, X_2, \dots, X_k) \\ &\quad - \dots - T(X_1, \dots, X_{k-1}, (X \wedge Y)X_k). \end{aligned}$$

Putting in the last formulas $T = R, T = S$ or $T = C$, we obtain the tensors $R \cdot R, R \cdot S, R \cdot C, Q(g, R), Q(g, S)$ and $Q(g, C)$, respectively. The tensor $C \cdot C$ we define in the same way as the tensor $R \cdot R$.

Let (M, g) be a Riemannian manifold covered by a system of charts $\{\mathcal{W}; x^i\}$. We denote by $g_{rs}, \Gamma'_{st}, R_{rstu}, S_{st}, G_{rstu} = g_{ru}g_{st} - g_{rt}g_{su}$ and

$$C_{rstu} = R_{rstu} - \frac{1}{n-2}(g_{ru}S_{st} - g_{rt}S_{su} + g_{st}S_{ru} - g_{su}S_{rt}) + \frac{\kappa}{(n-2)(n-1)}G_{rstu},$$

the local components of the metric g , the Levi-Civita connection ∇ , the Riemann-Christoffel curvature tensor R , the Ricci tensor S , the tensor G , and the Weyl conformal curvature tensor C of (M, g) , respectively, where $p, q, r, s, t, u, v, w \in \{1, 2, \dots, n\}$. The local components of the tensors $R \cdot R$ and $Q(g, R)$ are given by the following formulas

$$\begin{aligned} (R \cdot R)_{rstuvw} &= \nabla_w \nabla_v R_{rstu} - \nabla_v \nabla_w R_{rstu} \\ &= g^{pq} (R_{pstu} R_{qrvw} - R_{prt u} R_{qsvw} + R_{purs} R_{qtvw} - R_{ptrs} R_{quvw}), \\ Q(g, R)_{rstuvw} &= g_{rv} R_{wstu} + g_{sv} R_{rwtu} + g_{tv} R_{rsvu} + g_{uw} R_{rstv} \\ &\quad - g_{rw} R_{vstu} - g_{sw} R_{rvtu} - g_{tw} R_{rsvu} - g_{uw} R_{rstv}. \end{aligned}$$

A semi-Riemannian manifold (M, g) , $n \geq 2$, is said to be an Einstein manifold if the following condition

$$S = \frac{\kappa}{n} g \tag{4}$$

holds on M . According to [4] (p. 432), (4) is called the Einstein metric condition. Einstein manifolds form a natural subclass of various classes of semi-Riemannian manifolds determined by a curvature condition imposed on their Ricci tensor ([4], Table, pp. 432–433). For instance, every Einstein manifold belongs to the class of semi-Riemannian manifolds (M, g) realizing the following relation

$$\nabla \left(S - \frac{\kappa}{2(n-1)} g \right) (X, Y; Z) = \nabla \left(S - \frac{\kappa}{2(n-1)} g \right) (X, Z; Y), \tag{5}$$

which means that $S - (\kappa/(2(n-1)))g$ is a Codazzi tensor on M . Manifolds of dimension ≥ 4 fulfilling (5) are called manifolds with harmonic Weyl tensor ([4], p. 440). It is known that every warped product $S^1 \times_F M$ of the sphere S^1 , with a positive smooth function F , and an Einstein manifold (M, g) , $\dim M \geq 2$, realizes (5) ([4], p. 433). Such warped product is a non-Einstein manifold, in general. We say that (5) is a generalized Einstein metric condition ([4], chapter XVI). On the other hand, such warped product realizes a condition of pseudosymmetry type too. Namely, the warped product $S^1 \times_F M$ of the sphere S^1 , with a positive smooth function F , and an Einstein manifold (M, g) , $\dim M \geq 2$, is a Ricci-pseudosymmetric manifold ([24], Corollary 3.2). Thus, in particular, the warped product $S^1 \times_F CP^n$ of S^1 , with a positive smooth function \mathcal{F} , and the complex projective space CP^n (considered with its standard Riemannian locally symmetric metric) is a Ricci-pseudosymmetric manifold.

A semi-Riemannian manifold (M, g) , $n \geq 3$, is said to be Ricci-pseudosymmetric ([14], [24]) if at every point of M the following condition is satisfied:

$$(*)_2 \quad \text{the tensors } R \cdot S \text{ and } Q(g, S) \text{ are linearly dependent.}$$

Evidently, any Einstein manifold is Ricci-pseudosymmetric. Thus we see that $(*)_2$ is a generalized Einstein metric condition. The manifold (M, g) is Ricci-pseudosymmetric if and only if

$$R \cdot S = L_S Q(g, S) \tag{6}$$

holds on the set $U_S = \{x \in M \mid S - (\kappa/n)g \neq 0 \text{ at } x\}$, where L_S is some function on U_S . Warped products realizing $(*)_2$ were considered in [14], [17], [24] and [26]. Certain examples of compact and non-Einstein Ricci-pseudosymmetric manifolds were found in [26] and [30]. For instance, in [30] (Theorem 1) it was shown that the Cartan hypersurfaces M in the spheres S^7, S^{17} or S^{25} are non-pseudosymmetric, Ricci-pseudosymmetric manifolds with non-pseudosymmetric Weyl tensor. The Cartan hypersurfaces M in S^4 are non-semisymmetric, pseudosymmetric manifolds. Ricci-pseudosymmetric hypersurfaces immersed isometrically in a semi-Riemannian manifolds of constant curvature were investigated in [10].

A very important subclass of the class of Ricci-pseudosymmetric manifolds form pseudosymmetric manifolds. The semi-Riemannian manifold (M, g) , $n \geq 3$, is said to be pseudosymmetric ([21]) if at every point of M the following condition is satisfied:

$$(*)_3 \quad \text{the tensors } R \cdot R \text{ and } Q(g, R) \text{ are linearly dependent.}$$

The manifold (M, g) is pseudosymmetric if and only if

$$R \cdot R = L_R Q(g, R) \tag{7}$$

holds on the set $U_R = \{x \in M \mid R - (\kappa/(n(n-1)))G \neq 0 \text{ at } x\}$, where L_R is some function on U_R . It is clear that any semisymmetric manifold ($R \cdot R = 0$, [36]) is pseudosymmetric. Very recently the theory of Riemannian semisymmetric manifolds has been presented in [6]. The condition $(*)_3$ arose during the study of totally umbilical submanifolds of semisymmetric manifolds ([1]) as well as when we consider geodesic mappings of semisymmetric manifolds ([18], [37]). There exist many examples of pseudosymmetric manifolds which are not semisymmetric ([13], [19], [20], [21], [28]). Among these examples we can distinguish also compact pseudosymmetric manifolds (for instance, see [19], Example 3.1 and Theorem 4.1). Another example of a compact pseudosymmetric manifold is the warped product $S^1 \times_F S^{n-1}$, with a positive smooth function F , as well as n -dimensional

tori T^n with a certain metric (see [19], Examples 4.1 and 4.2). It is clear that if at a point x of a manifold (M, g) $(*)_3$ is satisfied then also $(*)_2$ holds at x . The converse statement is not true. E.g. every warped product $M_1 \times_F M_2$, $\dim M_1 = 1$, $\dim M_2 \geq 3$, of a manifold (M_1, \bar{g}) and a non-pseudosymmetric, Einstein manifold (M_2, \bar{g}) is a non-pseudosymmetric, Ricci-pseudosymmetric manifold (cf. [24], Remark 3.4 and [21], Theorem 4.1).

It is easy to see that if $(*)_3$ holds on a semi-Riemannian manifold (M, g) , $n \geq 4$, then at every point of M the following condition is satisfied:

$(*)_4$ the tensors $R \cdot C$ and $Q(g, C)$ are linearly dependent.

Manifolds fulfilling $(*)_4$ are called Weyl-pseudosymmetric. Weyl-pseudosymmetric manifolds has been studied in [15], [17] and [22]. The manifold (M, g) is a Weyl-pseudosymmetric manifold if and only if the relation $R \cdot C = L_2 Q(g, C)$ holds on the set $\mathcal{U}_C = \{x \in M \mid C \neq 0 \text{ at } x\}$, where L_2 is some function on \mathcal{U}_C .

A semi-Riemannian manifold (M, g) , $n \geq 4$, is said to be a manifold with pseudosymmetric Weyl tensor ([29]) if at every point of M the following condition is satisfied:

$(*)_5$ the tensors $C \cdot C$ and $Q(g, C)$ are linearly dependent.

Thus (M, g) is a manifold with pseudosymmetric Weyl tensor if and only if the relation $C \cdot C = L_3 Q(g, C)$ holds on the set \mathcal{U}_C , where L_3 is a certain function on \mathcal{U}_C . The condition $(*)_5$ arose during the study of 4-dimensional warped products ([17]). Namely, in [17] (Theorem 2) it was shown that at every point of a warped product $M_1 \times_F M_2$, with $\dim M_1 = \dim M_2 = 2$, $(*)_5$ is fulfilled. Many examples of manifolds satisfying $(*)_5$ are presented in [9]. For instance, the Cartesian product of two manifolds of constant curvature is a manifold realizing $(*)_5$. Warped products satisfying $(*)_5$ were considered in [29]. In [9] it was shown that the classes of manifolds realizing $(*)_3$ and $(*)_5$ do not coincide. However, there exist pseudosymmetric manifolds fulfilling $(*)_5$, e.g. Einsteinian pseudosymmetric manifolds ([9], Theorem 3.1). Curvature properties of pseudosymmetric manifolds with pseudosymmetric Weyl tensor were obtained in [31].

For a $(0, k)$ -tensor field $T, k \geq 1$, and a symmetric $(0, 2)$ -tensor field A , we define the $(0, k + 2)$ -tensor $Q(A, T)$ by

$$\begin{aligned} Q(A, T)(X_1, \dots, X_k; X, Y) &= ((X \wedge_A Y) \cdot T)(X_1, \dots, X_k) \\ &= -T((X \wedge_A Y)X_1, X_2, \dots, X_k) \\ &\quad - \dots - T(X_1, \dots, X_{k-1}, (X \wedge_A Y)X_k), \end{aligned}$$

where $X \wedge_A Y$ is the endomorphism defined by

$$(X \wedge_A Y)Z = A(Y, Z)X - A(X, Z)Y.$$

In particular, we have $X \wedge_g Y = X \wedge Y$. Putting in the above formula $A = S$ and $T = R, T = C$ or $T = G$, we obtain the tensors $Q(S, R), Q(S, C)$ and $Q(S, G)$, respectively.

A semi-Riemannian manifold (M, g) is said to be Ricci-generalized pseudosymmetric ([7]) if at every point of M the following condition is satisfied:

$$(*)_6 \quad \text{the tensors } R \cdot R \text{ and } Q(S, R) \text{ are linearly dependent.}$$

A very important subclass of Ricci-generalized pseudosymmetric manifolds form manifolds fulfilling the following relation $R \cdot R = Q(S, R)$ ([7], [8], [23]). Every 3-manifold (M, g) as well as every hypersurface M immersed isometrically in an $(n + 1)$ -dimensional semi-Euclidean space E_s^{n+1} , of index $s, n \geq 3$, fulfils the last equality, see [16] (Theorem 3.1) and [27] (Corollary 3.1), respectively.

As it was shown in [27], every hypersurface M in a semi-Riemannian space of constant curvature $M^{n+1}(c), n \geq 4$, fulfils (1). More precisely, we have the following

REMARK 2.1 ([27], Proposition 3.1). *Every hypersurface M immersed isometrically in a semi-Riemannian space of constant curvature $M^{n+1}(c), n \geq 4$, satisfies the equality $R \cdot R - Q(S, R) = -(((n - 2)\tilde{\kappa})/(n(n + 1)))Q(g, C)$, where $\tilde{\kappa}$ is the scalar curvature of $M^{n+1}(c)$ and R, S and C are the curvature tensor, the Ricci tensor and the Weyl tensor of M , respectively.*

Using Theorem 3.1 of [16], which was mentioned above, and the fact that the Weyl tensor of every 3-dimensional semi-Riemannian manifold vanishes identically, we conclude that $(*)_1$ is trivially satisfied on any 3-dimensional semi-Riemannian manifold. Recently, warped products realizing $(*)_1$ were considered in [11].

The relations $(*), (*), (*), (*)_6$ are called conditions of pseudosymmetry type. We refer to [12], [18] and [37] as the review papers on semi-Riemannian manifolds satisfying such conditions. A hypersurface fulfilling a curvature condition of pseudosymmetry type is said to be a hypersurface of pseudosymmetry type ([12]). We finish this section with the following

LEMMA 2.1. *Let $(M, g), n = \dim M \geq 3$, be a semi-Riemannian manifold.*

(i) ([13], Lemma 1.2; [23], Lemma 2) *If the Weyl tensor C of (M, g) vanishes at a point $x \in M$ then at x any of the following three identities is equivalent to each*

other:

$$R \cdot R = \alpha Q(g, R), \quad R \cdot S = \alpha Q(g, S),$$

$$\left(\frac{\kappa}{n-1} + (n-2)\alpha \right) \left(S - \frac{\kappa}{n} g \right) = S^2 - \frac{1}{n} \text{tr}(S^2)g,$$

where $\alpha \in \mathbf{R}$.

(ii) ([3], Lemma 3.1) The following identity is fulfilled on M : $Q(S, G) = -Q(g, U)$.

(iii) ([16], Theorem 3.1) If $\dim M = 3$ then $R \cdot R = Q(S, R)$ holds on M .

(iv) If the following conditions are fulfilled at a point $x \in M$: $C = 0$, $\text{rank}(S) = 1$ and $\kappa = 0$, then $R \cdot R = 0$ holds at x .

PROOF. (iv) The condition $\text{rank}(S) = 1$ we can present in the following form

$$S_{ij} = \beta u_i u_j, \quad u \in T_x^*(M), \quad \beta \in \mathbf{R}, \quad (8)$$

where u_i are the local components of u . From (8), by $\kappa = 0$, it follows that $\beta g^{ij} u_i u_j = 0$. Transvecting now (8) with $u^i = g^{ir} u_r$ we get $u^r S_{ij} = 0$. Next, transvecting (8) with S_k^i and using the last relation we get $S_{ij}^2 = 0$ which, in view of (i), completes the proof.

3. Warped Products

Let now (\bar{M}, \bar{g}) and (N, \tilde{g}) , $\dim \bar{M} = p$, $\dim N = n - p$, $1 \leq p < n$, be semi-Riemannian manifolds covered by systems of charts $\{\mathcal{U}; x^a\}$ and $\{\mathcal{V}; y^\alpha\}$, respectively. Let F be a positive smooth function on \bar{M} . The warped product $\bar{M} \times_F N$ of (\bar{M}, \bar{g}) and (N, \tilde{g}) ([5], [33]) is the product manifold $\bar{M} \times N$ with the metric $g = \bar{g} \times_F \tilde{g}$ defined by

$$\bar{g} \times_F \tilde{g} = \pi_1^* \bar{g} + (F \circ \pi_1) \pi_2^* \tilde{g},$$

where $\pi_1 : \bar{M} \times N \rightarrow \bar{M}$ and $\pi_2 : \bar{M} \times N \rightarrow N$ are the natural projections on \bar{M} and N , respectively. Let $\{\mathcal{U} \times \mathcal{V}; x^1, \dots, x^p, x^{p+1} = y^1, \dots, x^n = y^{n-p}\}$ be a product chart for $\bar{M} \times N$. The local components of the metric $g = \bar{g} \times_F \tilde{g}$ with respect to this chart are the following $g_{rs} = \bar{g}_{ab}$ if $r = a$ and $s = b$, $g_{rs} = F \tilde{g}_{\alpha\beta}$ if $r = \alpha$ and $s = \beta$, and $g_{rs} = 0$ otherwise, where $a, b, c, \dots \in \{1, \dots, p\}$, $\alpha, \beta, \gamma, \dots \in \{p+1, \dots, n\}$ and $r, s, t, \dots \in \{1, 2, \dots, n\}$. We will denote by bars (resp., by tildes) tensors formed from \bar{g} (resp., \tilde{g}). The local components Γ_{st}^r of the Levi-Civita connection ∇ of $\bar{M} \times_F N$ are the following ([34]):

$$\begin{aligned}\Gamma_{bc}^a &= \bar{\Gamma}_{bc}^a, & \Gamma_{\beta\gamma}^\alpha &= \tilde{\Gamma}_{\beta\gamma}^\alpha, & \Gamma_{\alpha\beta}^a &= -\frac{1}{2}\bar{g}^{ab}F_b\tilde{g}_{\alpha\beta}, & \Gamma_{\alpha\beta}^\alpha &= \frac{1}{2F}F_a\delta_\beta^\alpha, \\ \Gamma_{\alpha\beta}^a &= \Gamma_{\alpha\beta}^\alpha = 0, & F_a &= \partial_a F = \frac{\partial F}{\partial x^a}, & \partial_a &= \frac{\partial}{\partial x^a}.\end{aligned}$$

The local components

$$R_{rstu} = g_{rw}R_{stu}^w = g_{rw}(\partial_u\Gamma_{st}^w - \partial_t\Gamma_{su}^w + \Gamma_{st}^v\Gamma_{vu}^w - \Gamma_{su}^v\Gamma_{vt}^w), \quad \partial_u = \frac{\partial}{\partial x^u},$$

of the Riemann-Christoffel curvature tensor R and the local components S_{ts} of the Ricci tensor S of the warped product $\bar{M} \times_F N$ which may not vanish identically are the following:

$$R_{abcd} = \bar{R}_{abcd}, \quad R_{x\alpha\beta} = -\frac{1}{2}T_{\alpha\beta}\tilde{g}_{\alpha\beta}, \quad R_{\alpha\beta\gamma\delta} = F\tilde{R}_{\alpha\beta\gamma\delta} - \frac{1}{4}\Delta_1 F\tilde{G}_{\alpha\beta\gamma\delta}, \quad (9)$$

$$S_{ab} = \bar{S}_{ab} - \frac{n-p}{2}\frac{1}{F}T_{ab}, \quad S_{\alpha\beta} = \tilde{S}_{\alpha\beta} - \frac{1}{2}\left(\text{tr}(T) + \frac{n-p-1}{2F}\Delta_1 F\right)\tilde{g}_{\alpha\beta}, \quad (10)$$

where

$$T_{ab} = \bar{\nabla}_b F_a - \frac{1}{2F}F_a F_b, \quad \text{tr}(T) = \bar{g}^{ab}T_{ab}, \quad \Delta_1 F = \Delta_{1\bar{g}}F = \bar{g}^{ab}F_a F_b, \quad (11)$$

and T is the $(0, 2)$ -tensor with the local components T_{ab} . The scalar curvature κ of $\bar{M} \times_F N$ satisfies the following relation

$$\kappa = \bar{\kappa} + \frac{1}{F}\tilde{\kappa} - \frac{n-p}{F}\left(\text{tr}(T) + \frac{n-p-1}{4F}\Delta_1 F\right). \quad (12)$$

From now we assume that $\dim \bar{M} \times_F N = 4$ and $\dim \bar{M} = 1$. Then (9), (10) and (12) turn into

$$R_{x11\beta} = -\frac{1}{2}T_{11}\tilde{g}_{\alpha\beta}, \quad R_{\alpha\beta\gamma\delta} = F\tilde{R}_{\alpha\beta\gamma\delta} - \frac{1}{4}\Delta_1 F\tilde{G}_{\alpha\beta\gamma\delta}, \quad (13)$$

$$S_{11} = -\frac{3}{2F}T_{11}, \quad S_{\alpha\beta} = \tilde{S}_{\alpha\beta} - \frac{1}{2}\left(\text{tr}(T) + \frac{\Delta_1 F}{F}\right)\tilde{g}_{\alpha\beta}, \quad (14)$$

$$\kappa = \frac{1}{F}\tilde{\kappa} - \frac{3}{F}\left(\text{tr}(T) + \frac{1}{2}\frac{\Delta_1 F}{F}\right). \quad (15)$$

respectively. Further, by making use of (13), (14) and (15), we obtain the following relations (see [17], Lemma 6):

$$\begin{aligned}
C_{\alpha 11\beta} &= -\frac{1}{2}\bar{g}_{11}\left(\tilde{S}_{\alpha\beta} - \frac{1}{3}\tilde{\kappa}\tilde{g}_{\alpha\beta}\right), \\
C_{\alpha\beta\gamma\delta} &= \frac{1}{2}F(\tilde{g}_{\alpha\delta}\tilde{S}_{\beta\gamma} + \tilde{g}_{\beta\gamma}\tilde{S}_{\alpha\delta} - \tilde{g}_{\alpha\gamma}\tilde{S}_{\beta\delta} - \tilde{g}_{\beta\delta}\tilde{S}_{\alpha\gamma}) - \frac{1}{3}F\tilde{\kappa}\tilde{G}_{\alpha\beta\gamma\delta}. \quad (16)
\end{aligned}$$

On the other hand, from (3) it follows that

$$\tilde{C}_{\alpha\beta\gamma\delta} = \tilde{R}_{\alpha\beta\gamma\delta} - \tilde{U}_{\alpha\beta\gamma\delta} + \frac{1}{2}\tilde{\kappa}\tilde{G}_{\alpha\beta\gamma\delta}. \quad (17)$$

Since $\tilde{C}_{\alpha\beta\gamma\delta} = 0$, the last identity reduces to

$$\tilde{U}_{\alpha\beta\gamma\delta} = \tilde{R}_{\alpha\beta\gamma\delta} + \frac{1}{2}\tilde{\kappa}\tilde{G}_{\alpha\beta\gamma\delta}. \quad (18)$$

Now (16) turns into

$$C_{\alpha 11\beta} = -\frac{1}{2}\bar{g}_{11}(\tilde{S}_{\alpha\beta} - \frac{1}{3}\tilde{\kappa}\tilde{g}_{\alpha\beta}), \quad C_{\alpha\beta\gamma\delta} = \frac{1}{2}F\tilde{R}_{\alpha\beta\gamma\delta} - \frac{1}{12}F\tilde{\kappa}\tilde{G}_{\alpha\beta\gamma\delta}. \quad (19)$$

4. Preliminary Results

Let $\bar{M} \times_F N$ be a 4-dimensional warped product with 1-dimensional base manifold (\bar{M}, \bar{g}) . Using (13), (14), (18) and (19), we can verify that the local components of the tensors $R \cdot C$ and $Q(S, C)$ of the manifold $\bar{M} \times_F N$, which may not vanish identically are the following:

$$(R \cdot C)_{\alpha\beta\gamma\delta\lambda\mu} = \frac{1}{2}F(\tilde{R} \cdot \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} - \frac{1}{8}\Delta_1 F Q(\tilde{g}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu}, \quad (20)$$

$$\begin{aligned}
Q(S, C)_{\alpha\beta\gamma\delta\lambda\mu} &= -\frac{1}{4}F\left(\text{tr}(T) + \frac{\Delta_1 F}{F}\right)Q(\tilde{g}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} \\
&\quad + \frac{1}{2}FQ(\tilde{S}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} - \frac{1}{12}F\tilde{\kappa}Q(\tilde{S}, \tilde{G})_{\alpha\beta\gamma\delta\lambda\mu}, \quad (21)
\end{aligned}$$

$$(R \cdot C)_{1\alpha\beta\gamma 1\delta} = -\frac{1}{2}\frac{1}{F}T_{11}C_{\delta\alpha\beta\gamma} + \frac{1}{12}\tilde{\kappa}T_{11}\tilde{G}_{\delta\alpha\beta\gamma} - \frac{1}{4}T_{11}(\tilde{g}_{\gamma\delta}\tilde{S}_{\alpha\beta} - \tilde{g}_{\beta\delta}\tilde{S}_{\alpha\gamma}), \quad (22)$$

$$\begin{aligned}
Q(S, C)_{1\alpha\beta\gamma 1\delta} &= -\frac{3}{2}\frac{1}{F}T_{11}C_{\delta\alpha\beta\gamma} + \frac{1}{2}\bar{g}_{11}\left(\tilde{S}_{\delta\gamma}\tilde{S}_{\alpha\beta} - \tilde{S}_{\delta\beta}\tilde{S}_{\alpha\gamma} - \frac{1}{3}\tilde{\kappa}(\tilde{g}_{\alpha\beta}\tilde{S}_{\delta\gamma} - \tilde{g}_{\alpha\gamma}\tilde{S}_{\delta\beta})\right) \\
&\quad + \frac{1}{12}\tilde{\kappa}\left(\text{tr}(T) + \frac{\Delta_1 F}{F}\right)\bar{g}_{11}\tilde{G}_{\delta\alpha\beta\gamma} \\
&\quad - \frac{1}{4}\bar{g}_{11}\left(\text{tr}(T) + \frac{\Delta_1 F}{F}\right)(\tilde{g}_{\delta\gamma}\tilde{S}_{\alpha\beta} - \tilde{g}_{\delta\beta}\tilde{S}_{\alpha\gamma}), \quad (23)
\end{aligned}$$

$$(R \cdot C)_{\alpha 11\beta\gamma\delta} = -\frac{1}{2} \frac{1}{F} \bar{g}_{11} \left(F(\tilde{R} \cdot \tilde{S})_{\alpha\beta\gamma\delta} - \frac{1}{4} \Delta_1 F Q(\tilde{g}, \tilde{S})_{\alpha\beta\gamma\delta} \right), \quad (24)$$

$$Q(S, C)_{\alpha 11\beta\gamma\delta} = -\frac{1}{2} \bar{g}_{11} \left(\frac{1}{3} \tilde{\kappa} - \frac{1}{2} \left(\text{tr}(T) + \frac{\Delta_1 F}{F} \right) \right) Q(\tilde{g}, \tilde{S})_{\alpha\beta\gamma\delta}. \quad (25)$$

From Lemma 2.1(ii) it follows that $Q(\tilde{S}, \tilde{G})_{\alpha\beta\gamma\delta\lambda\mu} = -Q(\tilde{g}, \tilde{U})_{\alpha\beta\gamma\delta\lambda\mu}$, which by making use of (18), turns into $Q(\tilde{S}, \tilde{G})_{\alpha\beta\gamma\delta\lambda\mu} = -Q(\tilde{g}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu}$. Now (21) takes the form

$$Q(S, C)_{\alpha\beta\gamma\delta\lambda\mu} = \frac{1}{2} F \left(Q(\tilde{S}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} + \left(\frac{1}{6} \tilde{\kappa} - \frac{1}{2} \left(\text{tr}(T) + \frac{\Delta_1 F}{F} \right) \right) Q(\tilde{g}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} \right). \quad (26)$$

In view of Lemma 2.1(iii) we have also $(\tilde{R} \cdot \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} = Q(\tilde{S}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu}$. Substituting this in (26) we obtain

$$Q(S, C)_{\alpha\beta\gamma\delta\lambda\mu} = \frac{1}{2} F (\tilde{R} \cdot \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} + \left(\frac{1}{6} \tilde{\kappa} - \frac{1}{2} \left(\text{tr}(T) + \frac{\Delta_1 F}{F} \right) \right) Q(\tilde{g}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu},$$

whence

$$Q(S, C)_{\alpha\beta\gamma\delta\lambda\mu} = \frac{1}{2} F (\tilde{R} \cdot \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} - \frac{1}{2} F \tau Q(\tilde{g}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu}, \quad (27)$$

$$\tau_1 = \frac{1}{2} \left(-\frac{1}{3} \tilde{\kappa} + \text{tr}(T) + \frac{\Delta_1 F}{F} \right). \quad (28)$$

Now, the equality $(R \cdot C)_{\alpha\beta\gamma\delta\lambda\mu} = LQ(S, C)_{\alpha\beta\gamma\delta\lambda\mu}$, in virtue of (20) and (27), gives

$$(1 - L)(\tilde{R} \cdot \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} = \left(\frac{1}{4} \frac{\Delta_1 F}{F} - \tau_1 L \right) Q(\tilde{g}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu}. \quad (29)$$

By (22) and (23) the relation $(R \cdot C)_{1\alpha\beta\gamma 1\delta} = LQ(S, C)_{1\alpha\beta\gamma 1\delta}$ is equivalent to

$$\begin{aligned} & -\frac{1}{2} \frac{1}{F} T_{11} C_{\delta\alpha\beta\gamma} + \frac{1}{12} \tilde{\kappa} T_{11} \tilde{G}_{\delta\alpha\beta\gamma} - \frac{1}{4} T_{11} (\tilde{g}_{\gamma\delta} \tilde{S}_{\alpha\beta} - \tilde{g}_{\beta\delta} \tilde{S}_{\alpha\gamma}) \\ & = -\frac{3}{2} \frac{1}{F} L T_{11} C_{\delta\alpha\beta\gamma} + \frac{1}{2} L \bar{g}_{11} (\tilde{S}_{\gamma\delta} \tilde{S}_{\alpha\beta} - \tilde{S}_{\beta\delta} \tilde{S}_{\alpha\gamma}) + \frac{1}{12} \tilde{\kappa} \left(\text{tr}(T) + \frac{\Delta_1 F}{F} \right) L \bar{g}_{11} \tilde{G}_{\delta\alpha\beta\gamma} \\ & \quad + \frac{1}{2} L \bar{g}_{11} \left(-\frac{1}{2} \left(\text{tr}(T) + \frac{\Delta_1 F}{F} \right) (\tilde{g}_{\delta\gamma} \tilde{S}_{\alpha\beta} - \tilde{g}_{\delta\beta} \tilde{S}_{\alpha\gamma}) - \frac{1}{3} \tilde{\kappa} (\tilde{g}_{\alpha\beta} \tilde{S}_{\delta\gamma} - \tilde{g}_{\alpha\gamma} \tilde{S}_{\delta\beta}) \right). \quad (30) \end{aligned}$$

Further, we can check that the relation $(R \cdot C)_{\alpha 11\beta\gamma\delta} = L(S, C)_{\alpha 11\beta\gamma\delta}$ turns into

$$(\tilde{R} \cdot \tilde{S})_{\alpha\beta\gamma\delta} = \left(\frac{1}{4} \frac{\Delta_1 F}{F} - \tau_1 L \right) Q(\tilde{g}, \tilde{S})_{\alpha\beta\gamma\delta}. \quad (31)$$

This, in view of Lemma 2.1(iii), is equivalent to

$$(\tilde{R} \cdot \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} = \left(\frac{1}{4} \frac{\Delta_1 F}{F} - \tau_1 L \right) Q(\tilde{g}, \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu}. \quad (32)$$

Thus we have the following

PROPOSITION 4.1. *A 4-dimensional warped product $\bar{M} \times_F N$, $\dim \bar{M} = 1$, satisfies the condition $R \cdot C = LQ(S, C)$ if and only if (29), (30) and (32) hold on \mathcal{U} .*

5. Main Results

EXAMPLE 5.1. (i) *We present an example of a 4-dimensional warped product, with 1-dimensional base manifold, realizing $(*)$ and $(*)_3$. Let (N, \tilde{g}) , $\dim N = 3$, be a semi-Riemannian manifold such that its Ricci tensor \tilde{S} is of rank one and its scalar curvature $\tilde{\kappa}$ vanishes identically on N . Then, in view of Lemma 2.1(iv), (N, \tilde{g}) is a semisymmetric manifold. Furthermore, let F , defined by $F(x^1) = a \exp(bx^1)$, $a = \text{const.} > 0, b = \text{const.} \neq 0$, be a function on a 1-dimensional manifold (\bar{M}, g_1) . It is easy to check, that $\bar{M} \times_F N$ realizes (29), (30) and (32), with $L = 1/3$. Thus, in view of Proposition 4.1, $\bar{M} \times_F N$ fulfils $R \cdot C = (1/3)Q(S, C)$. From Corollary 4.2 of [21] it follows that the manifold $\bar{M} \times_F N$ is pseudosymmetric too. Next, using (3.12) of [21] and (15), we get $R \cdot R = (1/12)\kappa Q(g, R)$, where k is the scalar curvature of $\bar{M} \times_F N$.*

(ii) *We present an example of a 3-dimensional semisymmetric warped product such that the rank of its Ricci tensor is one and its scalar curvature vanishes identically. Let $M_2 = \{(x^2, x^3) : x^2, x^3 \in \mathbf{R}\}$ be a connected, non-empty, open subset of \mathbf{R}^2 , equipped with the metric tensor g_2 defined by $g_{2,22} = g_{2,33} = 0$, $g_{2,23} = g_{2,32} = 1$, and let $H = H(x^2)$ be a smooth function on M_2 . Moreover, let (M_3, g_3) be a 1-dimensional manifold. In [35] (see p. 177) it was shown that the rank of the Ricci tensor \tilde{S} of the warped product $M_2 \times_H M_3$ is equal to one and that the scalar curvature of this manifold vanishes identically. Moreover, we have (cf. [35], p. 177)*

$$\tilde{S}_{22} = -\frac{1}{H} \left(\frac{\partial H_2}{\partial x^2} - \frac{1}{2H} H_2 H_2 \right), \quad H_2 = \frac{\partial H}{\partial x^2}, \quad \tilde{S}_{33} = 0, \quad \tilde{S}_{44} = 0.$$

Furthermore, from Lemma 2.1 (iv) it follows that $M_2 \times_H M_3$ is a semisymmetric manifold. (iii) *We consider the warped product $\bar{M} \times_F N$, where $\dim \bar{M}$*

$= 1$, the warping function F is defined by $F(x^1) = a \exp(bx^1)$, $a = \text{const.} > 0$, $b = \text{const.} \neq 0$, and (N, \tilde{g}) is a semisymmetric manifold defined in (ii). We can verify that the tensor $S - (\kappa/4)g$ is of rank one, i.e. the warped product $\bar{M} \times_F N$ is a quasi-Einstein manifold.

In this section we prove, that under certain assumptions every 4-dimensional warped product $\bar{M} \times_F N$, $\dim \bar{M} = 1$, realizing $(*)$ is the manifold described in Example 5.1(i).

Symmetrizing (30) in α, δ we obtain

$$\left(\frac{1}{2}T_{11} - \tau L\tilde{g}_{11}\right)Q(\tilde{g}, \tilde{S})_{\alpha\beta\gamma\delta} = 0, \quad (33)$$

where

$$\tau = \frac{1}{2}\left(-\frac{2}{3}\tilde{\kappa} + \text{tr}(T) + \frac{\Delta_1 F}{F}\right). \quad (34)$$

From (19) it follows that the Weyl tensor C of every 4-dimensional warped product $\bar{M} \times_F N$, $\dim \bar{M} = 1$, vanishes at a point $x \in M_1 \times_F N$ if and only

$$\tilde{S}_{\alpha\beta} = \frac{1}{3}\tilde{\kappa}\tilde{g}_{\alpha\beta}. \quad (35)$$

holds at x . We note also that $Q(\tilde{g}, \tilde{S})$ vanishes at x if and only if (35) is satisfied at x . So, if the tensor C is non-zero at the point $x \in M_1 \times_F N$ then from (33) it follows that

$$\frac{1}{2}T_{11} = \tau L\tilde{g}_{11} \quad (36)$$

holds at x . Applying (36) in (30) we obtain

$$\begin{aligned} & -\frac{\tau}{F}C_{\delta\alpha\beta\gamma} + \frac{\tau}{6}\tilde{\kappa}\tilde{G}_{\delta\alpha\beta\gamma} - \frac{\tau}{2}(\tilde{g}_{\delta\gamma}\tilde{S}_{\alpha\beta} - \tilde{g}_{\delta\beta}\tilde{S}_{\alpha\gamma}) \\ & = -3\frac{\tau}{F}C_{\delta\alpha\beta\gamma} + \frac{1}{2}(\tilde{S}_{\delta\gamma}\tilde{S}_{\alpha\beta} - \tilde{S}_{\delta\beta}\tilde{S}_{\alpha\gamma}) + \frac{1}{12}\tilde{\kappa}\left(\text{tr}(T) + \frac{\Delta_1 F}{F}\right)\tilde{G}_{\delta\alpha\beta\gamma} \\ & \quad - \frac{1}{4}\left(\text{tr}(T) + \frac{\Delta_1 F}{F}\right)(\tilde{g}_{\delta\gamma}\tilde{S}_{\alpha\beta} - \tilde{g}_{\delta\beta}\tilde{S}_{\alpha\gamma}) - \frac{1}{6}\tilde{\kappa}(\tilde{g}_{\alpha\beta}\tilde{S}_{\gamma\delta} - \tilde{g}_{\alpha\gamma}\tilde{S}_{\delta\beta}). \end{aligned} \quad (37)$$

If $x \in U_L$ then the last equality reduces to

$$\begin{aligned} & \frac{1}{F} \tau(3L-1) C_{\delta\alpha\beta\gamma} + \frac{1}{6} \tilde{\kappa} \left(\tau - \frac{1}{2} \left(\text{tr}(T) + \frac{\Delta_1 F}{F} \right) \right) \tilde{G}_{\delta\alpha\beta\gamma} \\ &= \frac{1}{2} (\tilde{S}_{\gamma\delta} \tilde{S}_{\alpha\beta} - \tilde{S}_{\delta\beta} \tilde{S}_{\alpha\gamma}) - \frac{1}{6} \tilde{\kappa} (\tilde{g}_{\alpha\beta} \tilde{S}_{\delta\gamma} - \tilde{g}_{\alpha\gamma} \tilde{S}_{\beta\delta} + \tilde{g}_{\gamma\delta} \tilde{S}_{\alpha\beta} - \tilde{g}_{\delta\beta} \tilde{S}_{\alpha\gamma}), \end{aligned}$$

which, by (34), turns into

$$\frac{2}{F} \tau(3L-1) C_{\delta\alpha\beta\gamma} - \frac{1}{9} \tilde{\kappa}^2 \tilde{G}_{\delta\alpha\beta\gamma} = \tilde{S}_{\gamma\delta} \tilde{S}_{\alpha\beta} - \tilde{S}_{\delta\beta} \tilde{S}_{\alpha\gamma} - \frac{1}{3} \tilde{\kappa} \tilde{U}_{\delta\alpha\beta\gamma}. \quad (38)$$

On the other hand (18) and (19) give

$$C_{\delta\alpha\beta\gamma} = \frac{1}{2} F \left(\tilde{U}_{\delta\alpha\beta\gamma} - \frac{2}{3} \tilde{\kappa} \tilde{G}_{\delta\alpha\beta\gamma} \right).$$

Applying this in (38) we obtain

$$\tau(3L-1) \tilde{U}_{\delta\alpha\beta\gamma} - \frac{2}{3} \tau(3L-1) \tilde{\kappa} \tilde{G}_{\delta\alpha\beta\gamma} - \frac{1}{9} \tilde{\kappa}^2 \tilde{G}_{\delta\alpha\beta\gamma} = \tilde{S}_{\gamma\delta} \tilde{S}_{\alpha\beta} - \tilde{S}_{\delta\beta} \tilde{S}_{\alpha\gamma} - \frac{1}{3} \tilde{\kappa} \tilde{U}_{\delta\alpha\beta\gamma},$$

whence

$$\tilde{S}_{\gamma\delta} \tilde{S}_{\alpha\beta} - \tilde{S}_{\delta\beta} \tilde{S}_{\alpha\gamma} = \rho \tilde{U}_{\delta\alpha\beta\gamma} + \mu \tilde{G}_{\delta\alpha\beta\gamma}, \quad (39)$$

where

$$\rho = \tau(3L-1) + \frac{1}{3} \tilde{\kappa}, \quad \mu = -\frac{1}{3} \tilde{\kappa} \left(2\tau(3L-1) + \frac{1}{3} \tilde{\kappa} \right). \quad (40)$$

We put $\tilde{A}_{\alpha\beta} = \tilde{S}_{\alpha\beta} - \rho \tilde{g}_{\alpha\beta}$. Thus, by (39), we have

$$\begin{aligned} \tilde{A}_{\gamma\delta} \tilde{A}_{\alpha\beta} - \tilde{A}_{\beta\delta} \tilde{A}_{\alpha\gamma} &= \tilde{S}_{\gamma\delta} \tilde{S}_{\alpha\beta} - \tilde{S}_{\beta\delta} \tilde{S}_{\alpha\gamma} + \rho^2 \tilde{G}_{\delta\alpha\beta\gamma} \\ &\quad - \rho (\tilde{g}_{\gamma\delta} \tilde{S}_{\alpha\beta} + \tilde{g}_{\alpha\beta} \tilde{S}_{\gamma\delta} - \tilde{g}_{\beta\delta} \tilde{S}_{\alpha\gamma} - \tilde{g}_{\alpha\gamma} \tilde{S}_{\beta\delta}) \\ &= (\rho^2 + \mu) \tilde{G}_{\delta\alpha\beta\gamma}, \end{aligned} \quad (41)$$

which leads to

$$Q(\tilde{A}, A) = (\rho^2 + \mu) Q(\tilde{A}, \tilde{G}),$$

where the (0,4)-tensor A is defined by

$$A_{\alpha\beta\gamma\delta} = \tilde{A}_{\alpha\delta} \tilde{A}_{\beta\gamma} - \tilde{A}_{\alpha\gamma} \tilde{A}_{\beta\delta}.$$

Since the tensor $Q(\tilde{A}, A)$ vanishes identically, we have $(\rho^2 + \mu) Q(\tilde{A}, \tilde{G}) = 0$, whence we get easily $(\rho^2 + \mu)(\tilde{A} - (1/3)\text{tr}(\tilde{A})\tilde{g}) = 0$. Since $\tilde{S} \neq (1/3)\tilde{\kappa}\tilde{g}$ holds at

x , the last relation yields

$$\rho^2 + \mu = 0. \quad (42)$$

Further, using (29) and (32) we deduce that

$$(a) \quad (\tilde{R} \cdot \tilde{R})_{\alpha\beta\gamma\delta\lambda\mu} = 0, \quad (b) \quad \frac{1}{4} \frac{\Delta_1 F}{F} = \tau L, \quad (43)$$

hold at $x \in \mathcal{U}_L$. Further, contracting (39) with $\tilde{g}^{\alpha\beta}$ we obtain

$$\tilde{S}_{\gamma\delta}^2 = (\tilde{\kappa} - \rho)\tilde{S}_{\gamma\delta} + (2\mu + \rho\tilde{\kappa})\tilde{g}_{\gamma\delta},$$

which, by making use of (42), can be rewritten in the following form

$$\tilde{S}_{\gamma\delta}^2 - \frac{1}{3} \text{tr}(\tilde{S}^2)\tilde{g}_{\gamma\delta} = \left(\frac{\tilde{\kappa}}{2} + \alpha\right) \left(\tilde{S}_{\gamma\delta} - \frac{\tilde{\kappa}}{3}\tilde{g}_{\gamma\delta}\right), \quad \alpha = (\tilde{\kappa} - \rho) - \frac{\tilde{\kappa}}{2} = \frac{\tilde{\kappa}}{2} - \rho. \quad (44)$$

From (44), in view of Lemma 2.1(i), it follows that $(\tilde{R} \cdot \tilde{S})_{\alpha\beta\gamma\delta} = \alpha Q(\tilde{g}, \tilde{S})_{\alpha\beta\gamma\delta}$ holds at $x \in \mathcal{U}_L$ and in a consequence, $(\tilde{R} \cdot \tilde{R})_{\alpha\beta\gamma\delta\epsilon\mu} = \alpha Q(\tilde{g}, \tilde{R})_{\alpha\beta\gamma\delta\epsilon\mu}$. The last relation, by (43)(a), implies $\alpha = 0$, i.e. $\rho = \tilde{\kappa}/2$. Applying the last equality and (42) in (40) we find

$$(3L - 1)\tau = \frac{5}{6}\tilde{\kappa}, \quad \frac{2}{3}\tilde{\kappa}(3L - 1)\tau = -\frac{13}{36}\tilde{\kappa}^2, \quad (45)$$

which gives $(5/6)\tilde{\kappa}^2 = -(13/36)\tilde{\kappa}^2$, whence $\tilde{\kappa} = 0$. Now (45) reduces to $(3L - 1)\tau = 0$ and in a consequence, from (40) we get $\rho = \mu = 0$. So, (39) reduces to $\text{rank}(\tilde{S}) = 1$. Since $\tilde{\kappa} = 0$, (28) and (34) leads to $\tau_1 = \tau = (1/2)(\text{tr}(T) + (\Delta_1 F/F))$. Further, we denote by \mathcal{U}_F the set consisting of all points of \mathcal{U}_L at which $F' \neq 0$. We suppose that τ vanishes at $x \in \mathcal{U}_F$. Then (43)(b) implies $F' = 0$, a contradiction. Thus $L = 1/3$ holds on \mathcal{U}_F . We note that if $L = 1/3$ then only the functions F , defined by $F(x^1) = a \exp(bx^1)$, $a = \text{const.} > 0$, $b = \text{const.} \neq 0$, are non-constant solutions of (36) and (43)(b). Thus we have the following.

THEOREM 5.1. *Let the set \mathcal{U}_F be a dense subset of the set \mathcal{U}_L of a 4-dimensional warped product $\bar{M} \times_F N$, $\dim \bar{M} = 1$. Then the warped product $\bar{M} \times_F N$ satisfies the condition $R \cdot C = LQ(S, C)$ on the set $\mathcal{U}_L \subset \mathcal{U} \subset \bar{M} \times N$ if and only if $L = 1/3$, $F(x^1) = a \exp(bx^1)$, $a = \text{const.} > 0$, $b = \text{const.} \neq 0$, and (N, \tilde{g}) is a 3-dimensional semi-Riemannian manifold fulfilling $\text{rank}(\tilde{S}) = 1$ and $\tilde{\kappa} = 0$.*

REMARK 5.1. Let (N, \tilde{g}) , $\dim N = 3$, be a semisymmetric manifold with vanishing identically on N scalar curvature $\tilde{\kappa}$. Suppose that \tilde{g} is a Riemannian

metric. Using this fact we can easily deduce that the condition $\text{rank } \tilde{S} \leq 1$ implies $\tilde{S} = 0$. Therefore, if the assumption $\text{rank } \tilde{S} = 1$ is fulfilled on (N, \tilde{g}) then the metric \tilde{g} must be necessary indefinite, more precisely, \tilde{g} is a Lorentzian metric.

Now from Theorem 5.1, in view of the above remark, follows the following

COROLLARY 5.1. *If a generalized Robertson-Walker spacetime satisfies (*) then at every point of this spacetime at least one of the tensors $R \cdot C$ or $Q(S, C)$ must vanish.*

Let x be a point of a 4-dimensional warped product $\bar{M} \times_F N$, $\dim \bar{M} = 1$. If at x the conditions: $C \neq 0$ and $R \cdot C = 0$ are satisfied then $R \cdot R = 0$ holds at x ([17], Theorem 3). If at x the conditions: $C \neq 0, S \neq 0$ and $Q(S, C) = 0$ are satisfied then $R \cdot R = (\kappa/3)Q(g, R)$ holds at x ([25], Theorem 3.1). If at x the condition $S = 0$ is satisfied then $C = 0$ holds at x . This statement is an immediate consequence of (14) and (32). Finally, if at x the condition $C = 0$ is satisfied then $R \cdot R = \alpha Q(g, R)$, $\alpha \in \mathbf{R}$, holds at x ([13], Lemma 3.1). These facts, together with Corollary 5.1, leads to the following

THEOREM 5.2. *Every generalized Robertson-Walker spacetime satisfying (*) is a pseudosymmetric manifold.*

REMARK 5.2. (i) Theorem 2 of [29] implies that the warped product $\bar{M} \times_F N$, of a 1-dimensional base manifold (\bar{M}, \bar{g}) , a warping function F and a 3-dimensional manifold (N, \tilde{g}) with the Ricci tensor \tilde{S} of rank one realizes $(*)_5$, i.e. $\bar{M} \times_F N$ is a manifold with pseudosymmetric Weyl tensor.

(ii) We can also check that the Weyl tensor of the warped product defined above is not of harmonic curvature.

References

- [1] A. Adamów, R. Deszcz, On totally umbilical submanifolds of some class of Riemannian manifolds, *Demonstratio Math.*, **16** (1983), 39–59.
- [2] L. Alias, A. Romero and M. Sánchez, Compact spacelike hypersurfaces of constant mean curvature in generalized Robertson-Walker spacetimes, in: *Geometry and Topology of Submanifolds, VII*, World Scientific, River Edge, NJ, 1995, 67–70.
- [3] K. Arslan, R. Deszcz, R. Ezentaş and M. Hotloś, On a certain subclass of conformally flat manifolds. *Bull. Inst. Math. Acad. Sinica*, **26** (1998), 283–299.
- [4] A. L. Besse, *Einstein Manifolds*, Springer-Verlag, Berlin, Heidelberg, New York, 1987.
- [5] R. L. Bishop and B. O'Neill, Manifolds of negative curvature, *Trans. Amer. Math. Soc.*, **145** (1969), 1–49.

- [6] E. Boeckx, O. Kowalski and L. Vanhecke, *Riemannian Manifolds of Conullity Two*, World Scientific, Singapore, 1996.
- [7] F. Defever and R. Deszcz, On semi-Riemannian manifolds satisfying the condition $R \cdot R = Q(S, R)$, in: *Geometry and Topology of Submanifolds, III*, World Scientific, River Edge NJ, 1991, 108–130.
- [8] F. Defever and R. Deszcz, On warped product manifolds satisfying a certain curvature condition, *Atti Accad. Peloritana Pericolanti, Cl. Sci. Fis. Mat. Nat.*, **69** (1991), 213–236.
- [9] F. Defever and R. Deszcz, On Riemannian manifolds satisfying a certain curvature condition imposed on the Weyl curvature tensor, *Acta Univ. Palacki. Olomunc., Fac. Rerum Nat., Math.*, **110(32)** (1993), 27–34.
- [10] F. Defever, R. Deszcz, P. Dhooche, L. Verstraelen and Ş. Yaprak, On Ricci-pseudosymmetric hypersurfaces in spaces of constant curvature, *Results in Math.*, **27** (1995), 227–236.
- [11] F. Defever, R. Deszcz and M. Prvanović, On warped product manifolds satisfying some curvature condition of pseudosymmetry type, *Bull. Greek Math. Soc.*, **36** (1994), 43–67.
- [12] F. Defever, R. Deszcz, L. Verstraelen and Ş. Yaprak, Pseudosymmetry type curvature properties of hypersurfaces, in: *Geometry and Topology of Submanifolds, V*, World Scientific, River Edge, NJ, 1993, 109–131.
- [13] J. Deprez, R. Deszcz and L. Verstraelen, Examples of pseudosymmetric conformally flat warped products, *Chinese J. Math.*, **17** (1989), 51–65.
- [14] R. Deszcz, On Ricci-pseudosymmetric warped products, *Demonstratio Math.*, **22** (1989), 1053–1065.
- [15] R. Deszcz, Examples of four-dimensional Riemannian manifolds satisfying some pseudosymmetry curvature conditions, in: *Geometry and Topology of Submanifolds, II*, World Scientific, Teaneck, NJ, 1990, 134–143.
- [16] R. Deszcz, On conformally flat Riemannian manifold satisfying certain curvature conditions, *Tensor, N.S.*, **49** (1990), 134–145.
- [17] R. Deszcz, On four-dimensional Riemannian warped product manifolds satisfying certain pseudosymmetry curvature conditions, *Colloquium Math.*, **62** (1991), 103–120.
- [18] R. Deszcz, On pseudosymmetric spaces, *Bull. Soc. Math. Belg.*, **44** (1992), Ser. A, Fasc. 1, 1–34.
- [19] R. Deszcz, Curvature properties of a certain compact pseudosymmetric manifold, *Colloquium Math.*, **65** (1993), 139–147.
- [20] R. Deszcz, On pseudosymmetric warped product manifolds, in: *Geometry and Topology of Submanifolds, V*, World Scientific, River Edge, NJ, 1993, 132–146.
- [21] R. Deszcz and W. Grycak, On some class of warped product manifolds, *Bull. Inst. Math. Acad. Sinica*, **15** (1987), 311–322.
- [22] R. Deszcz and W. Grycak, On manifolds satisfying some curvature conditions, *Colloquium Math.*, **57** (1989), 89–92.
- [23] R. Deszcz and W. Grycak, On certain curvature conditions on Riemannian manifolds, *Colloquium Math.*, **58** (1990), 259–268.
- [24] R. Deszcz and M. Hotłoś, Remarks on Riemannian manifolds satisfying a certain curvature condition imposed on the Ricci tensor, *Prace Nauk. Pol. Szczec.*, **11** (1989), 23–34.
- [25] R. Deszcz and M. Hotłoś, On a certain extension of the class of semisymmetric manifolds, *Publ. Inst. Math. Beograd*, **63** (77) (1998), 115–130.
- [26] R. Deszcz, P. Verheyen and L. Verstraelen, On some generalized Einstein metric conditions, *Publ. Inst. Math. Beograd*, **60** (74) (1996), 108–120.
- [27] R. Deszcz and L. Verstraelen, Hypersurfaces of semi-Riemannian conformally flat manifolds, in: *Geometry and Topology of Submanifolds, III*, World Scientific, River Edge, NJ, 1991, 131–147.
- [28] R. Deszcz, L. Verstraelen and L. Vrancken, On the symmetry of warped product spacetimes, *Gen. Rel. Grav.*, **23** (1991), 671–681.
- [29] R. Deszcz, L. Verstraelen, and Ş. Yaprak, Warped products realizing a certain condition of pseudosymmetry type imposed on the Weyl curvature tensor, *Chinese J. Math.*, **22** (1994), 139–157.

- [30] R. Deszcz and Ş. Yaprak, Curvature properties of Cartan hypersurfaces, *Colloquium Math.*, **67** (1994), 91–98.
- [31] R. Deszcz and Ş. Yaprak, Curvature properties of certain pseudosymmetric manifolds, *Publ. Mat. Debrecen*, **45** (1994), 334–345.
- [32] P. E. Ehrlich, Y.-T. Jung and S.-B. Kim, Constant scalar curvatures on warped product manifolds, *Tsukuba J. Math.*, **20** (1996), 239–256.
- [33] G. I. Kruchkovich, On semi-reducible Riemannian spaces (in Russian), *Dokl. Akad. Nauk SSSR*, **115** (1957), 862–865.
- [34] G. I. Kruchkovich, On some class of Riemannian spaces (in Russian), *Trudy sem. po vekt. i tenz. analizu*, **11** (1961), 103–128.
- [35] G. I. Kruchkovich, On spaces of V. F. Kagan (in Russian), Appendix to the book: V. F. Kagan, *Subprojective spaces*, Moscow, 1961, 163–195.
- [36] Z. I. Szabó, Structure theorems on Riemannian spaces satisfying $R(X, Y) \cdot R = 0$. I. The local version, *J. Diff. Geom.*, **17** (1982), 531–582.
- [37] L. Verstraelen, Comments on pseudo-symmetry in the sense of Ryszard Deszcz, in: *Geometry and Topology of Submanifolds, VI*, World Scientific, River Edge, NJ, 1994, 199–209.

Ryszard Deszcz
Department of Mathematics
Agricultural University of Wrocław
ul. Grunwaldzka 53, PL-50-357 Wrocław
POLAND

Marek Kucharski
Institute of Mathematics
Wrocław University of Technology
Wybrzeże Wyspiańskiego 27, PL-50-376 Wrocław
POLAND