

# Meta-analysis of factors associated with antidiabetic drug prescribing for type 2 diabetes mellitus

Fatema Mahmoud<sup>1</sup>  | Alexander Mullen<sup>1</sup> | Chris Sainsbury<sup>2</sup> |  
Gordon F. Rushworth<sup>3</sup> | Haya Yasin<sup>1</sup> | Nouf Abutheraa<sup>4</sup> |  
Tanja Mueller<sup>1</sup> | Amanj Kurdi<sup>1,5,6,7</sup> 

<sup>1</sup>Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, UK

<sup>2</sup>Department of Diabetes, NHS Greater Glasgow & Clyde, Glasgow, UK

<sup>3</sup>Highland Pharmacy Education & Research Centre, NHS Highland, Inverness, UK

<sup>4</sup>The Institute of Applied Health Sciences, University of Aberdeen, Aberdeen, UK

<sup>5</sup>Department of Pharmacology and Toxicology, College of Pharmacy, Hawler Medical University, Erbil, Iraq

<sup>6</sup>Centre of Research and Strategic Studies, Lebanese French University, Erbil, Iraq

<sup>7</sup>Division of Public Health Pharmacy and Management, School of Pharmacy, Sefako Makgatho Health Sciences University, Pretoria, South Africa

## Correspondence

Fatema Mahmoud, University of Strathclyde, Strathclyde Institute of Pharmacy & Biomedical Sciences (SIPBS), 161 Cathedral St, G4 0RE, Glasgow, UK.  
Email: [fatema.mahmoud@strath.ac.uk](mailto:fatema.mahmoud@strath.ac.uk)

## Funding information

Jordan University of Science and Technology

## Abstract

**Background:** There is a lack of consensus on prescribing alternatives to initial metformin therapy and intensification therapy for type 2 diabetes mellitus (T2DM) management. This review aimed to identify/quantify factors associated with prescribing of specific antidiabetic drug classes for T2DM.

**Methods:** Five databases (Medline/PubMed, Embase, Scopus, Web of Science) were searched using the synonyms of each concept (patients with T2DM, anti-diabetic drugs and factors influencing prescribing) in both free text and Medical Subject Heading (MeSH) forms. Quantitative observational studies evaluating factors associated with antidiabetic prescribing of metformin, sulfonylurea, thiazolidinedione, Dipeptidyl-peptidase 4 inhibitors (DPP4-I), sodium glucose transporter 2 inhibitors (SGLT2-I), Glucagon-Like peptide receptor agonist (GLP1-RA) and insulin in outpatient settings and published from January 2009 to January 2021 were included. Quality assessment was performed using a Newcastle-Ottawa scale. The validation was done for 20% of identified studies. The pooled estimate was measured using a three-level random-effect meta-analysis model based on odds ratio [95% confidence interval]. Age, sex, body mass index (BMI), glycaemic control (HbA1c) and kidney-related problems were quantified.

**Results:** Of 2331 identified studies, 40 met the selection criteria. Of which, 36 and 31 studies included sex and age, respectively, while 20 studies examined baseline BMI, HbA1c and kidney-related problems. The majority of studies (77.5%, 31/40) were rated as good and despite that the overall heterogeneity for each studied factor was more than 75%, it is mostly related to within-study variance. Older age was significantly associated with higher sulfonylurea prescription (1.51 [1.29–1.76]), yet lower prescribing of metformin (0.70 [0.60–0.82]), SGLT2-I (0.57 [0.42–0.79]) and GLP1-RA (0.52 [0.40–0.69]); while higher baseline BMI showed opposite significant results (sulfonylurea: 0.76 [0.62–0.93], metformin: 1.22 [1.08–1.37], SGLT2-I: 1.88 [1.33–2.68], and GLP1-RA: 2.35 [1.54–3.59]). Both

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *European Journal of Clinical Investigation* published by John Wiley & Sons Ltd on behalf of Stichting European Society for Clinical Investigation Journal Foundation.

higher baseline HbA1c and having kidney-related problems were significantly associated with lower metformin prescription (0.74 [0.57–0.97], 0.39 [0.25–0.61]), but more insulin prescriptions (2.41 [1.87–3.10], 1.52 [1.10–2.10]). Also, DPP4-I prescriptions were higher for patients with kidney-related problems (1.37 [1.06–1.79]) yet lower among patients with higher HbA1c (0.82 [0.68–0.99]). Sex was significantly associated with GLP1-RA and thiazolidinedione prescribing (F:M; 1.38 [1.19–1.60] and 0.91 [0.84–0.98]).

**Conclusion:** Several factors were identified as potential determinants of anti-diabetic drug prescribing. The magnitude and significance of each factor differed by antidiabetic class. Patient's age and baseline BMI had the most significant association with the choice of four out of the seven studied antidiabetic drugs followed by the baseline HbA1c and kidney-related problems which had an impact on three studied antidiabetic drugs, whereas sex had the least impact on prescribing decision as it was associated with GLP1-RA and thiazolidinedione only.

#### KEYWORDS

antidiabetic drugs, meta-analysis, prescribing, prescribing criteria, type 2 diabetes

## 1 | INTRODUCTION

Diabetes mellitus (DM) is a chronic progressive disorder characterised primarily by persistent hyperglycaemia<sup>1</sup>; according to the International Diabetes Federation, in 2021, around 537 million adults were diagnosed with DM worldwide.<sup>2</sup> More than 90% of people with DM have type 2 DM (T2DM) which is characterised by chronic hyperglycaemia and insulin resistance, contributing to the development of diabetes-related life-threatening complications.<sup>3</sup> These complications can be prevented/attenuated by achieving adequate glycaemic control following an appropriate non-pharmacological and pharmacological care plan.<sup>4,5</sup>

Several groups of antidiabetic drugs (ADDs) with different effectiveness and safety profiles are currently available. The most commonly used ADDs are metformin, sulfonylurea (SU), thiazolidinedione (TZD), dipeptidyl-peptidase-4 inhibitors (DPP4-I), sodium glucose transporter-2 inhibitors (SGLT2-I), glucagon-like peptide receptor-1 agonists (GLP1-RA) and insulins.<sup>4–6</sup> All clinical guidelines have agreed on metformin as first-line therapy for patients newly diagnosed with T2DM.<sup>7–9</sup> However, the choice of intensifying therapy or alternative initial therapy in the presence of contraindications to metformin, is more variable, and prescribing decision could be influenced by several factors relevant to patients and drugs characteristics.<sup>7–9</sup>

Several observational studies have evaluated the association of multiple factors with ADD prescribing (ADP) in clinical practice, including, for example, patient's age,

sex, ethnicity, socioeconomic status, body mass index (BMI), glycaemic control (HbA1c), renal function and history of microvascular/macrovascular complications.<sup>10–14</sup> Nevertheless, no previous studies extensively quantified the impact of these different factors on prescribing decisions of different ADDs, which would be of interest especially after the introduction of newer ADDs which provided prescribers not only with wider options for T2DM management, but with ADDs that may have independent cardiac and renal protection effects.<sup>15–17</sup>

Generally, factors associated with drug prescribing may indirectly reflect prescriber's adherence to guideline recommendations and specific drug features. This highlights the importance of studying these factors in a systematic way to assess the process of patient care by investigating which and how factors contribute to the decision-making in clinical practice.<sup>18</sup> Therefore, this systematic review (SR) and meta-analysis (MA) aimed to summarise and quantify factors associated with ADP at both the initiation and intensification stages.

## 2 | METHODS

This SR&MA is presented following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist (Appendix S1<sup>19</sup>). The protocol was registered in the international prospective register of systematic reviews (PROSPERO; registration number CRD42020173917).

## 2.1 | Data sources and search strategy

The search strategy was developed using three main concepts: participants (patients with T2DM), intervention (antidiabetic drugs) and outcome (factors associated with ADP). Medline/PubMed, Embase, Scopus and Web of Science were searched for studies published between January 2009 and January 2021 (the date of starting data synthesis). Additional searches were conducted to ensure literature saturation on ProQuest, Open Grey database and the reference lists of included articles. The search strategy was independently reviewed by experienced researchers and an academic librarian. As an example, the full Medline search strategy is available in the Appendix S2.

## 2.2 | Eligibility criteria

Only quantitative observational studies reporting factors associated with ADP among adults with T2DM in primary-care/outpatient settings and published in English were included (Table 1). Literature was searched from 2009 onwards to ensure the inclusion of newer ADDs (GLP1-RA, DPP4-I and SGLT2-I), which have been introduced into market since 2009. Only adults ( $\geq 18$  years of age) were included to ensure that all people were subject to the same

treatment recommendations since different treatments are recommended for children with T2DM.

## 2.3 | Study selection, data extraction and quality assessment

Two stages of study selection were conducted using Covidence<sup>20</sup>: title/abstract screening; and full-text screening. Relevant data was extracted from included studies using an MS Excel extraction form that was initially piloted on a random 10% of included studies to assess whether it captures all relevant data. All identified factors were mapped into four categories: 1 – demographic factors; 2 – clinical factors; 3 – socioeconomic factors; and 4 – prescriber-related factors, which were initially developed based on the literature around factors affecting physician's prescribing decision and modified as appropriate to fit the current research question.<sup>21</sup> Along with data extraction, the included studies were evaluated for the risk of bias using the Newcastle-Ottawa scale (NOS) for cohort studies<sup>22</sup> and the adapted NOS for cross-sectional studies.<sup>23,24</sup> More details on the quality assessment and study scoring are available in the Appendix S5. At each step of screening, extraction and quality assessment, a total of 20% of included studies were validated by two independent reviewers.

TABLE 1 Study inclusion and exclusion criteria.

Category	Inclusion criteria
Language	English
Publication year	January 2009–January 2021
Publication type	Any studies reporting on factors associated with antidiabetic drugs' prescription or patients' characteristics prior or at the time of antidiabetic drugs' prescription
Methodology	Quantitative observational study designs
Diabetes type	Only type 2 diabetes mellitus
Patients	Adults ( $\geq 18$ years old) patients who were prescribed any of the following: Biguanide (metformin), Sulfonylurea (SU), thiazolidinedione (TZD), dipeptidyl-peptidase 4 inhibitors (DPP4-I), sodium glucose transporter 2 inhibitors (SGLT2-I), glucagon-like peptide 1 receptor agonists (GLP1-RA), and insulins
Category	Exclusion criteria
Language	Other than English
Publication year	Published before January 2009
Publication type	Reports, commentaries, editorials, book chapters, systematic reviews and meta-analysis
Patients	Studies on children, adolescents, pregnant or breastfeeding women
Outcome	Studies including type I DM, gestational diabetes Studies did not clearly state that factors were collected at baseline Studies with data on inpatient or admitted patients No relevant outcomes (e.g. switching medicine, discontinuation) Studies did not specify the type of antidiabetic groups being studied

## 2.4 | Data synthesis and meta-analysis

All factors related to ADP were identified from included studies. However, only factors reported by more than two studies for their association with the individual antidiabetic class were eligible for MA. Accordingly, MA was applied on five of the identified factors: age, sex, HbA1c, BMI and kidney-related problems.

Three-level random-effect models were used to combine the pooled estimates (presented as odds ratio (OR) and 95% confidence intervals (CI)) measuring the association of each factor with ADP; a three-level MA approach was used to address the presence of dependency or correlation of effect sizes arising from reporting more than one effect size per study due to examining the outcomes of more than one antidiabetic group.<sup>25,26</sup> Subsequently, the pooled estimate measuring the association of the individual factor with each type of antidiabetic class was calculated using a two-level random effect model. Studies to be included in the MA had to report the effects of the identified factors as OR from either binary or continuous data; or provide primary baseline data required for OR calculation. Appendix S3 provides details on the method of effect size computation.

Study heterogeneity was measured by conducting Higgins & Thompson's ( $I^2$ ) test over three levels to compute the overall heterogeneity as well as within-study (level-2) and between-studies (level-3) variance, with  $I^2 > 75\%$  indicating high heterogeneity.<sup>27</sup> Furthermore, the usefulness and performance of the three-level model was evaluated by conducting log-likelihood-ratio test.<sup>27-29</sup>

Moderator (sub-group) analysis was performed to explore any source of heterogeneity including the potential effect of several variables related to study characteristics on the overall estimate, such as class of ADDs, stage of treatment at which the outcome was assessed (initiation, intensification or not specified stage), quality of study, type of analysis used (adjusted vs. un-adjusted), study design and year of publication.<sup>28</sup> A  $p$ -value of  $<.05$  indicates a significant result. Some factors (age, BMI, HbA1c) were reported as a binary variable in some studies and as a continuous variable in other studies. The pooled estimate of those factors was initially computed including all studies presenting the outcome as binary or continuous data following the approach described by Cochrane guideline. Additional sub-group analysis based on the type of reported data was performed to assess whether there was a significant difference in the pooled estimates according to the data type.

Publication bias was assessed with the funnel plot and extended Eggers' test.<sup>30,31</sup> Moreover, the number of

outliers was measured and plotted as histogram for each MA and a sensitivity analysis was conducted to explore the influence of outliers on the pooled estimate. An effect size was considered as an outlier when its CI does not overlap with the CI of the pooled estimate.<sup>32</sup> Cook's distance ( $D$ ) test was also performed with the results presented as scatter plots to explore the influential impact of included studies.<sup>32</sup> A Cook's- $D$  value of  $\geq 4/k$  ( $k$ : the number of effect sizes) indicated an influential impact of a study on the overall estimate.<sup>33</sup> All statistical tests were performed using RStudio; for full R-syntax, please refer to Appendix S4.

## 3 | RESULTS

### 3.1 | Study selection

Of the 2331 identified studies which had title/abstract screened, 96 underwent full text screening and 40 articles met all inclusion criteria (Figure 1). The percentage of agreement between the independent reviewers from title/abstract and full-text screening was high (93.8% and 85.7%, respectively).

### 3.2 | Study characteristics

Table 2 shows the characteristics of included studies. More than two thirds of eligible studies ( $n = 33$ , 82.5%) were published from 2013 onwards.<sup>10,13,14,35,37-39,41,43,44,46,48-67,69,70</sup> The majority ( $n = 36$ , 90%) were of cohort study design<sup>10,13,14,34-42,44,45,47-53,55-62,64-70</sup> and more than one third ( $n = 15$ ; 37.5%) originated from the United States.<sup>14,36,40,42,46,47,54,56,57,59,60,64,65,67,68</sup> The total number of participants from the included studies was 5,327,502 people with T2DM, excluding one study which provided the number of visits rather than the number of patients.<sup>46</sup> Oral antidiabetics were examined in 90% of studies ( $n = 36/40$ ) whereas injectable drugs were evaluated in about half of included studies ( $n = 21$ , 52.5%). The most frequently investigated ADDs were SU ( $n = 21$ , 52.5%), metformin ( $n = 20$ , 50%) and DPP4-I ( $n = 19$ , 47.5%) while SGLT2-I was the least studied group ( $n = 11$ , 27.5%). Only 29 (72.5%) studies stated at which stage of treatment the outcome was observed; whether at initiation ( $n = 14$ ) or intensification ( $n = 15$ ). The quality assessment score of cohort studies ranged from 5 to 9 with the majority of studies ( $n = 29/37$ , 78.4%) rated as good. Of the three cross-sectional studies, two were rated as very good and one as satisfactory (Appendix S5).




FIGURE 1 PRISMA flow chart of screening process to identify relevant studies (January 2009–January 2021).

### 3.3 | Meta-analysis results

The following factors were identified in the included studies as factors associated with ADP (Table 3): demographic factors (patients' age, sex, ethnicity, smoking status, family history of diabetes and educational level), clinical factors (obesity, glycaemic status (HbA1c), kidney function, having macrovascular/microvascular complications or other comorbidities and diabetes duration), socioeconomic factors (deprivation level, income level, employment status, having insurance, area of living and type of medical facility) and prescriber-related factors (prescriber age, sex, speciality and practice experience).

However, it was possible to perform the MA on only five factors: age, sex, HbA1c, BMI and kidney-related problems.

### 3.4 | Sex

Out of the 40 eligible studies, 36 (90%) reported on sex association with ADP and all except one<sup>50</sup> were included in

the MA, contributing to a total of 96 effect sizes. Heintjes et al.<sup>50</sup> was excluded since the outcome was not reported as OR and insufficient data was available for OR computation. The three-level MA showed that overall, sex had no association with ADP (OR [95% CI]: 1.00 [0.86–1.16]), Figures 2A, 2B. However, subgroup analysis showed that the effect of sex varied significantly by the class of ADDs ( $p = .001$ ), with the most significant influence on GLP1-RA and TZD prescription. Female patients were significantly more likely to be prescribed GLP1-RA compared to male patients (OR [95%CI]: 1.38 [1.19–1.60]), yet less likely to get TZD prescription (OR [95% CI]: 0.91 [0.84–0.98]).

### 3.5 | Age

Age was evaluated in a total of 38 studies; 31 studies were included in the MA, contributing to a total of 88 effect sizes. Seven studies were excluded since they did not present the outcome as OR and did not provide the required data for OR computation.<sup>37,40,46,50,52,58,62</sup>

TABLE 2 Characteristics of the 40 studies which were eligible for inclusion.

Author, year, country	Study design	Study duration	# Of Participants/Age/Sex	Antidiabetic drug studied	Stage of treatment	Analysis method
Winkelmayr et al. (2010), Austria <sup>34</sup>	Retrospective cohort	1/2007–6/2008	39,077 patients/19–100 years; mean 63.4 years/F: 50.4%	Metformin vs. Other oral hypoglycaemic	Initiation	Multivariable logistic regression
Abdelmoneim et al. (2013), Canada <sup>35</sup>	Retrospective cohort	January 1998–December 2010	31,421 patients/≥65 years; mean: 74.8 years/F: 49%	Metformin vs. SU monotherapy	Initiation	Multivariable logistic regression models
Brouwer et al. (2012), United States <sup>36</sup>	Retrospective cohort	January 1998–December 2009	1972 patients/>21 years; median: 54 years/F: 52.5%	Metformin vs. SU, TZD, combination. SU vs. combination	Initiation	Multinomial regression model
Liu et al. (2017), Taiwan <sup>37</sup>	Retrospective cohort	January 2006–December 2010	28,640 patients/≥20 years; mean: 57.4/F: 47.3%	Non-metformin vs. metformin	Initiation	Logistic regression
Wang et al. (2013), Quebec, Canada <sup>38</sup>	Dynamic historical cohort study	January 2003–December 2011	1279 patients/≥18 years, 53.4% ≥65 years/F: 50.8%	Metformin vs. non metformin	Initiation	Multivariate generalised estimating equation analysis
Geier et al. (2014), Germany <sup>33</sup>	Retrospective cohort	June 2003–December 2009	10,657 patients/40–79 years; mean for both: 61.47 years/F: 51%	Metformin vs. SU	Initiation	Multiple logistic regression
Fujihara et al. (2017), Japan <sup>39</sup>	Retrospective cohort	December 2009–March 2015	2666 patients/≥20 years; overall mean: 60.9 years/F: 35.9%	Metformin, DPP-4Is vs. SU	Initiation	Multinomial logistic regression analysis
Desai et al. (2012), United States <sup>40</sup>	Retrospective cohort	January 2006–December 2008	254,973 patients/18–100 years; mean: 58.2 years/F: 47.3%	Metformin	Initiation	Multivariable logistic models
Grimes et al. (2015), Ireland <sup>41</sup>	Retrospective cohort	January 2008–December 2009	20,947 incident users of antidiabetic agents/≥40 years/F: 42.1%	Metformin vs. SU	Initiation	Adjusted logistic regression
Cai et al. (2010), United States <sup>42</sup>	Retrospective cohort	January 2006–June 2008	240,426 patients/26–88 years; mean: 54.4/F: 44.8%	Sitagliptin vs. non-Sitagliptin	Not specified	Chi-square statistics
Saine et al. (2015), United States and United Kingdom <sup>43</sup>	Cross-sectional/	THIN: October 2009–September 2012, US Medicare: August 2009–December 2011, HIRD: August 2009–July 2012	UK: 43,466, US: 631,273/Mean: UK: 58.8, US: 67.6 years/F: UK: 42.4%, US: 55%	Saxagliptin vs. other OAD	Not specified	Conditional logistic regression
Wilkinson et al. (2018), United Kingdom <sup>10</sup>	Retrospective cohort	January 2014–July 2017	14,149 individuals/≥18 years; Mean: 60 years/F: 40.3%	SGLT2-I, DPP 4-I vs. SU	1st intensification	Multinomial logistic regression, Sensitivity analyses
Grabner et al. (2015), United States <sup>14</sup>	Retrospective cohort	January 2011–September 2013	Overall: 27790 patients/≥18 years; mean overall: 55.03 years/F: 39.4%	Canagliflozine vs. DPP 4-I	Intensification not specified level of intensification	Multivariable logistic regression
OU et al. (2016), Taiwan <sup>44</sup>	Retrospective cohort	2011–2012	32,724 patients/≥20 years	DPP-4 I vs. other antidiabetics	Intensification, 1st and 2nd	Multiple logistic regression
Stargardt et al. (2009), Finland, France, Germany, Norway, Poland, Spain, UK <sup>45</sup>	Retrospective cohort	June 2006 and February 2007	1218 patients: 891; added SU 327 added TZD/≥30 years; Mean: SU, TZD: 61.0, 57.8 years	TZD vs. SU	1st intensification	Probit regression analysis

TABLE 2 (Continued)

Author, year, country	Study design	Study duration	# Of Participants/Age/Sex	Antidiabetic drug studied	Stage of treatment	Analysis method
Payk et al. (2015), United States <sup>46</sup>	Cross-sectional study	2003 to 2004 and 2007 to 2010	7042 visits (≥18 years; Mean: 61.6 years/F: 52%)	SU	Not specified	Weighted sampling, A multivariate logistic regression model
Zhang et al. (2010), United States <sup>47</sup>	Retrospective cohort	October 2006 – June 2008	41,836 patients/≥30 years; Mean overall: 60.08	Sitagliptin vs. Non-Sitagliptin	Not specified	Adjusted logistic regression analysis
Morita et al. (2019), Japan <sup>48</sup>	Retrospective cohort	October 2012–September 2016	Metformin and DPP4i users only: 74935,658	DPP4i vs. metformin	Initiation	Univariate logistic regression
Kim et al. (2019), Korea <sup>49</sup>	Retrospective cohort	2014–2016	3609 patients/≥20 years/F:48.9%	Newer (SGLT2i and DPP4i) vs. older (SU and TZD)	1st Intensification	Logistic regression analysis
Heintjes et al. (2017), Netherlands, Italy, Spain, United Kingdom <sup>50</sup>	Retrospective cohort	5 years: 2007 to 2011 (ES, IT, and NL) or 2008 to 2012 (UK)	485,570 patients/≥18 years	Metformin, SU, TZD, DPP4i, GLP-1ra, insulin.	Initiation and intensification; 1st line, 2nd line, 3rd line, 4th line	Poisson Regression
Nicolucci et al. (2019), 38 Countries <sup>51</sup>	Prospective cohort	December 2014–June 2016	14,668 patients/≥18 years; mean: overall: 57.5 years/F: 46.1%	DPP4i, SGLT2-I, GLP1-RA vs. SU	1st intensification (second line)	Firth logistic regression
Hartmann et al. (2019), Germany, Austria, Switzerland, Luxembourg <sup>52</sup>	Retrospective cohort	2000–2017	4770 patients/>18 years; median: BOT initiated, OAD w/o insulin, GLP w/o insulin: 64.0, 62.6, 55.3 years/F: overall:45.7%	GLP1-RA, basal insulin, OAD	1st Intensification	Multivariable linear and logistic regression models
Longato et al. (2020), Italy <sup>53</sup>	Retrospective cohort		12,996 patients/calculated mean for both groups: 62.84 years/F: 36.8%	SGLT2-I vs. GLP1-RA	Not specified	Chi-square, standardised mean difference, or many Whitney test
Ackermann et al. (2017), United States <sup>54</sup>	Multiple case-comparative study design	January 2011–June 2015	77,744 patients/≥18 years, 25.7% ≥65 years/F: 43.1%	DPP4-I, GLP1-RA, SGLT2-I, SU, TZD	1st intensification	Multinomial logistic regression
Whyte et al. (2019), England <sup>55</sup>	Retrospective cohort	January 2012–December 2016	49,380 patients/≥18 years; Mean: 68.7 years/F: 43.9%	SGLT2-I, GLP1-RA, metformin, insulin, SU, DPP-4-I, TZD	Not specified	Logistic regression, mixed effects model
Arnold et al. (2018), United States <sup>56</sup>	Retrospective cohort	Not stated	157,551 patients/≥18 years; Mean: 68.1 years/F:42.8	SGLT2-I, GLP1-RA, metformin, insulin, SU, DPP-4-I, TZD	Not specified	Poisson regression
Arnold et al. (2018), United States <sup>57</sup>	Retrospective cohort	2013–2016	456,106 patients/Mean: 67.6 years	SGLT2-I, GLP1-RA, metformin, insulin, SU, DPP-4-I, TZD	Not specified	Poisson regression

(Continues)

TABLE 2 (Continued)

Author, year, country	Study design	Study duration	# Of Participants/Age/Sex	Antidiabetic drug studied	Stage of treatment	Analysis method
Zaharan et al. (2013), Ireland <sup>58</sup>	Retrospective cohort	2008–2012	524,305	Metformin, SU, TZD, GLP1-RA, DPP4-I	Not specified	Adjusted logistic regression
Zoberi et al. (2017), United States <sup>59</sup>	Retrospective cohort	July 2008–July 2013	Patients: Overall: 1952/≥18 years Mean:59.3 years. F: 60.76%	Metformin, Insulin	Not specified	Chi-square and independent sample t-tests
Montvida et al. (2017), United States <sup>60</sup>	Retrospective cohort	2005–2016	1st line: 1,023,340 patients 2nd line: 357482 patients	GLP-1RA, SGLT2-I, MET, INS, TZD, DPP-4I, SU	Initiation and intensification	Descriptive only
Katakami et al. (2020), Japan <sup>61</sup>	Prospective cohort	September 2014–December 2015	1806 patients/mean:61.7 years/F:38.4%	Metformin, SU, TZD, gliitides, DPP4I, SGLT2-I, GLP1-RA, insulin	Intensification	Firth logistic regression models
Kostev et al. (2014), Germany <sup>62</sup>	Retrospective cohort	January 2003–December 2012	10, 223 patients/>40 years; Mean for both groups: 65.69 years/F for both groups: 49.7%	Insulin	Initiation intensification	A multivariate Cox regression model for insulin
Dhanaraj et al. (2013), India <sup>63</sup>	Cross-sectional	June 2007–March 2009	1185 patients/mean age: 55 years/F: 49%	Metformin, SU, insulin, pioglitazone	Not specified	Univariate logistic regression
Yu et al. (2017), United States <sup>64</sup>	Retrospective cohort	November 2014–February 2016	11,053 patients/≥18 years; Mean overall: 61.26 years/F: 51.5%	GLP-1-RA vs. basal insulin	Intensification; Not specified intensification level	Boosted regression models (GBM), logistic regression model
Levin et al. (2014), United States <sup>65</sup>	Retrospective cohort	January 2000–March 2011	51,771 patients/≥18 years; Mean overall: 55.6 years/F:40.2%	Insulin, GLP1-R vs. 3OHA	Intensification; Not specified intensification level	t-test or chi square
Gentile et al. (2018), Italy <sup>66</sup>	Retrospective cohort	2004–2011	Sample size included in the model 4 (n = 44,611)/≥18 years; mean: Overall: 65 years/F: 44.2%	Insulin vs. Non-insulin	Intensification; Not specified level of intensification	Cox proportional hazard model
Korytkowski et al. (2014), United States <sup>67</sup>	Retrospective cohort	June 2005–November 2011	1892 patients/calculated mean age for all groups: 54.78 years/F: 48.95%	Insulin, GLP1-RA vs. OHA	Intensification	t-test or chi square
Hirsch et al. (2011), United States <sup>68</sup>	Retrospective cohort	October 1, 2005, – January 2008	190, 444 patients; Sample size for multivariable model 51,048/≥18 years; mean: 62.4 years/F:51.9%	Exenatide vs. Non-exenatide	Not specified	Cox proportional hazard regression
Boom et al. (2020), Germany <sup>69</sup>	Retrospective cohort	January 2014–December 2018	10,497 patients/18-90 years, mean overall:63.6 years/F overall: 45.9%	Insulin vs. oral antidiabetic	Initiation	Multivariable logistic regression model
Moreno Juste et al. (2019), Italy <sup>70</sup>	Retrospective cohort	January 1 and December 31, 2016	12,753 patients/mean age for all groups: 63.9 years/F: 45.76%	Metformin, SU, DPP4-I	Initiation	t-test or chi square



TABLE 3 Description of the outcome of the 40 studies which were eligible for inclusion.

Author	Outcome definition	Patients-related	Clinical-related	Socioeconomic	Prescriber-related	Reported as
Winkelmayr et al. <sup>34</sup>	Associations of metformin initiation vs. any other oral hypoglycaemic medication.	Patients age, sex		Socioeconomic status	Age, sex, speciality	OR, 95% CI
Abdelmonem et al. <sup>35</sup>	Predictors of new monotherapy users	Age, sex	Microvascular (neuropathy, retinopathy, nephropathy) and macrovascular complications (as ischemic heart disease (IHD), cerebrovascular disease, heart failure (hf), peripheral vascular disease (PVD)), comorbidities (as hypertension (HTN) and dyslipidaemia)			OR, 95% CI
Brouwer et al. <sup>36</sup>	Factors influencing selection of initial oral hypoglycaemic medication	Age, sex, race	Glycaemic status (HbA1c), Serum creatinine			Probability ratio, 95% CI
Liu et al. <sup>37</sup>	Factors associated with non-metformin prescription as initial antidiabetic therapy.	Sex, age		Income level, medical facility features; accreditation level, ownership, location	Age, sex, speciality	OR, 95% CI
Wang et al. <sup>38</sup>	Predictors of starting metformin and the influence of guideline adherence on starting of oral hypoglycaemic agents	Age, sex	Comorbidities (renal and cardiovascular disease (CVD))		Sex, practice experience	OR, 95% CI
Longato et al. <sup>53</sup>	Difference in the baseline characteristics between patients newly initiated SGLT2-I vs. GLP1-RA	Age, sex	HTN, dyslipidaemia, microvascular and macrovascular complications, comorbidities			%/mean, <i>p</i> value
Geier et al. <sup>13</sup>	Predictors of Metformin vs. SU initiators	Age, sex, smoking status	Obesity, glycaemic status (HbA1c), diabetes duration			OR, 95% CI
Fujihara et al. <sup>39</sup>	Factors that influence the choice of each of three hypoglycaemic agents prescribed as initial monotherapy	Age, sex	Diabetes duration, body mass index (BMI), HTN, HbA1c			OR, 95% CI
Desai et al. <sup>40</sup>	Predictors of receiving metformin as initial oral hypoglycaemic therapy	Age, sex	Comorbidity	Income, Drug insurance cover		OR, 95% CI
Grimes et al. <sup>41</sup>	Socio-demographic factors association with initiation of metformin or SU	Age, sex				OR, 95% CI
Cai et al. <sup>42</sup>	Characteristics of patients prescribed Sitagliptin vs. other oral antidiabetics	Age, sex	Retinopathy, neuropathy, nephropathy, CVD (as HF, stroke, myocardial infarction (MI), PVD), other comorbid diseases (HTN), obesity			%, <i>p</i> value
Saine et al. <sup>43</sup>	Determinants of saxagliptin use	Age, sex, Smoking	HbA1c, obesity, nephropathy, neuropathy, retinopathy, CVD, PVD, cerebrovascular disease			OR, 95% CI/N, % for age and sex


(Continues)

TABLE 3 (Continued)

Author	Outcome definition	Patients-related	Clinical-related	Socioeconomic	Prescriber-related	Reported as
Wilkinson et al. <sup>10</sup>	Patient characteristics associated with the class of antidiabetic drug prescribed	Age, sex, Ethnicity, smoking	HbA1c, BMI, Kidney function (eGFR), CVD, retinopathy	SES		OR, 95% CI
Grabner et al. <sup>14</sup>	Baseline Characteristics with Initiation of Canagliflozin vs. DPP4-I		HbA1c, microvascular complications, dyslipidaemia, Obesity			OR, 95% CI
OU et al. <sup>44</sup>	Factors associated with the choice of DPP4-I rather than other antidiabetics	Age, sex	Comorbidities (HTN, dyslipidaemia, stroke, coronary artery disease (CAD), heart failure (HF))			Estimates, SE
Stargardt et al. <sup>45</sup>	Predicted probabilities of adding glitazone or sulfonylurea to metformin	Age, history of diabetes in family	History of macrovascular complication, HbA1c, weight		Specialty, years of experience	Predicted probability, <i>p</i> value
Payk et al. <sup>46</sup>	Predictors of SU use	Age, sex, race		Payment type	Specialty	OR, 95% CI
Zhang et al. <sup>47</sup>	Baseline characteristics of initiating Sitagliptin monotherapy compared to non-Sitagliptin monotherapy	Age	HbA1c, obesity, Microvascular conditions, Chronic renal disease, CVD			OR, 95% CI
Morita et al. <sup>48</sup>	Patient characteristics associated with the selection of DPP4-I vs. metformin	Age, sex	Renal disease, HbA1c, obesity, microvascular complications, CAD, and stroke			OR, 95% CI
Kim et al. <sup>49</sup>	The influencing factors in the selection of second oral antidiabetics added to metformin	Age, sex	CVD, renal failure, HF, dyslipidaemia	Insurance, institution	Specialty	OR, 95% CI
Heinjes et al. <sup>50</sup>	Factors associated with the choice of treatment at intensification	Age, sex, smoking status	Macrovascular complication, renal function, HbA1c, obesity			RR, 95% CI
Nicolucci et al. <sup>51</sup>	Factors associated with second-line treatment choices in patients prescribed metformin	Age, sex, Education, Health	Obesity, microvascular/macrovacular complications, diabetes duration, HbA1c, chronic kidney disease	Insurance coverage, employment status	Physician speciality	OR, 95% CI
Hartmann et al. <sup>52</sup>	Predictors of treatment escalation after metformin monotherapy failure	Age, sex	HbA1c, diabetes duration, microvascular/macrovacular disease, obesity			OR, 95% CI
Ackermann et al. <sup>54</sup>	Correlates of type 2 diabetes second line medication selection	Age, sex, race/ethnicity	HbA1c, obesity	Insurance	Specialty	Probability%, 95% C.I
Whyte et al. <sup>55</sup>	Disparity exists in drug prescribing	Ethnicity, sex		Socioeconomic status		OR, 95% CI
Arnold et al. <sup>56</sup>	The association of the variable of interest with the likelihood of being prescribed a glucose-lowering medication	Age	Obesity, Kidney function, CAD			Relative Risk/ 5years, 95% CI
Arnold et al. <sup>57</sup>	Glucose-Lowering Medication Use in T2D and HF		Heart failure			Relative risk, 95% CI
Zaharan et al. <sup>58</sup>	Variations in the prescribing of oral antidiabetic therapies	Age, sex				OR, 95% CI

TABLE 3 (Continued)

Author	Outcome definition	Patients-related	Clinical-related	Socioeconomic	Prescriber-related	Reported as
Zoberi et al. <sup>59</sup>	Characteristics of patients with diabetes by non-metformin prescription and by insulin prescription	Age, sex, Race, Smoking	HbA1c, obesity, Microvascular complications, CVD, Cerebrovascular disease, comorbidities (Hypertlipidaemia, HTN)			%, <i>p</i> value
Montvida et al. <sup>60</sup>	Patient characteristics according to antidiabetic therapy prescribed	Age, sex, ethnicity	HbA1c, obesity, CVD, chronic kidney disease			<i>N</i> , %
Katakami et al. <sup>61</sup>	Factors associated with the selection of second-line treatment	Age, sex	HbA1c, obesity, renal function (eGFR), CVD			OR, 95% CI
Kostev et al. <sup>62</sup>	Predictors of Insulin Initiation in Metformin and Sulfonylurea Users	Age, sex	Kidney function (eGFR), comorbidities (HTN, stroke, HF, Hyperlipidaemia)		Diabetologist care	HR, 95% CI; insulin
Dhanaraj et al. <sup>63</sup>	Choice of antidiabetic drug therapy and influencing factors	Age, sex, family history of diabetes	Obesity, HbA1c, microvascular complications, comorbidities, diabetes duration, serum creatinine.			OR, 95% CI
Yu et al. <sup>64</sup>	Factors that may predict choice of first injectable therapy.	Age, sex, ethnicity, smoking status	HbA1c, obesity, cardiovascular disease, chronic kidney disease, dyslipidaemia, hypertension			OR, 95% CI %, <i>p</i> value for some variables
Levin et al. <sup>65</sup>	Baseline Characteristics of patients with T2DM who added OAD, insulin, or GLP-1	Age, sex	HbA1c, HTN, Dyslipidaemia, HF, Microvascular complications (neuropathy, nephropathy, retinopathy), MI, PVD.			%, <i>p</i> value
Gentile et al. <sup>66</sup>	Predictors of initiating insulin therapy	Age, sex	Diabetes duration, HbA1c, obesity, retinopathy, kidney function (eGFR)			HR, <i>p</i> value
Korytkowski et al. <sup>67</sup>	Baseline characteristics of T2DM patients according to intensifying drugs	Sex, age, race	Obesity, HbA1c			Mean or %, <i>p</i> value
Hirsch et al. <sup>68</sup>	Predictors of exenatide use	Age, sex	Obesity, HbA1c, diabetes duration	Payer type		HR, 95% CI
Boom et al. <sup>69</sup>	Factors associated with the probability of receiving insulin	Age, sex	HbA1c, PAD, stroke, MI		Practice specialty	OR, 95% CI
Moreno Juste et al. <sup>70</sup>	Difference in the baseline characteristics among antidiabetics new users	Age, sex	microvascular/macrovascular complications	Area of living		Mean or %, <i>p</i> value




**FIGURE 2A** Forest plot of sex (female to male) association with antidiabetic drugs prescribing as overall and per antidiabetic group. CI, confidence interval; DPP4i, dipeptidyl peptidase 4 inhibitor ( $N=373,992$ ); GLP1-RA, glucagon-like peptide receptor agonist ( $N=107,128$ ); OR, odds ratio; insulin ( $N=390,711$ ).  $N$  represent all relevant studies except Whyte et al., Nicolucci et al. and Zaharan et al.

Despite that age showed an overall non-significant association with ADP (OR [95%CI]: 0.93 [0.66–1.32]), its effect varied significantly by class of ADDs ( $p < .0001$ ). For instance, SU was 51% more likely to be prescribed for older patients (reported either as continuous or categorical) (OR [95%CI]: 1.51 [1.30–1.76]). Contrastingly, patients at older age were significantly less likely to be prescribed GLP1-RA (OR [95%CI]: 0.52 [0.40–0.69]), SGLT2-I (OR

[95%CI]: 0.57 [0.42–0.79]) and metformin (OR [95%CI]: 0.70 [0.61–0.82]), **Figures 3A, 3B**.

### 3.6 | Baseline BMI


The influence of BMI on ADP was evaluated in 21 studies. All except one<sup>50</sup> were included in the MA,



**FIGURE 2B** Continued on Figure 2A for the remaining antidiabetic groups and overall estimate. Metformin ( $N = 1,186,718$ ); SGLT2-I, sodium glucose transporter 2 inhibitor ( $N = 51,874$ ); SU, sulfonyleurea ( $N = 602,435$ ); TZD, thiazolidinedione ( $N = 147,908$ ).  $N$  represent all relevant studies except Whyte et al., Nicolucci et al. and Zaharan et al.

contributing to a total of 66 effect sizes. Heintjes et al.<sup>50</sup> was excluded for the same reason stated previously. Despite that the overall MA showed no significant association of BMI with ADP (OR [95%CI]: 1.19 [0.85–1.67]),

the result varied significantly by the investigated antidiabetic group ( $p < .0001$ ). Figures 4A, 4B show that patients with higher BMI (reported either as a continuous or categorical) were more likely to be prescribed




**FIGURE 3A** Forest plot of age association with antidiabetic drugs prescribing as overall and per antidiabetic group. CI, confidence interval; DPP4i, dipeptidyl peptidase 4 inhibitor ( $N = 433,252$ ); GLP1-RA, glucagon-like peptide receptor agonist ( $N = 104,870$ ); insulin ( $N = 389,854$ ); OR, odds ratio.  $N$  represent all relevant studies except Arnold et al. and Nicolucci et al.

GLP1-RA, SGLT2-I, and metformin (OR [95%CI]: 2.35 [1.54–3.59], 1.89 [1.33–2.68], and 1.22 [1.08–1.37], respectively), but they were 24% less likely to use SU (OR [95%CI]: 0.76 [0.62–0.93]).

### 3.7 | Baseline glycaemic status (HbA1c)

A total of 62 effect sizes from 22 studies were included in the MA of HbA1c. Two studies were not included because of insufficient baseline data needed for OR

calculation.<sup>50,62</sup> Including all studies reporting HbA1c as continuous or binary, higher HbA1c value or category had no significant association with ADP (OR [95%CI]: 1.10 [0.81–1.49]). However, a significant difference in the pooled estimate per antidiabetic class was observed ( $p = .029$ ). People with higher baseline HbA1C were 2.41 times more likely to be prescribed insulin (OR [95%CI]: 2.41 [1.87–3.10]) yet less likely to get prescriptions of metformin, TZD and DPP4-I (OR [95%CI]: 0.74 [0.57–0.97], 0.76 [0.59–0.98], and 0.82 [0.68–0.99], respectively), Figures 5A, 5B.




**FIGURE 3B** Continued on Figure 3A for the remaining antidiabetic groups and overall estimate. Metformin ( $N=903,101$ ); SGLT2-I, sodium glucose transporter 2 inhibitor ( $N=51,874$ ); SU, sulfonylurea ( $N=602,435$ ); TZD, thiazolidinedione ( $N=147,908$ ).  $N$  represent all relevant studies except Arnold et al. and Nicolucci et al.

### 3.8 | Kidney-related problems

A total of 21 studies examined the impact of kidney-related problems in terms of chronic renal disease (CRD),

nephropathy or based on the estimated glomerular filtration rate (eGFR) of  $<60 \text{ mL/min/1.73 m}^2$ . Only one study<sup>50</sup> was excluded due to insufficient data necessary for OR calculation, thus 20 studies were included in the MA,



**FIGURE 4A** Forest plot of body mass index (BMI) association with antidiabetic drugs prescribing as overall and per antidiabetic groups. CI, confidence interval; DPP4i, dipeptidyl peptidase 4 inhibitor ( $N = 393,433$ ); GLP1-RA, glucagon-like peptide receptor agonist ( $N = 97,122$ ); insulin ( $N = 380,937$ ); OR, odds ratio.  $N$  represent all relevant studies except Arnold et al. and Nicolucci et al.

contributing to a total of 61 effect sizes. Of the included studies, nine reported the outcome as CRD ( $k = 28$ ) and six reported it as nephropathy ( $k = 14$ ) while the remaining five studies examined the renal function based on eGFR value ( $k = 19$ ).

The three-level MA showed that overall, kidney-related problems (either CRD, nephropathy or eGFR  $< 60$ ) had no significant association with ADP (OR [95%CI]: 0.89

[0.54–1.47]). Despite that the subgroup analysis showed a non-significant difference by class of ADDs ( $p = .079$ ); patients with kidney-related problems were significantly more likely to receive insulin (OR [95%CI]: 1.52 [1.10–2.10]) and DPP4-I (OR [95%CI]: 1.37 [1.06–1.79]) yet 61% less likely to get metformin prescriptions (OR [95%CI]: 0.39 [0.25–0.61]), **Figures 6A, 6B**. Additionally, a non-significant difference was observed in the overall estimates according







FIGURE 4B Continued on Figure 4A for the remaining antidiabetic groups and overall estimate. Metformin (N=821,416); SGLT2-I, sodium glucose transporter 2 inhibitor (N=46,007); SU, sulfonylurea (N=586,566); TZD, thiazolidinedione (N=117,355). N represent all relevant studies except Arnold et al. and Nicolucci et al.

to the type of kidney-related problems ( $p = .286$ ). The overall estimate of studies defining kidney problems as CRD was 0.64 [0.32–1.28], while the pooled estimates of studies defined kidney problems as nephropathy and eGFR were 1.13 [0.66–1.96] and 1.33 [0.55–3.12], respectively.

### 3.8.1 | Heterogeneity and three-level model fitness

Despite that overall heterogeneity was high for all studied factors (>75%), most of the total heterogeneity was



**FIGURE 5A** Forest plot of HbA1c association with antidiabetic drugs prescribing as overall and per antidiabetic groups. CI, confidence interval; DPP4-I, dipeptidyl peptidase 4 inhibitor ( $N = 323,684$ ); GLP1-RA, glucagon-like peptide receptor agonist ( $N = 95,944$ ); insulin ( $N = 380,360$ ); OR, odds ratio.  $N$  represent all relevant studies except Nicolucci et al.

related to within-study variance while between-study variance for all studied factors were  $< 75\%$  (Appendix S6). The results of the log likelihood ratio test (Appendix S6)

indicated that the three-level model had a better fit for variability in data and better estimation of the pooled estimate.





FIGURE 5B Continued on Figure 5A for the remaining antidiabetic groups and overall estimate. Metformin (N=821,416); SGLT2-I, sodium glucose transporter 2 inhibitor (N=44,396); SU, sulfonylurea (N=561,471); TZD, thiazolidinedione (N=144,871). N represent all relevant studies except Nicolucci et al.

### 3.8.2 | Moderator analysis

Tables 4 and 5 display the results of moderator analyses of all tested variables and the overall estimate within the levels of each variable for all synthesised factors. Of

all examined variables, only the type of statistical model used to assess the outcome (adjusted vs. un-adjusted) had a significant influence on the pooled estimate resulting from the MA of sex, age and kidney-related problems (p < .0001). On the other hand, there was no



**FIGURE 6A** Forest plot of kidney problem association with antidiabetic drugs prescribing as overall and per antidiabetic groups. CI, confidence interval; DPP4i, dipeptidyl peptidase 4 inhibitor ( $N=378,170$ ); GLP1-RA, glucagon-like peptide receptor agonist ( $N=91,916$ ); insulin ( $N=383,203$ ); OR, odds ratio.  $N$  represent all relevant studies except Arnold et al.


significant difference in the pooled estimate by country of study, stage of treatment and the other studied variables.

### 3.8.3 | Outliers/influential studies

A total of 15 out of 96 effect sizes of sex data, 27 out of 88 effect sizes of age data, 18 out of 66 effect sizes of BMI data and 12 out of 61 effect sizes of renal data were detected as outliers; moreover, about half of the effect sizes of HbA1c data were detected as outliers (31/62). Histogram plots of

all factors (Appendix S7) reflect that the potential outliers are not uniformly distributed around the pooled estimate. However, the results of the sensitivity analysis (Table 6) revealed a close overall OR and narrower but overlapped 95%CI of the pooled estimate after excluding the outliers compared to the one including the outliers. Nevertheless, it could not be determined whether the outliers did, in fact, bias the pooled estimate.

Cook's-D was measured for all factors (scatterplots in Appendix S7). None of the effect sizes included in the sex MA had a Cook's value exceeding 0.04(4/96), indicating that none had an influential effect on the pooled estimate.



**FIGURE 6B** Continued on Figure 6A for the remaining antidiabetic groups and overall estimate. Metformin (N=837,850); SGLT2-I, sodium glucose transporter 2 inhibitor (N=46,914); SU, sulfonylurea (N= 549,998); TZD, thiazolidinedione (N=143,382). N represent all relevant studies except Arnold et al. and Nicolucci et al.

In contrast, two effect sizes of age and HbA1c were considered as influential cases in the model as they have a distance value of >0.05(4/88) and >0.06(4/62), respectively.<sup>59,60,63,68</sup> For BMI MA, only one study presented a distance value larger than 0.06(4/66).<sup>68</sup> Lastly, three effect sizes included in the MA of kidney-related problems were considered to have influential effect in the model with a distance value of >0.07(4/61).<sup>49,61</sup>

### 3.8.4 | Publication bias

The funnel plots (Appendix S8) of all factors showed that all studies cluster at the top part of the plots, suggesting a possible presence of publication bias. Extended Eggers' test showed a significant possibility of asymmetry in the funnel plots of age, BMI and kidney-related problems (p<.0001, .0013 and <.0001, respectively), while the test

TABLE 4 Results of the moderator analysis of tested variables on the pooled estimate of each quantified factors.

Tested variable	Sex	Age	BMI	HbA1C	Kidney problem
Type of outcome variable					
Continuous	–	29/0.89 [0.72–1.10]	14/0.99 [0.67–1.45]	28/1.05 [0.59–1.87]	–
Binary		59/0.89 [0.62–1.28]	52/1.33 [0.92–1.94]	34/1.12 [0.71–1.74]	
		$p = .713$	$p = .115$	$p = .812$	
Type of analysis test					
Unadjusted	58/1.06 [0.86–1.31]	70/0.86 [0.76–1.27]	45/1.30 [0.88–1.93]	42/1.13 [0.77–1.64]	50/0.95 [0.59–1.52]
Adjusted	38/0.97 [0.86–1.20]	18/0.85 [0.64–1.13]	21/1.04 [0.83–1.31]	20/1.10 [0.87–1.40]	11/0.81 [0.36–1.86]
	$p < .0001$	$p < .0001$	$p = .518$	$p = .378$	$p < .0001$
Stage of treatment					
Initiation	30/0.98 [0.79–1.22]	28/1.16 [0.66–2.04]	15/0.93 [0.61–1.42]	16/0.87 [0.57–1.34]	11/1.35 [0.48–3.78]
Intensification	42/1.02 [0.82–1.27]	40/0.85 [0.58–1.25]	31/1.02 [0.75–1.38]	36/1.13 [0.84–1.51]	31/0.87 [0.43–1.75]
Not specified stage	24/1.00 [0.90–1.12]	20/1.04 [0.75–1.45]	20/1.57 [1.02–2.41]	10/1.21 [0.57–2.57]	19/1.05 [0.65–1.70]
	$p = .520$	$p = .415$	$p = .073$	$p = .179$	$p = .959$
Study design					
Retrospective cohort	70/0.99 [0.85–1.16]	63/0.98 [0.60–1.59]	41/1.17 [0.79–1.74]	39/1.13 [0.82–1.57]	42/0.97 [0.55–1.71]
Prospective cohort	11/0.97 [0.83–1.13]	11/1.04 [0.76–1.42]	11/1.10 [0.80–1.52]	11/1.00 [0.52–1.91]	11/0.74 [0.33–1.66]
Cross-sectional	9/0.99 [0.79–1.13]	8/1.03 [0.98–1.09]	8/1.14 [0.74–1.76]	6/1.00 [0.58–1.71]	8/1.06 [0.58–1.92]
Comparative multiple case	6 <sup>a</sup> /1.05 [0.78–1.41]	6 <sup>a</sup> /0.87 [0.63–1.20]	6 <sup>a</sup> /1.03 [0.78–1.37]	6 <sup>a</sup> /0.95 [0.62–1.47]	–
	$p = .9684$	$p = .902$	$p = .799$	$p = .844$	$p = .719$
Country					
United States	35/0.95 [0.76–1.20]	40/0.92 [0.56–1.50]	38/1.25 [0.81–1.93]	32/1.18 [0.78–1.80]	29/1.43 [0.72–2.85]
United Kingdom	16/1.06 [0.91–1.24]	5/0.87 [0.33–2.35]	3 <sup>c</sup> /1.69 [0.08–34.57]	3 <sup>c</sup> /0.87 [0.17–4.61]	3 <sup>c</sup> /0.53 [0.12–2.38]
Cross-national	11/0.97 [0.82–1.14]	9/0.89 [0.60–1.30]	8/1.49 [0.89–2.51]	9/1.39 [0.55–3.52]	10/0.80 [0.38–1.65]
Austria	4 <sup>b</sup> /1.03 [0.85–1.26]	4 <sup>b</sup> /0.89 [0.53–2.23]	–	–	–
Canada	2/0.94 [0.60–1.46]	2/0.81 [0.52–1.25]	–	–	2/0.24 [0.08–0.68]
Germany	2/1.10 [0.17–7.20]	2/1.39 [0.68–2.84]	1/1.09 [1.08–1.11]	2/1.55 [0.54–4.44]	–
Taiwan	3/1.07 [0.96–1.20]	2 <sup>c</sup> /0.87 [0.50–1.54]	–	–	–
Italy	5/0.07 [0.81–1.16]	5/1.18 [0.69–2.01]	1/0.70 [0.68–0.71]	1 <sup>g</sup> /2.61 [2.56–2.65]	2/0.94 [0.33–2.69]
Japan	10/1.05 [0.88–1.26]	11/0.95 [0.64–1.40]	11/1.16 [0.66–2.06]	11/1.05 [0.72–1.54]	7 <sup>d</sup> /1.05 [0.61–1.80]

TABLE 4 (Continued)

Tested variable	Sex	Age	BMI	HbA1C	Kidney problem
Korea	4 <sup>e</sup> /1.08 [0.67–1.76]	4 <sup>f</sup> /0.86 [0.42–1.75]	–	–	4 <sup>e</sup> /0.93 [0.24–3.59]
India	4 <sup>f</sup> /1.17 [0.85–1.60]	4 <sup>f</sup> /0.10 [0.99–1.01]	4 <sup>f</sup> /1.07 [0.75–1.53]	4 <sup>f</sup> /0.94 [0.40–2.22]	4 <sup>f</sup> /0.98 [0.438–2.18]
	<i>p</i> = .079	<i>p</i> = .763	<i>p</i> = .701	<i>p</i> = .853	<i>p</i> = .242
Quality of study					
Poor	25/1.02 [0.82–1.28]	25/0.92 [0.49–1.75]	19/1.15 [0.78–1.70]	20/1.01 [0.69–1.48]	20/1.27 [0.57–2.58]
Satisfactory	4 <sup>f</sup> /1.17 [0.85–1.60]	4 <sup>f</sup> /0.10 [0.99–1.01]	4 <sup>f</sup> /1.07 [0.75–1.53]	4 <sup>f</sup> /0.94 [0.40–2.22]	4 <sup>f</sup> /0.98 [0.44–2.18]
Good	62/1.00 [0.90–1.12]	55/0.99 [0.70–1.39]	39/1.22 [0.58–1.75]	36/1.18 [0.88–1.58]	33/0.85 [0.45–1.60]
Very good	5/0.87 [0.59–1.35]	4 <sup>g</sup> /1.05 [0.97–1.13]	4 <sup>g</sup> /1.18 [0.71–1.96]	2 <sup>h</sup> /1.21 [0.76–1.94]	4 <sup>g</sup> /1.24 [0.80–1.91]
	<i>p</i> = .6812	<i>p</i> = .976	<i>p</i> = .649	<i>p</i> = .685	<i>p</i> = .647
Year of publication	96/ <i>p</i> = .9537	88/ <i>p</i> = .06	66/ <i>p</i> = .080	62/ <i>p</i> = .143	61/ <i>p</i> = .409

Note: The result presented as the number of effect sizes (K)/Overall estimate per level (OR [95% C.I.])/p value.

Bold values indicate as *p* < 0.0001.

<sup>a</sup>Only one study.<sup>70</sup>

<sup>b</sup>4 effect sizes from one study.<sup>34</sup>

<sup>c</sup>3 effect sizes from one study.<sup>10</sup>

<sup>d</sup>7 effect sizes from one study.<sup>61</sup>

<sup>e</sup>4 effect size from one study.<sup>49</sup>

<sup>f</sup>4 effect sizes from one study.<sup>63</sup>

<sup>g</sup>4 effect sizes from one study.<sup>43</sup>

<sup>h</sup>2 effect sizes from one study.<sup>43</sup>

was non-significant for sex and HbA1c (*p* = .101 and .329, respectively).

## 4 | DISCUSSION

To our knowledge, no previous review either quantified the impact of several factors related to patients' characteristics on ADP; or compared their impact among different classes of ADDs. Age, baseline BMI and baseline HbA1c had the greatest impact on the selection of ADDs while patients' sex had the least impact.

The significant variability in the pooled estimate of sex by class of ADDs could be linked to the differences in the number of studies investigating each antidiabetic class, or to the differences in the pharmacological characteristics of ADDs (mainly their safety and tolerability profile). The observed higher prescriptions of GLP1-RA for female patients compared to male patients could be explained in part by previous findings that GLP1-RA was better tolerated and associated with a lower cardiovascular risk among female patients.<sup>71</sup> On the contrary, the significantly lower prescriptions of TZD for female patients

could be explained by the findings that female patients have experienced more side effects from TZD including weight gain, fracture and oedema.<sup>72,73</sup> This suggests a possible consideration of the variability in the effectiveness and tolerability of ADDs between female and male patients when making a decision on the appropriate ADDs in clinical practice. However, because of the limited number of studies examined the majority of antidiabetic classes, more studies are required to have a better understanding regarding the impact of sex on the choice of ADDs.

Despite the risk of SU-related hypoglycaemia being higher among older people, the pooled estimate of SU showed that older people were significantly more likely to use SU. The low cost of SU and the current availability of short-acting second-generation SU (e.g. glipizide) with fewer side effects might be partially responsible for the observed impact of age on SU prescription.<sup>74</sup> This could also reflect the legacy availability of SU for T2DM management as none of the newer ADDs were available 10 years ago, and patients started on SU may have stayed on the same regimen unless they developed intolerable side effects or required additional drug therapy.

Categorisation scheme of binary data	# Of effect sizes (Total k)	Pooled estimate (OR [95% CI])	p value
Age	59		
≥60 vs. <60 years	13	1.06 [0.61–1.83]	<i>p</i> = .942
≥65 vs. <65 years	29	1.01 [0.69–1.46]	
≥70 vs. <70 years	13	0.75 [0.30–1.84]	
≥55 vs. <55 years	4 <sup>a</sup>	0.96 [0.98–1.01]	
Body mass index (BMI)	52		
Obese (BMI ≥ 30 kg/m <sup>2</sup> ) vs. non-obese (BMI < 30 kg/m <sup>2</sup> )	32	1.175 [0.855–1.615]	<i>p</i> = .067
Overweight/obese (BMI ≥ 25 kg/m <sup>2</sup> ) vs. normal/underweight (BMI < 25 kg/m <sup>2</sup> )	13	1.545 [0.546–4.369]	
BMI ≥ 25 vs. BMI 22–25 kg/m <sup>2</sup>	7 <sup>b</sup>	1.018 [0.519–1.996]	
Glycaemic control (HbA1c)			
≥7% (≥53 mmol/mol) vs. <7% (<53 mmol/mol)	13	1.5 [0.34–6.75]	<i>p</i> = .916
≥8% (≥63.9 mmol/mol) vs. <8% (<63.9 mmol/mol)	21	1.05 [0.74–1.51]	

Abbreviations: CI, confidence interval; K, effect sizes; OR, odds ratio.

<sup>a</sup>All from one study.<sup>63</sup>

<sup>b</sup>All from one study.<sup>61</sup>

**TABLE 6** The pooled estimate of all quantified factors before and after excluding the outliers.

Studied factor	Pooled estimate without outliers	Pooled estimate with outliers
Sex	1.00 [0.86–1.16]	0.99 [0.92–1.07]
Age	0.93 [0.66–1.32]	0.96 [0.83–1.10]
BMI	1.19 [0.85–1.67]	1.21 [1.00–1.47]
HbA1c	1.10 [0.81–1.49]	1.06 [0.88–1.29]
Kidney problem	0.89 [0.54–1.47]	0.94 [0.69–1.29]

The safety of newer ADDs (GLP1-RA, SGLT2-I) in older adults was less studied thus prescribers might be less confident to prescribe the newer ADDs for older patients because of the higher concern that elderly patients are more susceptible to adverse reactions.<sup>74–76</sup> Furthermore, the higher cost of newer drugs, the cost of the required monitoring and the familiarity of prescribers with the update in clinical guidelines could contribute to the lower prescription of GLP1-RA and SGLT2-I for older patients. Therefore, further studies investigating prescribing quantity of newer ADDs for older patients are still required since older patients are more likely to have cardiovascular

**TABLE 5** Results of moderator analysis of categorisation scheme of binary data of age, body mass index and HbA1c meta-analysis.

and renal diseases, the situations where the newer ADDs are recommended. The negative significant association between metformin prescription and age could be related to the fact that metformin is not recommended to be prescribed for patients with gastrointestinal complaints, functional impairment or with renal insufficiency, conditions that are increasingly present with increasing age.<sup>75,77,78</sup> This might positively reflect clinical practice adherence to drug characteristics when prescribing metformin to older patients with T2DM.

GLP1-RA and SGLT2-I were reported to have weight loss effect and metformin was accepted to have weight neutral or slight weight loss effects, while SU is associated with weight gain.<sup>79–81</sup> Thereby, the weight effect of ADDs might be responsible for our findings that overweight/obese people were more likely to get a medication with weight neutral/loss effect (GLP1-RA, SGLT2-I, and metformin) but less likely to be prescribed a medication with weight gain effect (SU). Overall, these findings might indirectly reflect a consistency of ADD selection in clinical practice considering patient weight against drug features.

Baseline HbA1c level had the strongest association with insulin prescription where patients with higher baseline HbA1C were more likely to receive insulin, whereas



higher baseline HbA1c had negative weak significant associations with metformin, TZD and DPP4-I prescriptions. All aforementioned associations were consistent with the known effectiveness of each antidiabetic class relevant to HbA1c reduction, which partially indicate clinicians' consideration of disease severity (indicated by HbA1c) when selecting the most appropriate ADDs for each patient. Insulin is known to have the greatest effect on the reduction of HbA1c and this might explain the greater likelihood of prescribing insulin for patients with higher baseline HbA1c.<sup>82,83</sup>

Lastly, the management of diabetes in patients with kidney-related problems is challenging as the impairment in kidney function might affect glucose metabolism and alter drug clearance.<sup>84</sup> This further complicates the selection of an appropriate ADD, considering the need for more frequent adjustment of doses and monitoring for the risk of hypo/hyperglycaemia.<sup>84</sup> Insulin has been considered as the best choice for patients with T2DM and kidney problems, yet still requires close monitoring and dose adjustment.<sup>84</sup> Also, DPP4-I is among the most acceptable option for patients with kidney problems considering dose adjustment based on the agent and degree of impairment.<sup>84</sup> In contrast, metformin is not recommended for patients with kidney disease, and it is contraindicated when eGFR is  $<30\text{ mL/min/1.73 m}^2$ , because of the higher risk of lactic acidosis.<sup>84</sup> Collectively, that could explain the observed associations of higher prescription of insulin and DPP4-I and lower prescription of metformin for patients with kidney-related problems.

Despite that the use of SGLT2-I and GLP1-RA has been recently encouraged by several guidelines especially for patients with established or high risk of cardiovascular or renal diseases because of their cardioprotective and renal protective effects,<sup>9,85–88</sup> the pooled estimates of studies investigating the prescription of SGLT2-I and GLP1-RA for patients with kidney-related problems were not in line with the previous recommendations. Nevertheless, those recommendations are relatively recent while the majority of included studies were conducted early after the introduction of GLP1-RA and SGLT2-I. Therefore, more studies are still required to further investigate prescribing of newer classes for patients with kidney problems in clinical practice considering different stages and types of kidney disease.

#### 4.1 | Strength and limitations

To the best of our knowledge, this is the first SR/MA integrating the results of observational studies assessing the association of several factor with ADP to draw an overall estimate. This review provides a wide range of data by

investigating each factor on seven different antidiabetic classes. Additionally, applying a three-level MA approach to account for the presence of dependency among effect sizes gave an opportunity to answer the research question without losing valuable data and to directly compare different antidiabetic groups.

Nevertheless, all previous results should be interpreted cautiously because of several limitations of the study. First, limited number of studies examined certain classes of ADDs, especially the newer ones; thus, more studies are required to draw a more robust conclusion. Second, the possible presence of publication bias especially for age, BMI and kidney-related problems may have affected the reliability of findings; however, there is no agreed-upon method available to adjust for publication bias in the three-level MA model. Third, bias could have been introduced by including all studies in the pooled estimate regardless the type of data presentation (categorical vs. continuous) and the type of categorisation scheme; yet, subgroup analyses were done and showed no significant impact, and the pooled estimate of each sub-group was reported separately. Lastly, other important factors, including socioeconomic and prescriber-related factors, were much less frequently studied and further investigations are needed.

## 5 | CONCLUSION

In conclusion, all identified factors are crucial to be considered when making a decision regarding the most appropriate ADDs for patients with T2DM. The magnitude, direction and significance of influence of the identified factors on ADP varied according to the type of antidiabetic group. Age, baseline BMI and baseline HbA1c had the greatest impact on the selection of ADDs in which they had statistically significant associations with prescribing of four out of the seven investigated antidiabetic classes. On the other hand, sex had the least impact on ADDs selection which had only a significant influence on GLP1-RA and TZD prescriptions. The findings of this SR&MA could be helpful in determining the need of improving prescribing practice of ADDs by reflecting the consistency of prescribing decision of ADDs with guidelines recommendations and specific drugs features.

### AUTHOR CONTRIBUTIONS

The authors confirm their contribution to the paper as follows: Study concept and design: FM, AK, AM; Study screening, data collection and validation: FM, HY, NA; Data analysis: FM; interpretation of results: FM, AK, AM, TM, CS, GR; Draft manuscript preparation: FM, AK, TM, AM, CS, GR, HY, NA. All authors reviewed and revised the manuscript, and approved the final version.

## ACKNOWLEDGEMENTS

This work is part of a PhD project, and the authors acknowledge Jordan University of Science and Technology for sponsoring this PhD project at the University of Strathclyde.

## CONFLICT OF INTEREST STATEMENT

The authors have no conflict of interest to disclose.

## ORCID

Fatema Mahmoud  <https://orcid.org/0000-0003-2218-224X>

Amanj Kurdi  <https://orcid.org/0000-0001-5036-1988>

## REFERENCES

- Goyal R, Jialal I. *Diabetes Mellitus Type 2*. StatPearls; 2019.
- Sun H, Saeedi P, Karuranga S, et al. IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes Res Clin Pract.* 2021;183:109-119. doi:10.1016/j.diabres.2021.109119
- Papatheodorou K, Banach M, Bekiari E, Rizzo M, Edmonds M. Complications of Diabetes 2017. *J Diabetes Res.* 2018;2018:1-4. doi:10.1155/2018/3086167
- Marin-Penalver JJ, Martin-Timon I, Sevillano-Collantes C, Del Canizo-Gomez FJ. Update on the treatment of type 2 diabetes mellitus. *World J Diabetes.* 2016;7(17):354-395. doi:10.4239/wjcd.v7.i17.354
- Chaudhury A, Duvoor C, Reddy Dendi VS, et al. Clinical review of antidiabetic drugs: implications for type 2 Diabetes mellitus management. *Front Endocrinol (Lausanne).* 2017;8:6. doi:10.3389/fendo.2017.00006
- Davies MJ, D'Alessio DA, Fradkin J, et al. Management of Hyperglycemia in type 2 Diabetes, 2018. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care.* 2018;41(12):2669-2701. doi:10.2337/dci18-0033
- IDF Working Group. *Recommendations For Managing Type 2 Diabetes In Primary Care.* 2017 [www.idf.org/managing-type2-diabetes](http://www.idf.org/managing-type2-diabetes)
- American Diabetes Association (ADA). 9. Pharmacologic approaches to glycemic treatment: standards of medical Care in Diabetes-2020. *Diabetes Care.* 2020;43(Suppl 1):S98-S110. doi:10.2337/dc20-S009
- National Institute of Health and Care Excellence (NICE). *Type 2 diabetes in adults: management.* 2021. Accessed January 30, 2022. <https://www.nice.org.uk/guidance/gid-ng10160/documents/draft-guideline>
- Wilkinson S, Douglas IJ, Williamson E, et al. Factors associated with choice of intensification treatment for type 2 diabetes after metformin monotherapy: a cohort study in UK primary care. *Clin Epidemiol.* 2018;10:1639-1648. doi:10.2147/CLEP.S176142
- Heintjes EM, Overbeek JA, Hall GC, et al. Factors associated with type 2 Diabetes mellitus treatment choice across four European countries. *Clin Ther.* 2017;39(11):2296-2310 e14. doi:10.1016/j.clinthera.2017.09.016
- Chu WM, Ho HE, Huang KH, et al. The prescribing trend of oral antidiabetic agents for type 2 diabetes in Taiwan: an 8-year population-based study. *Medicine (Baltimore).* 2017;96(43):e8257. doi:10.1097/MD.00000000000008257
- Geier AS, Wellmann I, Wellmann J, et al. Patterns and determinants of new first-line antihyperglycaemic drug use in patients with type 2 diabetes mellitus. *Diabetes Res Clin Pract.* 2014;106(1):73-80. doi:10.1016/j.diabres.2014.07.014
- Grabner M, Peng X, Geremakis C, Bae J. Demographic and clinical profiles of type 2 Diabetes mellitus patients initiating canagliflozin versus DPP-4 inhibitors in a large U.S. Managed care population. *J Manag Care Spec Pharm.* 2015;21(12):1204-1212. doi:10.18553/jmcp.2015.21.12.1204
- Zinman B, Wanner C, Lachin JM, et al. Empagliflozin, cardiovascular outcomes, and mortality in type 2 Diabetes. *N Engl J Med.* 2015;373(22):2117-2128. doi:10.1056/NEJMoa1504720
- Neal B, Perkovic V, Mahaffey KW, et al. Canagliflozin and cardiovascular and renal events in type 2 Diabetes. *N Engl J Med.* 2017;377(7):644-657. doi:10.1056/NEJMoa1611925
- Marso SP, Daniels GH, Brown-Frandsen K, et al. Liraglutide and cardiovascular outcomes in type 2 Diabetes. *N Engl J Med.* 2016;375(4):311-322. doi:10.1056/NEJMoa1603827
- Davari M, Khorasani E, Tigabu BM. Factors influencing prescribing decisions of physicians: a review. *Ethiop J Health Sci.* 2018;28(6):795-804. doi:10.4314/ejhs.v28i6.15
- Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71. doi:10.1136/bmj.n71
- Covidence systematic review software. Veritas Health Innovation. [www.covidence.org](http://www.covidence.org)
- Sharifnia SHA, Mohammadzadeh M, Arzani G, et al. Main factors affecting Physicians' prescribing decisions: the Iranian experience. *Iran J Pharm Res.* 2018;17(3):1105-1115.
- Wells G, Shea B, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for Assessing the Quality of Non-Randomized Studies in Meta-Analysis. 2000.
- Chang WW, Boonhat H, Lin RT. Incidence of respiratory symptoms for residents living near a petrochemical industrial complex: a meta-analysis. *Int J Environ Res Public Health.* 2020;17(7):24-74. doi:10.3390/ijerph17072474
- Modesti PA, Reboldi G, Cappuccio FP, et al. Panethnic differences in blood pressure in Europe: a systematic review and meta-analysis. *PLoS One.* 2016;11(1):e0147601. doi:10.1371/journal.pone.0147601
- Van den Noortgate W, López-López JA, Marín-Martínez F, Sánchez-Meca J. Three-level meta-analysis of dependent effect sizes. *Behav Res Methods.* 2013;45(2):576-594. doi:10.3758/s13428-012-0261-6
- Cheung MW. Modeling dependent effect sizes with three-level meta-analyses: a structural equation modeling approach. *Psychol Methods.* 2014;19(2):211-229. doi:10.1037/a0032968
- Cheung MWL. *Meta-Analysis: A Structural Equation Modeling Approach.* John Wiley & Sons, Inc.; 2015.
- Assink M, Wibbelink CJM. Fitting three-level meta-analytic models in R: a step-by-step tutorial. *TQMP.* 2016;12(3):154-174. doi:10.20982/tqmp.12.3.p154
- Harrer M, Cuijpers P, Furukawa TA, Ebert DD. *Doing Meta-Analysis with R: A Hands-on Guide.* 1st ed. Chapman & Hall/CRC Press; 2021.
- Fernández-Castilla B, Declercq L, Jamshidi L, Beretvas SN, Onghena P, Van den Noortgate W. Detecting selection bias in

- meta-analyses with multiple outcomes: a simulation Study. *J Exp Educ.* 2021;89(1):125-144. doi:10.1080/00220973.2019.1582470
31. Page MJ, Sterne JAC, Higgins JPT, Egger M. Investigating and dealing with publication bias and other reporting biases in meta-analyses of health research: a review. *Res Synth Methods.* 2021;12(2):248-259. doi:10.1002/jrsm.1468
  32. Viechtbauer W, Cheung MW. Outlier and influence diagnostics for meta-analysis. *Res Synth Methods.* 2010;1(2):112-125. doi:10.1002/jrsm.11
  33. Cook RD. Detection of influential observation in linear regression. *Dent Tech.* 1977;19(1):15-18. doi:10.1080/00401706.1977.10489493
  34. Winkelmayr WC, Stedman MR, Pogantsch M, et al. Guideline-conformity of initiation with oral hypoglycemic treatment for patients with newly therapy-dependent type 2 diabetes mellitus in Austria. *Pharmacoepidemiol Drug Saf.* 2010;20(1):57-65. doi:10.1002/pds.2059
  35. Abdelmoneim AS, Eurich DT, Gamble J-M, Simpson SH. Use patterns of antidiabetic regimens by patients with type 2 diabetes. *Can J Diabetes.* 2013;37(6):394-400. doi:10.1016/j.jcjd.2013.04.008
  36. Brouwer ES, West SL, Kluckman M, et al. Initial and subsequent therapy for newly diagnosed type 2 diabetes patients treated in primary care using data from a vendor-based electronic health record. *Pharmacoepidemiol Drug Saf.* 2012;21(9):920-928. doi:10.1002/pds.2262
  37. Liu CH, Chen ST, Chang CH, Chuang LM, Lai MS. Prescription trends and the selection of initial oral antidiabetic agents for patients with newly diagnosed type 2 diabetes: a nationwide study. *Public Health.* 2017;152:20-27. doi:10.1016/j.puhe.2017.06.008
  38. Wang T-Y, Eguale T, Tamblyn R. Guidelines adherence in the treatment of patients with newly diagnosed type 2 diabetes: a historical cohort comparing the use of metformin in Quebec pre and post-Canadian Diabetes Association guidelines. *BMC Health Serv Res.* 2013;13(1):442. doi:10.1186/1472-6963-13-442
  39. Fujihara K, Igarashi R, Matsunaga S, et al. Comparison of baseline characteristics and clinical course in Japanese patients with type 2 diabetes among whom different types of oral hypoglycemic agents were chosen by diabetes specialists as initial monotherapy (JDDM 42). *Medicine.* 2017;96(7):e6122. doi:10.1097/md.0000000000006122
  40. Desai NR, Shrank WH, Fischer MA, et al. Patterns of medication initiation in newly diagnosed diabetes mellitus: quality and cost implications. *Am J Med.* 2012;125(3):302 e1-7. doi:10.1016/j.amjmed.2011.07.033
  41. Grimes RT, Bennett K, Tilson L, Usher C, Smith SM, Henman MC. Initial therapy, persistence and regimen change in a cohort of newly treated type 2 diabetes patients. *Br J Clin Pharmacol.* 2015;79(6):1000-1009. doi:10.1111/bcp.12573
  42. Cai B, Katz L, Alexander CM, Williams-Herman D, Girman CJ. Characteristics of patients prescribed sitagliptin and other oral antihyperglycaemic agents in a large US claims database. *Int J Clin Pract.* 2010;64(12):1601-1608. doi:10.1111/j.1742-1241.2010.02516.x
  43. Saine ME, Carbonari DM, Newcomb CW, et al. Determinants of saxagliptin use among patients with type 2 diabetes mellitus treated with oral anti-diabetic drugs. *BMC Pharmacol Toxicol.* 2015;16:8. doi:10.1186/s40360-015-0007-z
  44. Ou HT, Chang KC, Liu YM, Wu JS. Recent trends in the use of antidiabetic medications from 2008 to 2013: a nation-wide population-based study from Taiwan. *J Diabetes.* 2017;9(3):256-266. doi:10.1111/1753-0407.12408
  45. Stargardt T, Yin DD, Alexander CM. Treatment choice and effectiveness of adding sulphonylurea or glitazones to metformin for the treatment of type 2 diabetes mellitus. *Diabetes Obes Metab.* 2009;11(5):491-497. doi:10.1111/j.1463-1326.2008.00991.x
  46. Payk SL, Drew RH, Smith JD, Jiroutek MR, Holland MA. Sulphonylurea prescribing patterns after the introduction of DPP-4 inhibitors and GLP-1 receptor agonists. *Clin Ther.* 2015;37(7):1477-1482 e1. doi:10.1016/j.clinthera.2015.04.011
  47. Zhang Q, Rajagopalan S, Mavros P, et al. Baseline characteristic differences between patients prescribed sitagliptin vs. other oral antihyperglycemic agents: analysis of a US electronic medical record database. *Curr Med Res Opin.* 2010;26(7):1697-1703. doi:10.1185/03007995.2010.489029
  48. Morita Y, Murayama H, Odawara M, Bauer M. Treatment patterns of drug-naive patients with type 2 diabetes mellitus: a retrospective cohort study using a Japanese hospital database. *Diabetol Metab Syndr.* 2019;11:90. doi:10.1186/s13098-019-0486-y
  49. Kim J, Park S, Kim H, Je NK. National trends in metformin-based combination therapy of oral hypoglycaemic agents for type 2 diabetes mellitus. *Eur J Clin Pharmacol.* 2019;75(12):1723-1730. doi:10.1007/s00228-019-02751-9
  50. Heintjes EM, Overbeek JA, Hall GC, et al. Factors associated with type 2 Diabetes mellitus treatment choice across four European countries. *Clin Ther.* 2017;39(11):2296-2310.e14. doi:10.1016/j.clinthera.2017.09.016
  51. Nicolucci A, Charbonnel B, Gomes MB, et al. Treatment patterns and associated factors in 14 668 people with type 2 diabetes initiating a second-line therapy: results from the global DISCOVER study programme. *Diabetes Obes Metab.* 2019;21(11):2474-2485. doi:10.1111/dom.13830
  52. Hartmann B, Lanzinger S, van Mark G, et al. Treatment intensification strategies after initial metformin therapy in adult patients with type-2 diabetes: results of the DPV and DIVE registries. *Acta Diabetol.* 2020;57(2):229-236. doi:10.1007/s00592-019-01409-3
  53. Longato E, Di Camillo B, Sparacino G, Gubian L, Avogaro A, Fadini GP. Cardiovascular outcomes of type 2 diabetic patients treated with SGLT-2 inhibitors versus GLP-1 receptor agonists in real-life. *BMJ Open Diabetes Res Care.* 2020;8(1):e001451. doi:10.1136/bmjdr-2020-001451
  54. Ackermann RT, Wallia A, Brien MJ, et al. Correlates of second-line type 2 diabetes medication selection in the USA. *BMJ Open Diabetes Res Care.* 2017;5(1):e000421. doi:10.1136/bmjdr-2017-000421
  55. Whyte MB, Hinton W, McGovern A, et al. Disparities in glycaemic control, monitoring, and treatment of type 2 diabetes in England: a retrospective cohort analysis. *PLoS Med.* 2019;16(10):e1002942. doi:10.1371/journal.pmed.1002942
  56. Arnold SV, McGuire DK, Inzucchi SE, et al. Assessing use of patient-focused pharmacotherapy in glycemic management through the Diabetes collaborative registry (DCR). *J Diabetes Complicat.* 2018;32(11):1035-1039. doi:10.1016/j.jdiacomp.2018.02.009
  57. Arnold SV, Echouffo-Tcheugui JB, Lam CSP, et al. Patterns of glucose-lowering medication use in patients with type 2

- diabetes and heart failure. Insights from the Diabetes collaborative registry (DCR). *Am Heart J.* 2018;203:25-29. doi:10.1016/j.ahj.2018.05.016
58. Zaharan NL, Williams D, Bennett K. Prescribing of antidiabetic therapies in Ireland: 10-year trends 2003-2012. *Ir J Med Sci.* 2014;183(2):311-318. doi:10.1007/s11845-013-1011-1
  59. Zoberi KA, Salas J, Morgan CN, Scherrer JF. Comparison of family medicine and general internal medicine on Diabetes management. *Mo Med.* 2017;114(3):187-194.
  60. Montvida O, Shaw J, Atherton JJ, Stringer F, Paul SK. Long-term trends in Antidiabetes drug usage in the U.S.: real-world evidence in patients newly diagnosed with type 2 Diabetes. *Diabetes Care.* 2018;41(1):69-78. doi:10.2337/dc17-1414
  61. Katakami N, Mita T, Takahara M, et al. Baseline characteristics of patients with type 2 Diabetes initiating second-line treatment in Japan: findings from the J-DISCOVER Study. *Diabetes Therapy.* 2020;11(7):1563-1578. doi:10.1007/s13300-020-00846-6
  62. Kostev K, Dippel FW, Rathmann W. Predictors of insulin initiation in metformin and sulfonylurea users in primary care practices: the role of kidney function. *J Diabetes Sci Technol.* 2014;8(5):1023-1028. doi:10.1177/1932296814532616
  63. Dhanaraj E, Raval A, Yadav R, Bhansali A, Tiwari P. Erratum to: prescribing pattern of antidiabetic drugs and achievement of glycemic control in T2DM patients tertiary care hospital in North India. *Int J Diabetes Dev Countries.* 2013;33:140-146. doi:10.1007/s13410-013-0123-5
  64. Yu M, Mody R, Landó LF, et al. Characteristics associated with the choice of first injectable therapy among US patients with type 2 Diabetes. *Clin Ther.* 2017;39(12):2399-2408. doi:10.1016/j.clinthera.2017.11.001
  65. Levin P, Wei W, Zhou S, Xie L, Baser O. Outcomes and treatment patterns of adding a third agent to 2 OADs in patients with type 2 Diabetes. *J Managed Care Pharmacy.* 2014;20:501-512. doi:10.18553/jmcp.2014.20.5.501
  66. Gentile S, Strollo F, Viazzi F, et al. Five-year predictors of insulin initiation in people with type 2 Diabetes under real-life conditions. *J Diabetes Res.* 2018;2018:1-10. doi:10.1155/2018/7153087
  67. Korytkowski MT, Brooks M, Lombardero M, et al. Use of an electronic medical record (EMR) to identify glycemic intensification strategies in type 2 Diabetes. *J Diabetes Sci Technol.* 2014;9(3):593-601. doi:10.1177/1932296814564183
  68. Hirsch IB, Xu Y, Davis KL, Calingaert B. Patient factors associated with glucagonlike peptide 1 receptor agonist use with and without insulin. *Endocr Pract.* 2011;17(5):707-716. doi:10.4158/EP11014.OR
  69. van den Boom L, Kaiser M, Kostev K. Prevalence of insulin as a first-line therapy and associated factors in people with type 2 diabetes in German primary care practices. *Diabet Med.* 2020;37(8):1333-1339. doi:10.1111/dme.14338
  70. Moreno Juste A, Menditto E, Orlando V, et al. Treatment patterns of Diabetes in Italy: a population-based Study. *Front Pharmacol.* 2019;10:870. doi:10.3389/fphar.2019.00870
  71. Raparelli V, Elharram M, Moura CS, et al. Sex differences in cardiovascular effectiveness of newer glucose-lowering drugs added to metformin in type 2 diabetes mellitus. *J Am Heart Assoc.* 2020;9(1):e012940.
  72. Campesi I, Franconi F, Seghieri G, Meloni M. Sex-sex-related therapeutic approaches for cardiovascular complications associated with diabetes. *Pharmacol Res.* 2017;119:195-207. doi:10.1016/j.phrs.2017.01.023
  73. Joung K-I, Jung G-W, Park H-H, Lee H, Park S-H, Shin J-Y. Sex differences in adverse event reports associated with antidiabetic drugs. *Sci Rep.* 2020;10(1):17545. doi:10.1038/s41598-020-74000-4
  74. Yakaryılmaz FD, Öztürk ZA. Treatment of type 2 diabetes mellitus in the elderly. *World J Diabetes.* 2017;8(6):278-285. doi:10.4239/wjd.v8.i6.278
  75. Kim KS, Kim SK, Sung KM, Cho YW, Park SW. Management of type 2 diabetes mellitus in older adults. *Diabetes Metab J.* 2012;36(5):336-344. doi:10.4093/dmj.2012.36.5.336
  76. Lublóy Á. Factors affecting the uptake of new medicines: a systematic literature review. *BMC Health Serv Res.* 2014;14:469. doi:10.1186/1472-6963-14-469
  77. Schlender L, Martinez YV, Adeniji C, et al. Efficacy and safety of metformin in the management of type 2 diabetes mellitus in older adults: a systematic review for the development of recommendations to reduce potentially inappropriate prescribing. *BMC Geriatr.* 2017;17(1):227. doi:10.1186/s12877-017-0574-5
  78. American DA. 9. Pharmacologic approaches to glycemic treatment: standards of medical Care in Diabetes-2020. *Diabetes Care.* 2020;43(Suppl 1):S98-S110. doi:10.2337/dc20-S009
  79. Vilsboll T, Christensen M, Junker AE, Knop FK, Gluud LL. Effects of glucagon-like peptide-1 receptor agonists on weight loss: systematic review and meta-analyses of randomised controlled trials. *BMJ.* 2012;344:d7771. doi:10.1136/bmj.d7771
  80. Wang H, Yang J, Chen X, Qiu F, Li J. Effects of sodium-glucose cotransporter 2 inhibitor monotherapy on weight changes in patients with type 2 Diabetes mellitus: a Bayesian network meta-analysis. *Clin Ther.* 2019;41(2):322-334.e11. doi:10.1016/j.clinthera.2019.01.001
  81. Apovian CM, Okemah J, O'Neil PM. Body weight considerations in the Management of Type 2 Diabetes. *Adv Ther.* 2019;36(1):44-58. doi:10.1007/s12325-018-0824-8
  82. Sherifali D, Nerenberg K, Pullenayegum E, Cheng JE, Gerstein HC. The effect of Oral antidiabetic agents on A1C levels. *Diabetes Care.* 2010;33(8):1859-1864. doi:10.2337/dc09-1727
  83. Chaudhury A, Duvoor C, Reddy Dendi VS, et al. Clinical review of antidiabetic drugs: implications for type 2 Diabetes mellitus management. *Front Endocrinol (Lausanne).* 2017;8:6. doi:10.3389/fendo.2017.00006
  84. Betônico CCR, Titan SMO, Correa-Giannella MLC, Nery M, Queiroz M. Management of diabetes mellitus in individuals with chronic kidney disease: therapeutic perspectives and glycemic control. *Clinics (Sao Paulo).* 2016;71(1):47-53. doi:10.6061/clinics/2016(01)08
  85. Górriz JL, Soler MJ, Navarro-González JF, et al. GLP-1 receptor agonists and diabetic kidney disease: a call of attention to nephrologists. *J Clin Med.* 2020;9(4):947. doi:10.3390/jcm9040947
  86. Ninčević V, Omanović Kolaric T, Roguljić H, Kizivat T, Smolić M, Bilić ČI. Renal benefits of SGLT 2 inhibitors and GLP-1 receptor agonists: evidence supporting a paradigm shift in the medical Management of Type 2 Diabetes. *Int J Mol Sci.* 2019;20(23):5831. doi:10.3390/ijms20235831
  87. Li J, Albajrami O, Zhuo M, Hawley CE, Paik JM. Decision algorithm for prescribing SGLT2 inhibitors and GLP-1 receptor agonists for diabetic kidney disease. *Clin J Am Soc Nephrol.* 2020;15(11):1678-1688. doi:10.2215/CJN.02690320

88. American Diabetes Association. Introduction: Standards of medical Care in Diabetes-2020. *Diabetes Care*. 2020;43(Suppl 1):S1-s2. doi:[10.2337/dc20-Sint](https://doi.org/10.2337/dc20-Sint)

### SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Mahmoud F, Mullen A, Sainsbury C, et al. Meta-analysis of factors associated with antidiabetic drug prescribing for type 2 diabetes mellitus. *Eur J Clin Invest*. 2023;00:e013997. doi:[10.1111/eci.13997](https://doi.org/10.1111/eci.13997)