

# The highly diverse plasmid population found in *Escherichia coli* colonizing travellers to Laos and its role in antimicrobial resistance gene carriage

Snaith, Ann E; Dunn, Steven J; Moran, Robert A; Newton, Paul N; Dance, David A B; Davong, Viengmon; Kuenzli, Esther; Kantele, Anu; Corander, Jukka; McNally, Alan

DOI:

[10.1099/mgen.0.001000](https://doi.org/10.1099/mgen.0.001000)

License:

Creative Commons: Attribution (CC BY)

## Document Version

Publisher's PDF, also known as Version of record

## Citation for published version (Harvard):

Snaith, AE, Dunn, SJ, Moran, RA, Newton, PN, Dance, DAB, Davong, V, Kuenzli, E, Kantele, A, Corander, J & McNally, A 2023, 'The highly diverse plasmid population found in *Escherichia coli* colonizing travellers to Laos and its role in antimicrobial resistance gene carriage', *Microbial Genomics*, vol. 9, no. 5, 001000. <https://doi.org/10.1099/mgen.0.001000>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# The highly diverse plasmid population found in *Escherichia coli* colonizing travellers to Laos and its role in antimicrobial resistance gene carriage

Ann E. Snaith<sup>1</sup>, Steven J. Dunn<sup>1</sup>, Robert A. Moran<sup>1</sup>, Paul N. Newton<sup>2,3,4</sup>, David A. B. Dance<sup>2,3,4</sup>, Viengmon Davong<sup>2</sup>, Esther Kuenzli<sup>5,6</sup>, Anu Kantele<sup>7,8,9</sup>, Jukka Corander<sup>10,11,12</sup> and Alan McNally<sup>1,\*</sup>

## Abstract

Increased colonization by antimicrobial-resistant organisms is closely associated with international travel. This study investigated the diversity of mobile genetic elements involved with antimicrobial resistance (AMR) gene carriage in extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli* that colonized travellers to Laos. Long-read sequencing was used to reconstruct complete plasmid sequences from 48 isolates obtained from the daily stool samples of 23 travellers over a 3 week period. This method revealed a collection of 105 distinct plasmids, 38.1% ( $n=40$ ) of which carried AMR genes. The plasmids in this population were diverse, mostly unreported and included 38 replicon types, with F-type plasmids ( $n=23$ ) the most prevalent amongst those carrying AMR genes. Fine-scale analysis of all plasmids identified numerous AMR gene contexts and emphasized the importance of IS elements, specifically members of the IS6/IS26 family, in the evolution of complex multidrug resistance regions. We found a concerning convergence of ESBL and colistin resistance determinants, with three plasmids from two different F-type lineages carrying *bla*<sub>CTX-M</sub> and *mcr* genes. The extensive diversity seen here highlights the worrying probability that stable new vehicles for AMR will evolve in *E. coli* populations that can disseminate internationally through travel networks.

## DATA SUMMARY

Long-read sequencing data are available through the National Center for Biotechnology Information (NCBI) under the BioProject PRJNA853172. Complete plasmid sequences have been uploaded to GenBank with accession numbers in Table S1 (available in the online version of this article). The authors confirm all supporting data, code and protocols have been provided within the article or through supplementary data files.

Received 24 August 2022; Accepted 24 February 2023; Published 12 May 2023

**Author affiliations:** <sup>1</sup>Institute of Microbiology and Infection, College of Medical and Dental Sciences, University of Birmingham, Birmingham, B15 2TT, UK; <sup>2</sup>Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit, Microbiology Laboratory, Mahosot Hospital, Rue Mahosot, Vientiane, Laos; <sup>3</sup>Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, UK; <sup>4</sup>Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, UK; <sup>5</sup>Department of Medicine, Swiss Tropical and Public Health Institute of Basel, Basel, Switzerland; <sup>6</sup>University of Basel, Basel, Switzerland; <sup>7</sup>Meilahti Infectious Diseases and Vaccine Research Center, MeVac, Biomedicum 1, Haartmaninkatu 8, Helsinki University Hospital and University of Helsinki, Helsinki 00290, Finland; <sup>8</sup>Multidisciplinary Center of Excellence in Antimicrobial Resistance Research, University of Helsinki, Helsinki, Finland; <sup>9</sup>Human Microbiome Research Program, Faculty of Medicine, Haartmaninkatu 4, 00014 University of Helsinki, Helsinki, Finland; <sup>10</sup>Parasites and Microbes Programme, Wellcome Sanger Institute, Wellcome Genome Campus, Cambridge, UK; <sup>11</sup>Department of Biostatistics, Faculty of Medicine, University of Oslo, Oslo, Norway; <sup>12</sup>Department of Mathematics and Statistics, Helsinki Institute of Information Technology, University of Helsinki, Helsinki, Finland.

\*Correspondence: Alan McNally, a.mcnally.1@bham.ac.uk

**Keywords:** plasmid; antimicrobial resistance; Laos.

**Abbreviations:** AMR, antimicrobial resistance; CTAB, cetyltrimethylammonium bromide; ESBL, extended-spectrum beta-lactamase; IS, insertion sequence; LB, lysogeny broth; MDR, multidrug-resistant; MGE, mobile genetic element; MITE, miniature inverted-repeat tandem element; ST, sequence type; TE, transposable element.

**Data statement:** All supporting data, code and protocols have been provided within the article or through supplementary data files. Four supplementary figures and eight supplementary tables are available with the online version of this article.

001000 © 2023 The Authors



This is an open-access article distributed under the terms of the Creative Commons Attribution License. This article was made open access via a Publish and Read agreement between the Microbiology Society and the corresponding author's institution.

### Impact Statement

The global spread of antimicrobial resistance (AMR) is closely associated with international travel. AMR is a severe global concern and has compromised treatment options for many bacterial pathogens, among them pathogens carrying extended-spectrum beta-lactamase (ESBL) and colistin resistance genes. Colonizing multidrug-resistant (MDR) organisms have the potential to cause serious consequences. Infections caused by antimicrobial-resistant bacteria, including MDR bacteria, are associated with longer hospitalization, poorer patient outcomes, greater mortality and higher costs compared to infections with susceptible bacteria. This study elucidates the numerous different types of plasmids carrying AMR genes in colonizing ESBL-producing *Escherichia coli* isolates found in faecal samples from travellers to Vientiane, Laos. Here we add to known databases of AMR plasmids by adding these MDR plasmids found in Southeast Asia, an area of high AMR prevalence. We characterized novel AMR plasmids, including complex ESBL ( $bla_{CTX-M}$ ) and colistin (*mcr*) resistance co-carriage plasmids, emphasizing the potential exposure of travellers to Laos to a wide variety of mobile genetic elements that may facilitate global AMR spread. This in-depth study has revealed further detail concerning the numerous factors that may influence AMR transfer, and therefore potential routes of AMR spread internationally, and is a step towards finding methods to combat AMR spread.

## INTRODUCTION

Infections caused by antimicrobial-resistant organisms are harder to treat, lengthen hospital stays, increase mortality rates and place a significant financial burden on healthcare institutes [1]. It is increasingly important to characterize the mechanisms that allow antimicrobial resistance (AMR) to spread worldwide, compromising treatment options for many bacterial pathogens. The rapid spread of AMR has been closely associated with international travel [2–5]. Clinically relevant AMR determinants are commonly found in Gram-negative bacteria colonizing travellers returning from regions with the highest AMR prevalence, including Southeast Asia [3, 6, 7].

*Escherichia coli*, a Gram-negative human gut commensal and opportunistic extraintestinal pathogen, is an important vector for AMR [8, 9]. As exemplified by pandemic multidrug-resistant (MDR) lineages such as ST131 [10, 11], *E. coli* is capable of acquiring and maintaining multiple AMR determinants and exhibiting resistance to multiple classes of antibiotics. The presence of combinations of AMR genes can significantly impact therapeutic options. The limited treatment options for ESBL-resistant organisms make these a cause for concern. Colistin is a last-resort antibiotic included in the Reserve category of the World Health Organization (WHO) Essential Medicines List [12].

In *E. coli* and other Gram-negative bacteria, carriage of both colistin resistance genes and production of extended spectrum beta-lactamases (ESBLs) is concerning, as it suggests a potentially stable environment for the accumulation of further resistance, for example, carbapenemases, severely limiting treatment options [9, 13, 14]. ESBL resistance genes, including  $bla_{CTX-M}$  variants, and *mcr* genes that confer colistin resistance, are commonly found in *E. coli* carried by returning travellers [3, 6, 7, 15]. AMR genes can move intra- or inter-cellularly and accumulate at single sites in association with mobile genetic elements (MGEs) [16–19]. Plasmids are extrachromosomal genetic elements that can transfer horizontally between bacteria of the same or different species and are strongly associated with the spread of AMR [20, 21]. In *E. coli* and other members of the Enterobacterales, AMR genes have been found in many different plasmid types [20]. Reports of stable, epidemic and internationally distributed plasmids [22–24] highlight the threat successful plasmid lineages pose and the importance of understanding the mechanisms by which they acquire and accumulate AMR genes.

We recently characterized the dynamics of the acquisition of MDR Gram-negative organisms in real time during travel to Vientiane, Laos. These MDR Gram-negative organisms had a surprisingly high co-prevalence of ESBLs and colistin resistance genes [7]. Here we explore the pattern of AMR carriage and context in *E. coli* from that cohort using long-read sequencing to understand the contexts of AMR genes and the role of MGEs, particularly plasmids, in the acquisition of drug-resistant *E. coli* by travellers to a region of high AMR prevalence. The 48 representative isolates for long-read sequencing in this study were selected for their AMR and plasmid replicon profiles and were collected on a daily basis in an area of high AMR prevalence, enabling continuous monitoring of the drug-resistant *E. coli* that colonized study participants. Continuous sampling facilitated the examination of common and circulating plasmids in the *E. coli* population, across a variety of sequence types (STs) and at different study time points [7]. Our data show the diversity and widespread distribution of numerous distinct AMR plasmids acquired by these travellers in Vientiane, Laos and the multiple different potential routes of AMR spread by plasmids and highlighted the complex nature of plasmids carrying both ESBL resistance and *mcr* genes.

## METHODS

### Study design and sample source

The *E. coli* isolates used here were collected as part of a study [7] looking at the dynamics of gut colonization of 21 volunteers attending a medical course in Vientiane, Laos. Faecal samples were taken daily during the 22 day period. Samples were processed, shipped, stored and handled as previously described by Kantele *et al.* [7]. Here ESBL-positive isolates were cultured from faecal samples after initial screening on CHROMagar ESBL agar plates (CHROMagar, Paris, France) at the Microbiology Laboratory of Mahosot Hospital, Vientiane, Laos and after transportation further screened with chromID ESBL chromogenic medium (bioMérieux) by the University of Helsinki, Helsinki, Finland [7]. These ESBL isolates [7] were used in this study. In order to explore the pattern of AMR in this traveller dataset and the role of MGEs we prepared hybrid assemblies and annotated plasmid sequences identified to locate resistance genes and potential routes for spread.

### Selection of *E. coli* isolates for MinION sequencing

Whole-genome sequences of 292 ESBL-positive Gram-negative isolates from the Laos study were previously generated [7] and are available under National Center for Biotechnology Information (NCBI) BioProject accession number PRJNA558187. Illumina-generated whole-genome sequences of *E. coli* isolates ( $n=219$ ) were sequence typed using mlst (v2.15) (<https://github.com/tseemann/mlst>). Isolates were selected for MinION sequencing ( $n=48$ ) by deduplication of longitudinal patient samples, and by using ABRicate (v0.8.10) (<https://github.com/tseemann/abricate>) to identify unique elements such as plasmid replicons [PlasmidFinder (v0.8.10) [25]] and antibiotic resistance genes [ResFinder (v0.8.10) [26]] from the short-read assemblies generated with SPAdes (v. 3.13.0) [27].

### DNA extraction and sequencing

*E. coli* were cultured overnight on UTI chromogenic agar (Sigma) at 37 °C. After purity checks single colonies were subcultured overnight in lysogeny broth (LB) (Miller) (shaking, 37 °C). For the majority of isolates DNA was extracted using Monarch Genomic DNA Extraction kit (NEB), but in some instances a lower quantity and quality yield was obtained. In these instances, we observed atypical precipitates and opted for extraction using phenol/chloroform with cetyltrimethylammonium bromide (CTAB). The extracted DNA was sequenced over four runs of MinION (Oxford Nanopore Technology) using R9.4.1 flow cells. Three runs were prepared using Ligation Protocol (LSK-SQK109) and one run was prepared with the Rapid Barcoding Sequencing kit (SQK-RBK004), with both protocols modified to a one-pot implementation.

### Long-read sequencing analysis

Data was basecalled using Guppy (v0.5.1) <https://github.com/nanoporetech/pyguppyclient>). The quality of the data was examined using NanoPlot (v1.28.2) [28]. In read files where coverage was high, filtlong (>100×) (v0.2.0) (<https://github.com/rrwick/Filtlong>) was used to select the best-read files available and the coverage was limited to 100×. Barcodes were trimmed using qcat (v1.0.1) (<https://github.com/nanoporetech/qcat>). Hybrid assemblies using the Illumina reads were created with Unicycler (v0.4.7) [29] and visualized in Bandage (v0.8.1) [30]. In some instances, Unicycler was unable to fully resolve the assembly (generating only circular sequences); in these cases we took a long-read-first approach, using Flye (v2.6) [31] to assemble the long read set into a gfa file that was then provided to the Unicycler pipeline. Assemblies were analysed using ABRicate (v0.8.10) with the ResFinder [26, 32] and PlasmidFinder databases [25]. TigSPLIT (<https://github.com/stevenjdunn/TigSPLIT>) was used to extract contigs, allowing detailed analysis of the location of plasmid replicons in relation to the resistance genes present in each isolate.

### Analysis of plasmid sequences

Plasmid contigs were annotated using Prokka (v1.14.6) [33]. ABRicate (v0.8.10) was used to identify plasmid replicons with the PlasmidFinder database and AMR genes were identified with ResFinder (85% coverage and identity cutoffs). In situations where PlasmidFinder did not provide a type, plasmid contigs were compared to known plasmid replicons for type assignments using BLAST. Where possible, plasmid replicons were sub-typed using the PubMLST database (<https://pubmlst.org/>) [34, 35]. SnapGene (v5.2.4) was used to visualize Prokka-annotated plasmid contigs. ISFinder [36] was used to identify insertion sequence (IS) elements. NCBI BLAST (v2.5.0+) was used to compare plasmids to reference plasmids (Table S2). The NCBI non-redundant nucleotide database was queried with BLASTn to determine whether plasmids identified here had been seen elsewhere. BLASTn and tBLASTn were used to compare similar plasmid regions and to identify homologues of known plasmid genes. IQ-TREE (v2.0.3) [37] and Panaroo (v1.2.3) [38] were used to compare plasmids. ISEScan (v1.7.2.3) was used to quantify IS elements in plasmid assemblies at an element-family level [39].

### Identification of small plasmids from Illumina dataset

NCBI BLAST (v2.5.0+) was used to search draft genomes for plasmid-specific 100 bp signature sequences (Table S3). Signature sequences, as described previously [22], comprise 50 bp each of a mobile genetic element and its immediately adjacent

sequences such that they span the specific junction generated by a given insertion or deletion event. This facilitated the detection of specific plasmid lineages, whether they were represented by single or multiple contigs in the Illumina dataset [7].

## RESULTS

### Isolates from Laos contain a broad diversity of plasmids and resistance genes

A total of 163 complete plasmids were obtained from the hybrid-assembled genomes of 48 *E. coli* isolates that colonized 21 participants and their contacts (Table S4). Isolates harboured one to eight plasmids each, except for a single ST1722 isolate that did not contain any plasmids. The 163 plasmids were typed by size, replicon type and AMR gene carriage, and identical or almost-identical plasmids (Table S3) were deduplicated, resulting in a total of 105 distinct plasmids. These 105 plasmids were named pLAO1–pLAO105 (Tables S1 and S4) and ranged in size from 1531 to 259739 bp, with around half (53.3%,  $n=56$ ) smaller than 25 kb. Forty of the 105 plasmids (38.1%) contained one or more AMR genes. Plasmids that did not carry AMR genes were generally smaller (mean size: 23 554 bp), than those that did (mean size: 97 260 bp). However, pLAO10 (GenBank accession OP242224), pLAO78 (OP242289) and pLAO84 (OP242239) are notable ColE1-like plasmids that carry one–five AMR genes each and range in size from just 5540 bp to 22368 bp (Table S4).


Thirty-eight different replicon types were identified in the collection of 105 plasmids. Types present amongst the small plasmids included: theta-replicating plasmids that initiate replication with RNA primers ( $\theta$ -RNA,  $n=26$ ) or replication proteins ( $\theta$ -Rep,  $n=9$ ), rolling-circle plasmids ( $n=6$ ) and Q-type plasmids that replicate by strand displacement ( $n=3$ ). Amongst large plasmids (>25 kb), F-types ( $n=22$ ) were dominant, and their replicons were sub-typed using the PubMLST database (Table S4). Other large plasmids include phage plasmids ( $n=9$ , including Y- or p0111 types) and X-types ( $n=5$ ). Nine large plasmids were co-integrates, which included multiple replicons of disparate types and all contained one or more AMR genes (Fig. 1). Ten plasmids could not be assigned a type (Fig. 1). In total, 76.3% ( $n=29$ ) of plasmid types contained AMR genes, with the most prevalent being FII-46:FIB-20 ( $n=4$ ), X1 ( $n=3$ ),  $\theta$ -RNA ( $n=3$ ), FII-2 ( $n=2$ ), FII-46:FIB-10 ( $n=2$ ), p0111 ( $n=2$ ) and Y ( $n=2$ ) (Fig. 1, Table S4). AMR plasmids were present across all participants throughout the study (Tables S4 and S5), with carriage occurring sporadically and transiently during the study period.

We found evidence suggesting the circulation of plasmids within Lao *E. coli* in the traveller population. One example, the ColE1-like plasmid pLAO84 (GenBank accession: OP242239), which carries the tetracycline resistance determinant *tet(A)*, was found in the complete genomes of *E. coli* of two different STs, ST195 and ST34, that were acquired by two different participants (Pt33 and Pt40). Mapping contigs assembled from Illumina data against the complete pLAO84 sequence confirmed plasmid circulation, with its complete or fragmented sequence detected in seven additional *E. coli* isolates (Fig. S1). These isolates were obtained from six different participants (including Pt33 and Pt40) over a 10 day period. It is not possible to determine where or when pLAO84 transferred between its hosts, or how the hosts were acquired by these participants. Another example of a potential circulating plasmid, the Q1 plasmid pLAO60 (GenBank accession OP242237) that carries aminoglycoside and beta-lactam resistance genes (Table S4), was present in the complete genome of the ST542 isolate (LA124) and in one genome (LA230) in the wider Illumina dataset (Table S4 and Fig. S2). The Q1 plasmid in isolate LA230 was missing a 101 bp segment that appears to have been lost in a homologous recombination event.

Another notable finding was the presence of phage and virulence plasmids in this collection. A Y-type phage-plasmid, pLAO59 (GenBank accession OP242238), appears to have lost key genes for phage body synthesis, potentially in deletion events mediated by *ISKpn26* and *IS1294*, and instead carries genes that confer resistance to six antibiotic classes and genes that confer resistance to mercury, copper and silver (Fig. S3). The FII-18:FIB-1 pLAO32 is related to the colicin V (ColV) virulence-resistance plasmid pCERC3 [40] and contains virulence genes, including those for the aerobactin and Sit siderophore systems, in addition to multiple drug resistance genes (Fig. S3), but lacks the genes for ColV.

### Variation within individual plasmid types, with diverse and complex resistance regions

In addition to the variety of different plasmid types, long-read sequence data allowed us to observe diversity amongst plasmids of the same type (Tables S4 and S5, Figs S1–S4). There was considerable genetic diversity in plasmids carrying the FII-2 replicon (Fig. 2). FII-2 plasmids pLAO44 (GenBank accession OP242233) and pLAO37 (OP242229) in ST69 and ST101 strains only contained FII-2 replicons, while FII-2 plasmid pLAO82 (OP242230) in an ST34 isolate carried an additional  $\theta$ -RNA replicon and plasmids pLAO100 (ST40, OP242240) and pLAO103 (ST457, OP242243) carried an additional FIB-10 replicon (Figs 2 and 3). Diversity also occurred in the resistance regions of these plasmids (Fig. 2 and Fig. S4). FII-2 plasmids carried multiple resistance genes, including various combinations of *bla*<sub>CTX-M-27</sub>, *bla*<sub>CTX-M-55</sub> and *mcr-3.4* (Figs 2 and 3). Plasmid types FII-2 (pLAO37), FII-2: $\theta$ -RNA (pLAO82) and FII-46:FIB-like (pLAO69, OP242232) incorporated resistance regions with an accumulation of multiple resistance genes and co-carriage of *mcr* and *bla*<sub>CTX-M</sub> genes (Fig. 3b). FII-2: $\theta$ -RNA, pLAO82, demonstrated even more complexity with further accumulation of resistance genes as it also carried an additional *mcr-3.4* located between copies of *IS26* and *ISKpn40* (Fig. 3). There is widespread global distribution of the resistance regions found




**Fig. 1.** An overview of plasmid diversity showing plasmid size and AMR gene carriage (a) and plasmid type within sequence type (b) for all plasmids identified.

in these FII-2 plasmids, which have been identified in multiple countries, some of which are not close to Laos (Table S6). We found no link between the ST of the isolate and the type of plasmid AMR genes carried by that ST (Fig. 3a).


**Co-carriage of *bla*<sub>CTX-M</sub> and *mcr* genes occurs in multiple resistance region configurations**

Three plasmids in this collection carried both *bla*<sub>CTX-M</sub> and *mcr* genes (Fig. 3b). All three *bla*<sub>CTX-M</sub>/*mcr* co-carriage plasmids were F-types, but they differed in size and plasmid replicon type. The 98 237 bp plasmid, pLAO37, only carried an FII-2 replicon and was found in five ST101 *E. coli* isolates. The 110 949 bp pLAO82 was a multi-replicon co-integrate plasmid carrying both FII-2 and θ-RNA replicons and was found in one ST34 *E. coli* isolate. The 105 425 bp plasmid, pLAO69, carried FII-46 and FIB-20 replicons, and was found in two ST10 *E. coli* isolates. All three of these plasmids have typical F-type plasmid backbones that include




**Fig. 2.** Diversity within FII-2 plasmid replicon group showing variable plasmid sizes and resistance genes. (a) Annotated FII-2 plasmid group maps indicate the location of *bla<sub>CTX-M</sub>* and *mcr* genes. Maroon arrows represent Prokka-annotated genes. All other brightly coloured arrows represent antibiotic resistance genes. Transposable elements (TEs) are displayed as brightly coloured boxes. Notable elements are IS26 (bright green), *ISEcp1* (bright pink), Tn1721 (pale blue) and ISKpn26 (lavender). The plasmid transfer (*tra*) region is highlighted with an orange outline. (b) IQ-TREE of FII-2 plasmids core alignment showing AMR genes and gene/presence absence matrix. The participant (Pt) and ST of the isolate from which the plasmid was identified are displayed in different colours. Resistance genes identified using Abricate (grey, present) are shown alongside the gene presence/absence profile from Panaroo (royal blue, present). Forty-six core genes were classified by Panaroo in the phylogeny.

(a)




(b)


i) pLAO37



ii) pLAO82



iii) pLAO69



**Fig. 3.** Antimicrobial resistance gene carriage diversity by (a) ST and (b) within *mcr* and *bla*<sub>CTX-M</sub> co-carriage plasmids. All maroon genes are prokka-annotated genes. All other brightly coloured genes are antibiotic resistance genes. Transposable elements (e.g. transposons, insertion sequences) are displayed as brightly coloured boxes. The plasmid transfer (*tra*) region is highlighted with an orange outline.



genes for replication, stable maintenance, conjugative transfer and establishment in new hosts [40, 41]. Each plasmid contained a complete and uninterrupted transfer region, suggesting that all three have the capacity for self-mediated conjugation [42].

Detailed comparison of the FII-2 plasmids pLAO37 and pLAO82 showed that the backbones are almost identical apart from a recombination patch (approximately 7 kb). pLAO37 and pLAO82 have the same FII-2 *repA1* gene. The AMR genes in both pLAO37 and pLAO82 are located in a complex region bounded by IS26 at the left end and Tn1721 at the right end (Fig. 3b). These regions comprise sequences from multiple mobile genetic elements with distinct origins and include genes that confer resistance to beta-lactams (*bla*<sub>CTX-M</sub>), colistin (*mcr-3.4*), aminoglycosides (*aacC2d*), chloramphenicol (*catA1*) and quinolones (*qnrS1*). An additional  $\theta$ -RNA replicon in pLAO82 is part of a small plasmid that has been captured and incorporated into the resistance region. pLAO37 and pLAO82 contain an extra partial Tn21 and partial IS26 region inserted between the *tmrB* and *mcr* genes (Fig. 3b). The pLAO82 resistance region also includes an additional copy of *mcr-3.4*. The *mcr-3.4* in pLAO37 was truncated by ISKpn26, which removed the terminal 32 bp of the gene. This configuration of ISKpn26 and *mcr-3.4* is not present in any other sequence deposited in the GenBank non-redundant nucleotide database. A novel transposon, Tn7514, was present in both pLAO37 and pLAO82 but was inserted in two different backbone locations (Fig. 2b and Tables S7 and S8) [43].

Although largely syntenic, the nucleotide identity of the pLAO69 backbone differs significantly from that of pLAO37 and pLAO82, consistent with its distinct replicon type. The FII-46 *repA1* gene of pLAO69 is only 94% identical to FII-2 *repA1* of pLAO37 and pLAO82. The pLAO69 transfer region also differed significantly, matching only 89% of pLAO37 and pLAO82 transfer region with an overall identity of 97.2% in a BLASTn comparison. pLAO69 carried *mcr-3.1*, which differs from the *mcr-3.4* found in pLAO37 and pLAO82 (Fig. 3b). However, inspection of the resistance region in pLAO69 that contains *mcr-3.1* revealed that it is largely composed of the same confluence of mobile elements found in the resistance regions of pLAO37 and pLAO82 (Fig. 3b). In addition to their identities, the configuration of these elements was the same in pLAO69 and pLAO37/pLAO82.

### Insertion sequence type, abundance and diversity are associated with AMR gene carriage

The vast majority ( $n=36$ ) of the 40 AMR plasmids contained one or more IS. Three exceptions were plasmids that contained AMR genes found in association with gene cassettes [*bla*<sub>VEB</sub> in pLAO60 and *aadA2*, *ant(3'')*-Ia, *cmlA1*, *dfrA12* in pLAO78], or miniature inverted-repeat tandem elements (MITEs) [*tet(A)* in pLAO84]. The remaining plasmid, pLAO41, contained *mcr-1* but no detectable IS elements.


AMR plasmids carried almost three times more unique families of IS than their non-AMR counterparts (Fig. 4a), with an average of 6.75 unique IS families in AMR plasmids (range 1–11), vs 2.4 in non-AMR plasmids (range 1–8).

AMR plasmids contained an abundance of IS6/26 and IS1-family elements. Of the 36 AMR plasmids that did carry IS elements, all but one carried at least one IS6-family element (Fig. 4a), and the IS1 family was present in 83% ( $n=30$ ). IS3 and IS5 were also common, present in 75% ( $n=27$ ) and 78% ( $n=28$ ) of AMR plasmids, respectively. Differences in IS6 family element carriage were seen between AMR and non-AMR plasmids, with AMR plasmids containing a mean of five IS6 family elements per plasmid (range 0–9). In contrast, only 24% ( $n=4$ ) of non-AMR plasmids carried IS6 family elements, with only 1–2 IS6 elements per plasmid (Fig. 4a). IS6 family elements were present in all plasmids that co-carry *bla*<sub>CTX-M</sub> and *mcr* genes (Fig. 4b), where they were the most prevalent IS family. There was no clear association between IS families and co-carriage of the *mcr* and *bla*<sub>CTX-M</sub> genes (Fig. 4b). In plasmids where there was co-carriage of *mcr* and *bla*<sub>CTX-M</sub> genes, IS6 and IS3 were the most abundant IS families, with all co-carriage plasmids additionally carrying IS1380, IS1 and IS4. In each co-carriage plasmid, more than six different IS families were identified, and of the IS identified between 30–46% were of the IS6 family (Fig. 4b), with IS26 the most prominent element.

## DISCUSSION

In this study we conducted an in-depth investigation of the role of plasmids in the alarmingly high levels of AMR found in *E. coli* that colonized the gastrointestinal tract of travellers to Laos [7]. We have revealed an enormous diversity in the plasmids of this *E. coli* population, particularly in their resistance regions, many of which contained multiple AMR genes. Concerningly, the majority of the 40 AMR plasmids ( $n=30$ , 75%) contained ESBL genes, a colistin resistance gene or both. Our data showed the abundance and importance of F-type, X-type, Q-type and ColE1-like plasmids as vectors for AMR gene spread in *E. coli* in Laos. AMR plasmids accounted for 38.1% ( $n=40$ ) of the 105 distinct plasmids and 17 of these 40 AMR plasmids identified were F-type (42.5%), highlighting their importance as carriers of AMR genes (including ESBLs, *mcr*). F-type plasmids are known to carry AMR genes [44] and are an important factor in the high incidence of AMR carriage in this study. Although four types dominated, the AMR plasmid population identified in travellers to this region of Laos was extremely diverse, with 29 different plasmid types, including phage and virulence plasmids, found to carry AMR genes. Both AMR and non-AMR plasmids were identified throughout the study, including several from baseline faecal samples [7], where plasmids may have been acquired from travel to Laos or potentially from other travel.

A variety of less anticipated vehicles for AMR genes were identified in these Lao *E. coli*. Small ColE1-like  $\theta$ -RNA plasmids are common in *E. coli* [21, 45] and their high prevalence would be expected. These  $\theta$ -RNA ( $n=26$ ) were the most prevalent



**Fig. 4.** IS families present in (a) all plasmid types displaying AMR gene presence/absence and IS linked to (b) *mcr* and *bla*<sub>CTX-M</sub> co-carriage plasmids.

plasmids in the collection, followed by F-types ( $n=23$ ) (Fig. 1), but in contrast to F-type plasmids, only a small number of  $\theta$ -RNA plasmids ( $n=3$ ) carried AMR genes. This is consistent with previous findings that ColE1-like plasmids occasionally carry resistance genes, including aminoglycoside and beta-lactam resistance determinants [46]. While relatively uncommon, the importance of small, high-copy-number plasmids for AMR should not be underestimated, as these can serve as platforms for the evolution of new resistance phenotypes [46, 47]. The *tet(A)*-carrying ColE1-like plasmid pLAO84 clearly demonstrates the capacity these small plasmids have for disseminating resistance genes. pLAO84 was found in multiple *E. coli* STs in this collection, suggesting that it was circulating in the local *E. coli* population before being acquired by multiple study participants (Fig. S1). Additionally, the presence of a plasmid that was almost identical to pLAO84 in GenBank (GenBank accession CP057097.1) indicates that this ColE1-like plasmid lineage has already spread internationally, as it was present in *Escherichia fergusonii* isolated from pig faeces in the UK.

A Y-type phage-plasmid (pLAO59, OP242238, File S8) highlighted an additional opportunity for AMR spread, since there is a chance for co-selection of this MDR plasmid due to carriage of metal resistance genes, e.g. for silver and copper. pLAO59 appeared not only to have lost key phage genes required for the lytic lifestyle [48], apparently as a result of deletions by insertion sequences, but carried multiple AMR genes and heavy metal resistance genes. Co-resistance and cross-resistance can cause co-selection of bacteria carrying metal resistance and AMR genes, with the metal resistance gene causing maintenance of the AMR gene [49, 50]. In Vientiane, Laos it has been reported that environmental samples sourced near municipal solid waste landfill showed heavy contamination with heavy metals, including copper at levels higher than WHO permissible standards [51], indicating the possibility that co-selection by metal resistance genes is a real environmental pressure in the Vientiane area from which these isolates were collected. Silver and copper can co-select for various AMR genes in *E. coli*, including tetracycline and sulphonamide resistance genes, which were also found in pLAO59 [52, 53].

Multiple combinations of AMR resistance genes were found in multiple genetic contexts associated with various mobile genetic elements. *mcr* genes, for example, were found next to multiple TEs (Fig. 3, Table S4) and mainly on F-type plasmids, consistent with the literature indicating that *mcr-3.4* is associated with the FII-type [54, 55]. *ISEcp1* is known to contribute to the spread of *bla*<sub>CTX-M</sub> [56], but interestingly, only one complete *ISEcp1* was identified (pLAO32) as part of a *bla*<sub>CTX-M-55</sub>-containing transposition unit flanked by the 5 bp target site duplication TAACA. Most *bla*<sub>CTX-M</sub> genes in this collection were associated with complete copies of IS26 and partial *ISEcp1* (Figs 2–4, Figs S3 and S4). IS26 from the IS6/26 family is known to play a key role in AMR gene dissemination [57–60]. IS26 carriage predisposes plasmids to insertion of additional IS26 and any associated AMR genes, which facilitates the accumulation of AMR genes at single sites, leading to multidrug resistance phenotypes [57, 61]. This appears to be the case with pLAO37, pLAO82 and pLAO69 featuring IS26 and, based on what is known of IS26 behaviour [57], IS26 is likely to have played an important role in the assembly of their complex co-carriage resistance regions (Fig. 3).

Analysis of plasmid contigs from 49 *E. coli* hybrid assemblies has confirmed the vast and diverse genetic context of AMR in the Kantele *et al.* dataset isolated in Laos, and highlighted that in addition to multiple unique colonizing strains [7], there are multiple distinct plasmids present in this dataset. Long-read sequencing has been critical in highlighting the key role insertion sequence elements may play in the formation of this complex set of MDR plasmids that risk spreading AMR. This previously unreported cohort of MDR plasmids offers the alarming prospect that one of these will create a stable configuration for the creation of a successful pandemic MDR plasmid.

#### Funding information

AS was funded by the Wellcome Antimicrobial and Antimicrobial Resistance (AAMR) DTP (108876B15Z). The Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit is funded by the Wellcome Trust (grant number 106698/Z/14/Z). AK was supported by the Finnish Governmental Subsidy for Health Science Research, the Sigrid Jusélius Foundation, and the Academy of Finland (grant number 346127). JC was funded by the ERC grant no. 742158. RM is funded by Medical Research Council grant awarded to AM (MRS0136601). SJD was funded by Biotechnology and Biological Sciences Research Council grant awarded to AM (BBR0062611).

#### Acknowledgements

We are very grateful to the volunteers and the staff of the Microbiology Laboratory, Mahosot Hospital, Vientiane, Laos, who processed the samples and for the support of the directors and staff of Mahosot Hospital and the Ministry of Health, Lao PDR.

#### Author contribution

A.M. and J.C. conceived the idea for this analysis. A.K., E.K., D.A.B.D., P.N.P. and V.D. collected samples, isolated the strains and provided the isolates and epidemiological data in Laos. A.S. and S.D. performed long-read genomic sequencing and sequence processing. A.S., S.D. and R.M. performed genomic analysis, data interpretation and data visualization. A.S., R.M. and S.D. drafted the article. A.M. discussed results and edited the draft. J.C., A.K., P.N. and D.D. critically reviewed the draft. All authors read and approved the final manuscript.

#### Conflicts of interest

The authors declare that there are no conflicts of interest.

#### Ethical statement

The study protocol (see Appendix 1 of reference [7]) was approved by the Ethics Committee of the Helsinki University Hospital, Helsinki, Finland, and the Ethics Committee of Northwest and Central Switzerland, Basel, Switzerland. All participants provided written informed consent.

#### References

1. Cosgrove SE. The relationship between antimicrobial resistance and patient outcomes: mortality, length of hospital stay, and health care costs. *Clin Infect Dis* 2006;42 Suppl 2:S82–9.
2. Arcilla MS, Van Hattem JM, Bootsma MCJ, van Genderen PJJ, Goorhuis A, *et al.* Prevalence and risk factors for carriage of ESBL-producing *Enterobacteriaceae* in a population of Dutch travellers: a cross-sectional study. *Travel Med Infect Dis* 2020;33:101547.
3. Bevan ER, McNally A, Thomas CM, Piddock LJV, Hawkey PM. Acquisition and loss of CTX-M-producing and non-producing *Escherichia coli* in the fecal microbiome of travelers to South Asia. *mBio* 2018;9:e02408-18.
4. Bokhary H, Pangesti KNA, Rashid H, Abd El Ghany M, Hill-Cawthorne GA. Travel-related antimicrobial resistance: a systematic review. *Trop Med Infect Dis* 2021;6:11.
5. Woerther PL, Andremont A, Kantele A. Travel-acquired ESBL-producing *Enterobacteriaceae*: impact of colonization at individual and community level. *J Travel Med* 2017;24:S29–S34.
6. Arcilla MS, van Hattem JM, Haverkate MR, Bootsma MCJ, van Genderen PJJ, *et al.* Import and spread of extended-spectrum

- $\beta$ -lactamase-producing *Enterobacteriaceae* by international travellers (COMBAT study): a prospective, multicentre cohort study. *Lancet Infect Dis* 2017;17:78–85.
7. Kantele A, Kuenzli E, Dunn SJ, Dance DAB, Newton PN, et al. Dynamics of intestinal multidrug-resistant bacteria colonisation contracted by visitors to a high-endemic setting: a prospective, daily, real-time sampling study. *Lancet Microbe* 2021;2:e151–e158.
  8. Chaudhuri RR, Henderson IR. The evolution of the *Escherichia coli* phylogeny. *Infect Genet Evol* 2012;12:214–226.
  9. Rawat D, Nair D. Extended-spectrum  $\beta$ -lactamases in Gram negative bacteria. *J Glob Infect Dis* 2010;2:263–274.
  10. Stoesser N, Sheppard AE, Pankhurst L, De Maio N, Moore CE, et al. Evolutionary history of the global emergence of the *Escherichia coli* epidemic clone ST131. *mBio* 2016;7:e02162.
  11. Petty NK, Ben Zakour NL, Stanton-Cook M, Skippington E, Totsika M, et al. Global dissemination of a multidrug resistant *Escherichia coli* clone. *Proc Natl Acad Sci* 2014;111:5694–5699.
  12. Organisation WH. WHO Model List of Essential Medicines; 2017
  13. Meletis G. Carbapenem resistance: overview of the problem and future perspectives. *Ther Adv Infect Dis* 2016;3:15–21.
  14. Gharabeh MH, Shatnawi SQ. An overview of colistin resistance, mobilized colistin resistance genes dissemination, global responses, and the alternatives to colistin: a review. *Vet World* 2019;12:1735–1746.
  15. Matamoros S, van Hattem JM, Arcilla MS, Willemse N, Melles DC, et al. Global phylogenetic analysis of *Escherichia coli* and plasmids carrying the *mcr-1* gene indicates bacterial diversity but plasmid restriction. *Sci Rep* 2017;7:15364.
  16. Nikaido H. Multidrug resistance in bacteria. *Annu Rev Biochem* 2009;78:119–146.
  17. Aghapour Z, Gholizadeh P, Ganbarov K, Bialvaei AZ, Mahmood SS, et al. Molecular mechanisms related to colistin resistance in *Enterobacteriaceae*. *Infect Drug Resist* 2019;12:965–975.
  18. Furniss RCD, Kostrzewa M, Mavridou DAI, Larrouy-Maumus G. The clue is in the lipid A: rapid detection of colistin resistance. *PLoS Pathog* 2020;16:e1008331.
  19. Huddleston JR. Horizontal gene transfer in the human gastrointestinal tract: potential spread of antibiotic resistance genes. *Infect Drug Resist* 2014;7:167–176.
  20. Partridge SR, Kwong SM, Firth N, Jensen SO. Mobile genetic elements associated with antimicrobial resistance. *Clin Microbiol Rev* 2018;31:e00088-17.
  21. Rozwandowicz M, Brouwer MSM, Fischer J, Wagenaar JA, Gonzalez-Zorn B, et al. Plasmids carrying antimicrobial resistance genes in *Enterobacteriaceae*. *J Antimicrob Chemother* 2018;73:1121–1137.
  22. Hu Y, Moran RA, Blackwell GA, McNally A, Zong Z. Fine-scale reconstruction of the evolution of FII-33 multidrug resistance plasmids enables high-resolution genomic surveillance. *mSystems* 2022;7:e0083121.
  23. León-Sampedro R, DelaFuente J, Díaz-Agero C, Crellen T, Musicha P, et al. Pervasive transmission of a carbapenem resistance plasmid in the gut microbiota of hospitalized patients. *Nat Microbiol* 2021;6:606–616.
  24. Harmer CJ. HI1 and I1 resistance plasmids from *Salmonella enterica* serovar *Typhimurium* strain SRC27 are epidemic. *Microb Drug Resist* 2021;27:1495–1504.
  25. Carattoli A, Hasman H. Plasmidfinder and in silico pMLST: identification and typing of plasmid replicons in Whole-Genome Sequencing (WGS). *Methods Mol Biol* 2020;2075:285–294.
  26. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, et al. ResFinder 4.0 for predictions of phenotypes from genotypes. *J Antimicrob Chemother* 2020;75:3491–3500.
  27. Prijbelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes de novo assembler. *Curr Protoc Bioinformatics* 2020;70:e102.
  28. De Coster W, D'Hert S, Schultz DT, Cruys M, Van Broeckhoven C. NanoPack: visualizing and processing long-read sequencing data. *Bioinformatics* 2018;34:2666–2669.
  29. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 2017;13:e1005595.
  30. Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of *de novo* genome assemblies. *Bioinformatics* 2015;31:3350–3352.
  31. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat graphs. *Nat Biotechnol* 2019;37:540–546.
  32. Borowiak M, Fischer J, Hammerl JA, Hendriksen RS, Szabo I, et al. Identification of a novel transposon-associated phosphoethanolamine transferase gene, *mcr-5*, conferring colistin resistance in d-tartrate fermenting *Salmonella enterica* subsp. *enterica* serovar Paratyphi B. *J Antimicrob Chemother* 2017;72:3317–3324.
  33. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014;30:2068–2069.
  34. Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res* 2018;3:124.
  35. Villa L, García-Fernández A, Fortini D, Carattoli A. Replicon sequence typing of IncF plasmids carrying virulence and resistance determinants. *J Antimicrob Chemother* 2010;65:2518–2529.
  36. Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. ISfinder: the reference centre for bacterial insertion sequences. *Nucleic Acids Res* 2006;34:D32–6.
  37. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 2015;32:268–274.
  38. Tonkin-Hill G, MacAlasdair N, Ruis C, Weimann A, Horesh G, et al. Producing polished prokaryotic pangenomes with the Panaroo pipeline. *Genome Biol* 2020;21:180.
  39. Xie Z, Tang H. ISEScan: automated identification of insertion sequence elements in prokaryotic genomes. *Bioinformatics* 2017;33:3340–3347.
  40. Moran RA, Holt KE, Hall RM. pCERC3 from a commensal ST95 *Escherichia coli*: a ColV virulence-multiresistance plasmid carrying a *sul3*-associated class 1 integron. *Plasmid* 2016;84–85:11–19.
  41. Cox KEL, Schildbach JF. Sequence of the R1 plasmid and comparison to F and R100. *Plasmid* 2017;91:53–60.
  42. Frost LS, Leplae R, Summers AO, Toussaint A. Mobile genetic elements: the agents of open source evolution. *Nat Rev Microbiol* 2005;3:722–732.
  43. Tansirichaiya S, Rahman MA, Roberts AP. The transposon registry. *Mob DNA* 2019;10:40.
  44. Matlock W, Chau KK, AbuOun M, Stubberfield E, Barker L, et al. Genomic network analysis of environmental and livestock F-type plasmid populations. *ISME J* 2021;15:2322–2335.
  45. Riley MA, Gordon DM. A survey of Col plasmids in natural isolates of *Escherichia coli* and an investigation into the stability of Col-plasmid lineages. *J Gen Microbiol* 1992;138:1345–1352.
  46. Ares-Arroyo M, Rocha EPC, Gonzalez-Zorn B. Evolution of ColE1-like plasmids across  $\gamma$ -Proteobacteria: from bacteriocin production to antimicrobial resistance. *PLoS Genet* 2021;17:e1009919.
  47. Moran RA, Hall RM. Analysis of pCERC7, a small antibiotic resistance plasmid from a commensal ST131 *Escherichia coli*, defines a diverse group of plasmids that include various segments adjacent to a multimer resolution site and encode the same NikA relaxase accessory protein enabling mobilisation. *Plasmid* 2017;89:42–48.
  48. Łobocka MB, Rose DJ, Plunkett G 3rd, Rusin M, Samojedny A, et al. Genome of bacteriophage P1. *J Bacteriol* 2004;186:7032–7068.
  49. Pal C, Asiani K, Arya S, Resning C, Stekel DJ, et al. Metal resistance and its association with antibiotic resistance. *Adv Microb Physiol* 2017;70:261–313.
  50. Cheng G, Ning J, Ahmed S, Huang J, Ullah R, et al. Selection and dissemination of antimicrobial resistance in Agri-food production. *Antimicrob Resist Infect Control* 2019;8:158.
  51. Vongdala N, Tran HD, Xuan TD, Teschke R, Khanh TD. Heavy metal accumulation in water, soil, and plants of municipal solid waste landfill in Vientiane, Laos. *Int J Environ Res Public Health* 2018;16:22.

52. Yang H, Wei SH, Hobman JL, Dodd CER. Antibiotic and metal resistance in *Escherichia coli* isolated from pig Slaughterhouses in the United Kingdom. *Antibiotics* 2020;9:746.
53. Falgenhauer L, Ghosh H, Guerra B, Yao Y, Fritzenwanker M, et al. Comparative genome analysis of IncHI2 VIM-1 carbapenemase-encoding plasmids of *Escherichia coli* and *Salmonella enterica* isolated from a livestock farm in Germany. *Vet Microbiol* 2017;200:114–117.
54. Ahmed S, Olsen JE, Herrero-Fresno A, Moreno-Hagelsieb G. The genetic diversity of commensal *Escherichia coli* strains isolated from non-antimicrobial treated pigs varies according to age group. *PLoS One* 2017;12:e0178623.
55. Yamaguchi T, Kawahara R, Harada K, Teruya S, Nakayama T, et al. The presence of colistin resistance gene mcr-1 and -3 in ESBL producing *Escherichia coli* isolated from food in Ho Chi Minh City, Vietnam. *FEMS Microbiol Lett* 2018;365:11.
56. Rossolini GM, D'Andrea MM, Mugnaioli C. The spread of CTX-M-type extended-spectrum beta-lactamases. *Clin Microbiol Infect* 2008;14 Suppl 1:33–41.
57. Harmer CJ, Moran RA, Hall RM. Movement of IS26-associated antibiotic resistance genes occurs via a translocatable unit that includes a single IS26 and preferentially inserts adjacent to another IS26. *mBio* 2014;5:e01801–14.
58. Ford PJ, Avison MB. Evolutionary mapping of the SHV beta-lactamase and evidence for two separate IS26-dependent blaSHV mobilization events from the *Klebsiella pneumoniae* chromosome. *J Antimicrob Chemother* 2004;54:69–75.
59. Hordijk J, Bosman AB, van Essen-Zandbergen A, Veldman K, Dierikx C, et al. qnrB19 gene bracketed by IS26 on a 40-kilobase IncR plasmid from an *Escherichia coli* isolate from a veal calf. *Antimicrob Agents Chemother* 2011;55:453–454.
60. Post V, Hall RM. AbaR5, a large multiple-antibiotic resistance region found in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2009;53:2667–2671.
61. Harmer CJ, Hall RM. IS26-mediated formation of transposons carrying antibiotic resistance genes. *mSphere* 2016;1:e00038-16.

#### Five reasons to publish your next article with a Microbiology Society journal

1. When you submit to our journals, you are supporting Society activities for your community.
2. Experience a fair, transparent process and critical, constructive review.
3. If you are at a Publish and Read institution, you'll enjoy the benefits of Open Access across our journal portfolio.
4. Author feedback says our Editors are 'thorough and fair' and 'patient and caring'.
5. Increase your reach and impact and share your research more widely.

Find out more and submit your article at [microbiologyresearch.org](https://microbiologyresearch.org).