

Meta-DiSc 2.0

Plana, Maria N.; Arevalo-Rodriguez, Ingrid; Fernández-García, Silvia; Soto, Javier; Fabregate, Martin; Pérez, Teresa; Roqué, Marta; Zamora, Javier

DOI:

[10.1186/s12874-022-01788-2](https://doi.org/10.1186/s12874-022-01788-2)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Plana, MN, Arevalo-Rodriguez, I, Fernández-García, S, Soto, J, Fabregate, M, Pérez, T, Roqué, M & Zamora, J 2022, 'Meta-DiSc 2.0: a web application for meta-analysis of diagnostic test accuracy data', *BMC Medical Research Methodology*, vol. 22, no. 1, 306. <https://doi.org/10.1186/s12874-022-01788-2>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

SOFTWARE

Open Access



Meta-DiSc 2.0: a web application for meta-analysis of diagnostic test accuracy data

Maria N. Plana^{1,2*} , Ingrid Arevalo-Rodriguez^{2,3}, Silvia Fernández-García³, Javier Soto⁴, Martin Fabregate⁵, Teresa Pérez^{6,7}, Marta Roqué^{2,8} and Javier Zamora^{2,3,9}

Abstract

Background: Diagnostic evidence of the accuracy of a test for identifying a target condition of interest can be estimated using systematic approaches following standardized methodologies. Statistical methods for the meta-analysis of diagnostic test accuracy (DTA) studies are relatively complex, presenting a challenge for reviewers without extensive statistical expertise. In 2006, we developed Meta-DiSc, a free user-friendly software to perform test accuracy meta-analysis. This statistical program is now widely used for performing DTA meta-analyses. We aimed to build a new version of the Meta-DiSc software to include statistical methods based on hierarchical models and an enhanced web-based interface to improve user experience.

Results: In this article, we present the updated version, Meta-DiSc 2.0, a web-based application developed using the R Shiny package. This new version implements recommended state-of-the-art statistical models to overcome the limitations of the statistical approaches included in the previous version. Meta-DiSc 2.0 performs statistical analyses of DTA reviews using a bivariate random effects model. The application offers a thorough analysis of heterogeneity, calculating logit variance estimates of sensitivity and specificity, the bivariate I-squared, the area of the 95% prediction ellipse, and the median odds ratios for sensitivity and specificity, and facilitating subgroup and meta-regression analyses. Furthermore, univariate random effects models can be applied to meta-analyses with few studies or with non-convergent bivariate models.

The application interface has an intuitive design set out in four main menus: file upload; graphical description (forest and ROC plane plots); meta-analysis (pooling of sensitivity and specificity, estimation of likelihood ratios and diagnostic odds ratio, sROC curve); and summary of findings (impact of test through downstream consequences in a hypothetical population with a given prevalence).

All computational algorithms have been validated in several real datasets by comparing results obtained with STATA/SAS and MetaDTA packages.

Conclusion: We have developed and validated an updated version of the Meta-DiSc software that is more accessible and statistically sound. The web application is freely available at www.metadisc.es.

Keywords: Systematic review, Meta-analysis, Sensitivity, Specificity, Diagnosis, Heterogeneity

*Correspondence: nieves.plana@salud.madrid.org

¹ Health Technology Assessment Unit, Hospital Universitario Ramón Y Cajal, IRYCIS, Madrid, Spain

Full list of author information is available at the end of the article

Background

The evaluation of the role and properties of diagnostic tools has become a priority for global health policy and decision-making, driven mainly by the development



of new technologies for well-known diseases and the emergence of new deleterious conditions affecting large-scale populations [1, 2]. Diagnostic evidence of the accuracy of a test for detecting a target condition of interest can be appraised using systematic approaches following standardized methodologies [3]. Briefly, diagnostic studies focus on estimating the ability of the index tool to identify subjects with or without the condition of interest [3]; evidence synthesis then requires two quantities: test sensitivity and specificity and the correlation between them [4]. The statistical approach used depends on the choice between estimating accuracy for a common threshold (i.e. an average operating point), or an expected curve across many thresholds (i.e. a summary ROC curve) [5–7], using commercial software packages with the analytical characteristics needed for fitting complex hierarchical models.

We recently found that the statistical synthesis of accuracy data was one of the methods more frequently omitted during the development of rapid reviews of diagnostic tests [8]. This, then, would be a potential bottleneck for the extended evaluation of diagnostic tools [8]. For several years, Meta-DiSc software has been one of the most widely used statistical programs in the meta-analysis of diagnostic data, with more than 1300 citations in peer-reviewed scientific articles [9]. It is a freely available, easy-to-use tool, that enables reviewers to apply statistical methods for the meta-analysis of diagnostic test accuracy (DTA) within an evidence synthesis framework. This software implemented the statistical methods recommended during its development, including the linear model proposed by Littenberg and Moses, and the univariate I-squared index to quantify heterogeneity. Hierarchical models are currently the method of choice for overcoming the limitations of previous statistical approaches [3]. These methodological developments have prompted us to update Meta-DiSc to include current statistical methods for the meta-analysis of test accuracy systematic reviews and an enhanced web-based interface to improve user experience. Our objective was to develop a new version of the Meta-DiSc software as a web application (app) to summarize DTA results by applying statistical methods based on hierarchical models.

Implementation

We have developed a web-based app using R Shiny software. Shiny can be used to build R-based interactive applications directly on RStudio, the integrated development environment for R. The application has been deployed using the shinyapps.io platform.

Estimating pooled diagnostic accuracy indices

The app performs statistical analysis of DTA reviews using a bivariate random effects model [5] and the *glmer* function of the *lme4* package [10] for fitting a generalized linear mixed effect model. Summary points (average sensitivity and specificity) and the parameters are derived to depict the sROC curve. Positive and negative likelihood ratios (LR) and the diagnostic odds ratio (DOR) estimates are obtained from model parameters. The Delta method as implemented in the *msm* package [11] is used to compute the standard error of the estimates parameters. Forest plots and ROC plots have been implemented using the functionalities of the *meta*, *ggplot2* and *plotly* packages [12–14].

The program also offers the possibility of using a univariate random effects model. Although separate pooling is not recommended for DTA meta-analysis since it fails to account for the correlation between sensitivity and specificity, we have included this option because univariate models, in some instances, have a role in DTA reviews. This is the case, for example, when it is difficult to estimate all parameters of a bivariate model or when the focus of the analysis is only on one of the accuracy indices (i.e. sensitivity or specificity) [15].

Quantifying heterogeneity

Meta-DiSc 2.0 implements a thorough analysis of heterogeneity. In addition to the estimates of logit variances of sensitivity and specificity [16], the software calculates a bivariate I-squared index [17], the area of the 95% prediction ellipse using the *polyarea* function of the *pracma* package [18], and finally, the median odds ratios for sensitivity and specificity [16].

Exploring heterogeneity: subgroup and meta-regression analyses

The app can be used to perform subgroup and meta-regression analysis. For this purpose, additional columns need to be included in the dataset to define dichotomous covariates (one each time), which will be used to split the dataset and obtain the accuracy estimates for each subgroup. Exploring these individual results gives the reviewer insights into the between-group difference in sensitivity and specificity and the between-study variances in both indices. The meta-regression option compares the accuracy estimates obtained for these subgroups (i.e., sensitivity and specificity) [19]. The bivariate model includes interaction terms with both sensitivity and specificity and compares the statistical significance of these effects using the *lmtree* package [20]. For simplicity, meta-regression analysis implemented in Meta-DiSc 2.0 assumes that between-study variances are equal.

Therefore, authors should check how appropriate is this assumption by comparing the between study variances in each subgroup.

Results

Meta-DiSc 2.0 is freely available from www.metadisc.es. The user interface design is intuitive and easy-to-use. The left lateral panel organizes the workspace in four main menus: *File upload*, *Graphical description*, *Meta-analysis*, and *Summary of findings*. The app also includes a short user-guide video to show the practical use of the application.

File upload menu

The app can import data as either comma-delimited (i.e.,.csv) or Excel files (.xlsx files). The file must include data from 2 × 2 tables of individual studies in four columns named TP, FP, FN, TN, representing the number of true positives, false positives, false negatives and true negatives, respectively. The file must also include a unique identifier for each study (ID). It may also incorporate additional columns that will be considered as covariates to explore sources of between-study variability (Fig. 1). Figs. 2, 3, 4 and 5 show different app screens for the analysis of a published diagnostic accuracy systematic review on pulse oximetry screening for a critical congenital heart defects dataset [21].

Graphical description menu

The app generates forest plots of sensitivity and specificity of individual studies to evaluate heterogeneity graphically. Studies of the forest plots are ordered in the same way as defined in the uploaded file. The ROC plot represents individual sensitivity and specificity, and offers the option of adding error bars, as either horizontal or vertical lines. This graphical description can be presented by subgroups defined by the covariates included in the file. All figures are downloadable as.png and.svg formats.

Meta-analysis menu

All analyses are obtained from the meta-analysis menu. The first option of this menu is to fit the bivariate model of sensitivity and specificity. Results are shown in the corresponding tabs: i) statistics, ii) sROC curve, iii) subgroup analysis, and iv) sensitivity analysis.

In the statistics tab, users will find the pooled accuracy estimates (sensitivity and specificity, positive and negative likelihood ratios, diagnostic odds ratio and false-positive rate) along with their corresponding 95% CI (Fig. 2A). Additionally, the app provides model parameters estimates (logit sensitivity, logit specificity, standard errors, logits variances, covariance and correlation), which can be easily transferred to the Cochrane Review manager system (RevMan [22]) (Fig. 2B). Finally, the app shows the heterogeneity statistics, including variances of the logit sensitivity and specificity along with corresponding





Fig. 1 File upload menu



median odds ratios (MOR) [16], bivariate I-squared [17], and the area of 95% prediction ellipse [16] (Fig. 2C).

After visualizing these numerical results, users can obtain graphical summary results by moving to the next tab named sROC curve (Fig. 3), where the ROC plane graphic can be visualized and downloaded. Different display options can be selected or omitted, e.g., summary point, confidence and prediction ellipses, summary ROC curve, and individual study results.

The subgroup analysis tab fits a new bivariate model, including additional parameters to assess whether sensitivity and specificity differ between subgroups. After showing the coefficients of the estimated model, a formal comparison between subgroups can be made using the meta-regression tab. The app shows the relative sensitivity and specificity along with 95% confidence intervals (LCI and UCI) and p-values of likelihood ratio tests to compare the subgroups formed according to the selected covariate (Fig. 4).

A final sensitivity analysis tab can be used to restrict the analysis to certain specific studies, by simply selecting the level of the dummy variable that will be employed as the inclusion criterion from the dropdown menu.


If two independent univariate analyses of sensitivity and specificity are selected, the results of both random effects models are displayed in a series of screens showing pooled estimates, heterogeneity statistics, and forest plots.

Summary of findings menu

To describe the absolute impact of a diagnostic test in a population with a given prevalence and fix a hypothetical sample size, the app calculates the number of false-positive and false-negative test results observed [23]. Users can download a figure that shows the outcomes (TP, FP, FN, TN) obtained (Fig. 5).

As a worked example, we have used a dataset that corresponds to a systematic review that evaluates the diagnostic accuracy of pulse oximetry as a screening method for detecting critical congenital heart defects (CCHD) in asymptomatic newborn infants [21]. The published meta-analysis included nineteen studies and was performed using the METADAS macro for SAS that uses Proc NLMIXED [24]. To assess potential sources of heterogeneity, we performed subgroup analyses and meta-regression. The overall sensitivity of pulse oximetry for the detection of CCHD was 76.3% (95% CI 69.5 to 82.0), while specificity was 99.9% (95% CI 99.7 to 99.9). We measured total between-study variability in sensitivity and specificity through variances of the random effects for logit(sensitivity) and logit(specificity), and their covariance. We also provided 95% confidence and prediction ellipses.

We replicated the published analysis using Meta-DiSc 2.0, extending the heterogeneity description to include the area of the 95% prediction ellipse [16], the median odds ratio for sensitivity and specificity [16] and I2 bivariate [17] (Fig. 2C). We also replicated the subgroup analysis for the covariate "test of timing" (within 24 h of birth vs after 24 h from birth) (Fig. 3). Summary estimates of sensitivity and specificity of studies that performed screening after 24 h were 73.6% (95% CI 62.8 to 82.1) and 99.9% (95% CI 99.9 to 100). For studies that performed screening within 24 h, summary estimates of sensitivity and specificity were 79.5% (95% CI 70.0 to 86.6) and 99.6% (95% CI 99.1 to 99.8). The relative specificity for the detection of CCHD was significantly higher when newborn pulse oximetry was performed more than 24 h after birth (Fig. 4). Validation of the analyses using Meta-DiSc 2.0 produced the same results as those obtained with METADAS macro [24]. The comparison of the numerical results obtained with Meta-DiSc 2.0 and the results obtained with other




Statistics

Summary statistics RevMan parameters **Metaregression**

Parameter	Estimate	LCI	UCI	p-value
Relative sensitivity level within 24h vs after 24h	1.08	0.905	1.29	0.366
Relative specificity level within 24h vs after 24h	0.996	0.993	1	0.001
Global test comparison				0.005

[Download](#)

Fig. 4 Metarregresion results (Meta-analysis menu)



software (METADAS in SAS [24], METANDI in Stata [25] and MetaDTA [26]) are shown in Table 1. We have further evaluated the app, replicating the analysis of four systematic reviews published in the literature [27–30] (Table 1).

Discussion

Our goal was to update a previous version of the Meta-DiSc software [9]. After this update, we are confident that MetaDiSc 2.0 can be in the league of available DTA meta-analysis software. The application unifies the main

Table 1 Metanalysis results using Meta-DiSc 2.0, METANDI (STATA), METADAS (SAS) and MetaDTA statistical software

First author, year	Software	Sensitivity (95% CI)	Specificity (95% CI)	LR + (95% CI)	LR- (95% CI)	DOR (95% CI)
Fahey et al. 1995 [27]	Meta-DiSc 2.0	0.66 (0.58; 0.73)	0.75 (0.68; 0.80)	2.58 (2.14; 3.11)	0.46 (0.39; 0.55)	5.59 (4.34; 7.19)
	METANDI	0.66 (0.58; 0.72)	0.75 (0.68; 0.80)	2.58 (2.14; 3.12)	0.46 (0.39; 0.55)	5.58 (4.33; 7.20)
	METADAS	0.66 (0.58; 0.72)	0.75 (0.68; 0.80)	2.58 (2.14; 3.12)	0.46 (0.39; 0.54)	5.59 (4.34; 7.20)
	MetaDTA	0.66 (0.58; 0.73)	0.75 (0.68; 0.80)	2.58 (2.14; 3.11)	0.46 (0.39; 0.55)	5.59 (4.34; 7.19)
Scheidler et al. 1997 [29]	Meta-DiSc 2.0	0.67 (0.61; 0.74)	0.84 (0.76; 0.89)	4.14 (2.70; 6.36)	0.39 (0.31; 0.49)	10.64 (5.83; 19.40)
	METANDI	0.67 (0.60; 0.74)	0.84 (0.76; 0.89)	4.15 (2.69; 6.40)	0.39 (0.31; 0.49)	10.65 (5.81; 19.54)
	METADAS	0.67 (0.60; 0.74)	0.84 (0.76; 0.89)	4.14 (2.69; 6.38)	0.39 (0.31; 0.49)	10.64 (5.82; 19.46)
	MetaDTA	0.67 (0.61; 0.74)	0.84 (0.76; 0.89)	4.14 (2.70; 6.36)	0.39 (0.31; 0.49)	10.64 (5.83; 19.40)
Honest et al. 2002 [28]	Meta-DiSc 2.0	0.58 (0.45; 0.71)	0.84 (0.78; 0.88)	3.57 (2.75; 4.63)	0.50 (0.38; 0.67)	7.14 (4.59; 11.13)
	METANDI	0.58 (0.44; 0.71)	0.84 (0.78; 0.88)	3.57 (2.75; 4.64)	0.50 (0.37; 0.67)	7.15 (4.57; 11.19)
	METADAS	0.57 (0.43; 0.70)	0.84 (0.77; 0.88)	3.49 (2.67; 4.56)	0.51 (0.38; 0.68)	6.82 (4.34; 10.71)
	MetaDTA	0.58 (0.45; 0.71)	0.84 (0.78; 0.88)	3.57 (2.75; 4.63)	0.50 (0.38; 0.67)	7.14 (4.59; 11.13)
Verde 2010 [30]	Meta-DiSc 2.0	0.96 (0.95; 0.97)	0.95 (0.94; 0.97)	20.61 (14.65; 29.0)	0.05 (0.04; 0.06)	452.28 (296.93; 688.90)
	METANDI	0.96 (0.95; 0.97)	0.95 (0.93; 0.97)	20.59 (14.59; 29.06)	0.05 (0.04; 0.06)	451.76 (295.52; 690.60)
	METADAS	0.96 (0.95; 0.97)	0.95 (0.94; 0.97)	20.61 (14.64; 29.01)	0.05 (0.04; 0.06)	452.28 (296.78; 689.26)
	MetaDTA	0.96 (0.95; 0.97)	0.95 (0.94; 0.97)	20.61 (14.65; 29.0)	0.05 (0.04; 0.06)	452.28 (296.93; 688.90)
Plana 2018 [21]	Meta-DiSc 2.0	0.76 (0.70; 0.82)	1.00 (1.00; 1.00)	535.19 (281.45; 1017.67)	0.24 (0.18; 0.31)	2257.34 (1183.08; 4307.05)
	METANDI	0.76 (0.70; 0.82)	1.00 (1.00; 1.00)	535.27 (279.68; 1024.40)	0.24 (0.18; 0.31)	2254.82 (1174.19; 4329.97)
	METADAS	0.76 (0.70; 0.82)	1.00 (1.00; 1.00)	535.55 (280.27; 1023.35)	0.24 (0.18; 0.31)	2258.72 (1179.04; 4327.09)
	MetaDTA	0.76 (0.70; 0.82)	1.00 (1.00; 1.00)	535.19 (281.45; 1017.67)	0.24 (0.18; 0.31)	2257.34 (1183.08; 4307.05)

standard routines for diagnostic accuracy meta-analysis and prevents reviewers from choosing among the variety of R packages available for this purpose, since not all of them have the currently recommended methods for DTA meta-analysis. Additionally, novel reviewers using MetaDiSc 2.0 could well avoid the steeped learning curve associated with using R. Another Shiny web application, MetaDTA [26], developed by the United Kingdom National Institute for Health Research (NIHR) Complex Review Support Unit in 2019, is available to conduct DTA meta-analyses. Meta-DiSc 2.0 has an advantage over the MetaDTA software because of its capacity to perform meta-regression analyses and calculate additional measures to quantify heterogeneity.

The app has several limitations. The meta-regression analysis implemented is based on the assumption of equal variances for the random effects of the logit sensitivities and the logit specificities of the compared subgroups. This assumption may be reasonable in many situations, although it may not be in some reviews. It is worth noting that the bivariate I-squared statistic depends on sample size. For this reason, the comparison of I-squared values among meta-analyses with a different number of studies and a different number of diseased and non-diseased participants is limited.

The app does not allow comparing the accuracy of two diagnostic tests, and the current version does not incorporate the risk of bias assessment using the QUADAS-2 tool [31].

The development of this web application was led by the Clinical Biostatistics Unit of the Ramón y Cajal Research Institute (IRYCIS), a unit that has broad experience in diagnostic test synthesis research focused on supporting informed decision-making in the healthcare area. This constitutes a collaborative project for knowledge transfer between IRYCIS and the Complutense University of Madrid and is supported by an intramural project funded by the Ramón y Cajal Research Institute ("*Rapid diagnostic reviews for decision-making in healthcare: analysis of critical points and software development*", 2018). This project has also been funded by Instituto de Salud Carlos III through the project "PI19/00481" (Co-funded by European Regional Development Fund/European Social Fund; "A way to make Europe"/"Investing in your future"). The Biomedical Research Networking Center in Epidemiology and Public Health (CIBERESP) funds the subscription to the shinyapps.io platform where the app is hosted.

Conclusion

We developed an updated version of Meta-DiSc for performing diagnostic test accuracy meta-analyses. All computational algorithms have been validated by comparing different statistical tools and published meta-analyses.

Abbreviations

app: Application; CCHD: Critical congenital heart defects; CI: Confidence interval; CIBERESP: Biomedical research networking center in epidemiology and public health; DOR: Diagnostic odds ratio; DTA: Diagnostic test accuracy; FN: False negatives; FP: False positives; ID: Identifier; IRYCIS: Ramón y Cajal research institute; LCI: Low confidence interval; LR: Likelihood ratios; MOR: Median odds ratios; NIHR: National institute for health research; sROC: Summary receiver operating characteristic; TN: True negatives; TP: True positives; UCI: Upper confidence interval.

Acknowledgements

Not applicable.

Availability and requirements

Project name: Meta-DiSc 2.0
Project home page: <https://metadisc.sourceforge.io>
Operating system(s): Platform independent
Programming language: R
Other requirements: Internet browser
License: GNU General Public License version 3.0 (GPLv3)
Any restrictions to use by non-academics: None

Authors' contributions

JZ, MNP and IA conceived the idea. SF, JS, MNP and JZ developed the software. IA, TP, MF and MR validated the software on a number of reviews and gave suggestions for improvements. All authors participated in preparing this manuscript and have read and approved the final version. The author(s) read and approved the final manuscript.

Funding

Meta-DiSc 2.0 has been developed with funding from an intramural project by the Ramón y Cajal Research Institute ("*Rapid diagnostic reviews for decision-making in healthcare: analysis of critical points and software development*", 2018). It has also been funded by Instituto de Salud Carlos III through the project "PI19/00481" (Co-funded by European Regional Development Fund/European Social Fund; "A way to make Europe"/"Investing in your future"). The Biomedical Research Networking Center in Epidemiology and Public Health (CIBERESP) funds the subscription to the shinyapps.io platform where the app is hosted. The funding body played no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Availability of data and materials

Meta-DiSc 2.0 is a free web application and can be accessed at www.metadisc.es. R code and example data are available at <https://metadisc.sourceforge.io>.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Health Technology Assessment Unit, Hospital Universitario Ramón Y Cajal, IRYCIS, Madrid, Spain. ²CIBER of Epidemiology and Public Health, Madrid, Spain. ³Clinical Biostatistics Unit, Hospital Universitario Ramón Y Cajal, IRYCIS, Madrid, Spain. ⁴Radiology Department, Hospital Universitario Ramón Y Cajal, IRYCIS, UPM, Madrid, Spain. ⁵Internal Medicine Department, Hospital Universitario Ramón Y Cajal, IRYCIS, Madrid, Spain. ⁶Department of Statistics and Data Science, Complutense University of Madrid, Madrid, Spain. ⁷Barts Research Centre for Women's Health, WHO Collaborating Centre, Queen Mary University of London, London, UK. ⁸Iberoamerican Cochrane Center, Biomedical Research Institute Sant Pau (IIB Sant Pau), Barcelona, Spain. ⁹WHO Collaborating Centre for Global Women's Health, Institute of Metabolism and Systems Research, Institute of Applied Health Research, University of Birmingham, Birmingham, UK.

Received: 4 March 2022 Accepted: 10 November 2022
Published online: 28 November 2022

References

- Knottnerus J, Frank B. The evidence base of clinical diagnosis. 2nd ed. London: BMJ Books; 2009.
- Bossuyt PM. Testing COVID-19 tests faces methodological challenges. *J Clin Epidemiol*. 2020;126:172–6.
- Deeks J, Bossuyt P, Gatsonis CE. *Cochrane handbook for systematic reviews of diagnostic test accuracy*. London: The Cochrane Collaboration; 2010.
- Harbord RM, Deeks JJ, Egger M, Whiting P, Sterne JA. A unification of models for meta-analysis of diagnostic accuracy studies. *Biostatistics*. 2007;8(2):239–51.
- Chu H, Cole SR. Bivariate meta-analysis of sensitivity and specificity with sparse data: a generalized linear mixed model approach. *Journal of clinical epidemiology*. 2006;59(12):1331–2 (author reply 2–3).
- Reitsma JB, Glas AS, Rutjes AW, Scholten RJ, Bossuyt PM, Zwinderman AH. Bivariate analysis of sensitivity and specificity produces informative summary measures in diagnostic reviews. *J Clin Epidemiol*. 2005;58(10):982–90.
- Riley RD, Abrams KR, Sutton AJ, Lambert PC, Thompson JR. Bivariate random-effects meta-analysis and the estimation of between-study correlation. *BMC Med Res Methodol*. 2007;7:3.
- Arevalo-Rodriguez I, Steingart KR, Tricco AC, Nussbaumer-Streit B, Kaune-D, Alonso-Coello P, et al. Current methods for development of rapid reviews about diagnostic tests: an international survey. *BMC Med Res Methodol*. 2020;20(1):115.
- Zamora J, Abraira V, Muriel A, Khan K, Coomarasamy A. Meta-DiSc: a software for meta-analysis of test accuracy data. *BMC Med Res Methodol*. 2006;6:31.
- Bates D, Maechler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. *J Stat Softw*. 2015;67:1–48.
- Jackson CH. Multi-State Models for Panel Data: The msm Package for R. *J Stat Softw*. 2011;38(8):1–29.
- Wickham H. *ggplot2: Elegant Graphics for Data Analysis*. New York: 2016. Available from: <https://ggplot2.tidyverse.org>.
- Sievert C. *Interactive Web-Based Data Visualization with R, plotly, and shiny*. CRC: Chapman and Hall; 2020. Available from: <https://plotly-r.com>.
- Balduzzi S, Rucker G, Schwarzer G. How to perform a meta-analysis with R: a practical tutorial. *Evid Based Ment Health*. 2019;22(4):153–60.
- Takwoingi Y, Guo B, Riley RD, Deeks JJ. Performance of methods for meta-analysis of diagnostic test accuracy with few studies or sparse data. *Stat Methods Med Res*. 2017;26(4):1896–911.
- Plana MN, Pérez T, Zamora J. New measures improved the reporting of heterogeneity in diagnostic test accuracy reviews: a metaepidemiological study. *J Clin Epidemiol*. 2021;131:101–12.
- Zhou Y, Dendukuri N. Statistics for quantifying heterogeneity in univariate and bivariate meta-analyses of binary data: the case of meta-analyses of diagnostic accuracy. *Stat Med*. 2014;33(16):2701–17.
- Borchers HW. *pracma: Practical Numerical Math Functions 2021 [2.3.3]*. Available from: <https://CRAN.R-project.org/package=pracma>.
- Macaskill P, Gatsonis C, Deeks JJ, Harbord RM, Takwoingi Y. Chapter 10: analysing and presenting results. In: Deeks JJ, Bossuyt PM, Gatsonis C, editors. *Cochrane handbook for systematic reviews of diagnostic test accuracy version 1.0*. The Cochrane Collaboration; 2010. Available from: <http://srdta.cochrane.org/>.
- Zeileis A, Hothorn T. Diagnostic checking in regression relationships. *R News*. 2002;2:7–10. <http://CRAN.R-project.org/doc/Rnews/>.
- Plana MN, Zamora J, Suresh G, Fernandez-Pineda L, Thangaratinam S, Ewer AK. Pulse oximetry screening for critical congenital heart defects. *Cochrane Database Syst Rev*. 2018;3(3):CD011912.
- Review Manager Web (RevMan Web) [Computer Program]. Version 1.22.0. The Cochrane Collaboration; 2020. Available at revman.cochrane.org.
- Hultcrantz M, Mustafa RA, Leeflang MMG, Lavergne V, Estrada-Orozco K, Ansari MT, et al. Defining ranges for certainty ratings of diagnostic accuracy: a GRADE concept paper. *J Clin Epidemiol*. 2020;117:138–48.
- METADAS: A SAS macro for meta-analysis of diagnostic accuracy studies. User guide version 1.0 beta. December 2008. Available at: <http://srdta.cochrane.org/en/clib.html>. Accessed 3 July 2009.
- Harbord RM, Whiting P. metandi: Meta-analysis of diagnostic accuracy using hierarchical logistic regression. *Stata Journal*. 2009;9(2):211–29.
- Freeman SC, Kerby CR, Patel A, Cooper NJ, Quinn T, Sutton AJ. Development of an interactive web-based tool to conduct and interrogate meta-analysis of diagnostic test accuracy studies: MetaDTA. *BMC Med Res Methodol*. 2019;19(1):81.
- Fahey MT, Irwig L, Macaskill P. Meta-analysis of Pap test accuracy. *Am J Epidemiol*. 1995;141(7):680–9.
- Honest H, Bachmann LM, Gupta JK, Kleijnen J, Khan KS. Accuracy of cervicovaginal fetal fibronectin test in predicting risk of spontaneous preterm birth: systematic review. *BMJ*. 2002;325(7359):301.
- Scheidler J, Hricak H, Yu KK, Subak L, Segal MR. Radiological evaluation of lymph node metastases in patients with cervical cancer. A meta-analysis. *Jama*. 1997;278(13):1096–101.
- Verde PE. Meta-analysis of diagnostic test data: a bivariate Bayesian modeling approach. *Stat Med*. 2010;29(30):3088–102.
- Whiting PF, Rutjes AW, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, Leeflang MM, Sterne JA, Bossuyt PM, QUADAS-2 Group. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med*. 2011;155(8):529–36.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

