



DNA replication-dependent binding of CTCF plays a critical role in adenovirus genome functions

著者	Komatsu Tetsuro, Sekiya Takeshi, Nagata Kyosuke
journal or publication title	Scientific reports
volume	3
page range	2187
year	2013-07
権利	(C) 2013 Nature Publishing Group, a division of Macmillan Publishers Limited.
URL	http://hdl.handle.net/2241/119599

doi: 10.1038/srep02187



OPEN

DNA replication-dependent binding of CTCF plays a critical role in adenovirus genome functions

SUBJECT AREAS:

VIRAL PROTEINS

CHROMOSOMES

ADENOVIRUS

CHROMATIN STRUCTURE

Tetsuro Komatsu^{1,2}, Takeshi Sekiya¹ & Kyosuke Nagata¹

¹Department of Infection Biology, Faculty of Medicine and Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tsukuba 305-8575, Japan, ²Microbiologie Fondamentale et Pathogénicité, MFP CNRS UMR 5234, Université Bordeaux SEGALEN, Bordeaux 33076, France.

Received
13 May 2013Accepted
25 June 2013Published
15 July 2013

The expression of adenovirus late genes is shown to require viral DNA replication, but its mechanism remains elusive. Here we found that knockdown of CTCF suppresses viral DNA replication as well as late, but not early, gene expression. Chromatin immunoprecipitation assays indicated that CTCF binds to viral chromatin depending on viral DNA replication. These findings depict CTCF as a critical regulator for adenovirus genome functions in late phases of infection.

Correspondence and requests for materials should be addressed to K.N. (knagata@md.tsukuba.ac.jp)

In the cell nucleus, the genomic DNA forms chromatin structure. It is being clarified that the higher-order chromatin structure, such as the DNA looping, plays an important role in a dynamic property of the chromatin¹. One of the chromatin organizing proteins, CTCF (CCCTC-binding factor), is a well-characterized chromatin-binding factor involved in the formation of the long-range interactions of chromatin². CTCF has eleven zinc fingers and therefore binds to divergent DNA sequences, as indicated by chromatin immunoprecipitation (ChIP) in combination with tiling arrays (ChIP-on-chip)³ or high-throughput sequencing analyses (ChIP-seq)⁴. A variety of chromatin-related proteins are reported as binding partners of CTCF, including cohesin complexes^{5,6}, a nucleolar protein B23/nucleophosmin, and CTCF itself⁷. These interactions are thought to enable CTCF binding sites to contact each other and/or be tethered to the subnuclear domains, resulting in the formation of intra- and interchromatin interaction². In addition to the role on the cellular chromatin, recent reports have revealed the involvement of CTCF on viral proliferation, as Lieberman and co-workers recently demonstrated the CTCF-mediated formation of chromatin loops on Kaposi's sarcoma-associated Herpesvirus (KSHV) and Epstein-Barr virus (EBV) genomes^{8,9}. It is shown that CTCF regulates the latency-specific chromatin conformation of KSHV and EBV genomes, and siRNA-mediated depletion of CTCF or mutations in the CTCF binding sites disrupt the chromatin architecture and de-regulate latent gene expression^{8,9}. Thus, CTCF could impact on the regulation of not only cellular but also viral chromatin.

The adenovirus (Ad) has a linear double-stranded DNA genome that forms chromatin-like structure in the virion¹⁰. Previously, we have reported that viral chromatin structure regulates the expression of viral early genes (e.g. E1A, E4 genes) in early phases of infection^{11,12}. The expression of the late genes (e.g. major late genes) are hardly observed during early phases of infection, while concomitantly with the onset of viral DNA replication, those genes are fully activated. Thomas and Mathews demonstrated that the expression of the late genes requires viral DNA replication in *cis*¹³. In addition, we have shown the regulatory mechanism of the viral chromatin structure during DNA replication and proposed a possible role of viral DNA replication in the activation of late genes¹⁴. Thus, it is suggested that the regulation of viral chromatin structure has a significant role in the DNA replication-dependent activation of viral genes. In spite of these evidences, however, the functional relationship between viral gene expression and DNA replication in infected cells remains largely unclear. In this study we sought to further clarify the role of chromatin structure and/or chromatin-related factors on the Ad genome DNA. As described above, it is shown that CTCF plays a role on the chromatin of some DNA viruses^{8,9}. These lead us to hypothesize that CTCF could function also on Ad chromatin.

Results

CTCF is required for viral DNA replication and late gene expression. To study a role of CTCF, we carried out knock down (KD) of the expression of CTCF by siRNA treatment (Fig. 1A). Either control siRNA (siCont) or




Figure 1 | Effects of CTCF KD on viral DNA replication and gene expression. (A) Western blotting and knockdown of CTCF. Cell lysates were prepared from HeLa cells treated with siCont (lanes 3) or siCTCF (lane 4) and subjected to western blot analyses using anti-CTCF (top panel) and anti- β -actin antibodies (bottom panel). For siCont-treated cells, 25% (lane 1) and 50% (lane 2) volume of lysates were also loaded. Full-size images are shown in Supplementary figure 1. (B) Amounts of viral DNA. HeLa cells treated with siCont or siCTCF were infected with HAdV5 at an MOI of 100, and total DNAs were purified at 6 and 12 hpi. The amount of viral DNA was measured by qPCR using primers for the E1A promoter. The amount at 12 hpi relative to that at 6 hpi was graphed. Mean values with s.d. were obtained from three independent experiments. (C) RT-qPCR assays. Total RNAs were purified at 6 and 12 hpi, and subjected to RT-qPCR using indicated primer sets. The mRNA levels relative to those of control cells at 12 hpi were graphed. Mean values with s.d. were obtained from three independent experiments. *P*-values are calculated using Student's *t*-test. "n.s." indicates "not statistically significant".

siRNA targeted for CTCF (siCTCF) was introduced into HeLa cells, and then cell lysates were prepared and subjected to western blot analyses using anti-CTCF antibody. Only a single band corresponding to CTCF was detected, demonstrating the specificity of the antibody. Under our experimental condition, the expression level of CTCF in siCTCF-treated cells was decreased to approximately 25% of that in control cells (Fig. 1A, compare lane 4 with lanes 1–3).

To test whether CTCF plays a role in Ad DNA replication and gene expression, we carried out CTCF KD followed by quantitative PCR (qPCR) of viral DNA and RT-qPCR (Fig. 1B and C). Under our experimental condition, the onset of viral DNA replication can be observed around 8 hpi (hours post infection)^{12,14}. siCont- or siCTCF-treated cells were mock-infected or infected with human adenovirus type 5 (HAdV5) at an MOI (multiplicity of infection) of 100, and at 6 (for early phases) and 12 hpi (for late phases of infection) total DNAs and RNAs were purified. We first measured viral DNA amounts by qPCR using a primer set targeted for the E1A promoter region (E1A pro, see Table 2) to evaluate the efficiency of viral DNA replication (Fig. 1B). In siCont-treated cells, the amount of viral DNA was increased by ~30 fold through viral DNA replication. In contrast, siCTCF-treated cells allowed only ~9 fold amplification of viral DNA.

Next, we performed RT-qPCR assays using several primer sets for cellular and viral genes (Fig. 1C). Under the condition employed here, the mRNA level of *GAPDH* was unaffected by Ad infection and siRNA treatment, and that of *CTCF* was specifically decreased by siCTCF treatment (Fig. 1C, *GAPDH* and *CTCF*). The mRNA levels of viral early genes were not significantly affected by CTCF KD (Fig. 1C, E1A and E4), suggesting that CTCF is not involved in viral early gene expression. In contrast, the level of mRNA transcribed from the major late promoter (MLP) was drastically decreased by siCTCF treatment (Fig. 1C, MLP). Similarly, the mRNA level of E2A was reduced by CTCF KD, particularly at 12 hpi (Fig. 1C, E2A). It is noted that E2 gene transcription is regulated by early and late promoters and transcription from the E2 late promoter depends on viral DNA replication¹⁵ (Unpublished data). Therefore, it is reasonable to

assume that CTCF KD could predominantly affect the transcription from the E2 late promoter, although in this study we did not discriminate E2A mRNAs transcribed from two promoters. Collectively, these results suggest that CTCF is critical for viral DNA replication as well as late, but not early, gene expression.

CTCF binds to viral chromatin in a DNA replication-dependent manner. To examine whether CTCF functions directly on viral chromatin, ChIP assays were performed using anti-CTCF antibody (Fig. 2). Since the effect of CTCF KD was observed in late phases of infection (Fig. 1), first we studied using infected cells at 12 hpi for ChIP assays (Fig. 2A). We used a variety of primer sets for the Ad genome to test the genome-wide binding of CTCF (see Fig. 2B). We found that CTCF is recruited into several regions of the virus genome, including the MLP region and the ORF regions of the viral structural protein (Hexon) and the E4 ORF3 gene (E4 orf). In addition, a weak binding of CTCF at the E1A pro region was observed. Next, we focused on three CTCF binding sites, the E1A pro, MLP, and Hexon regions, and performed ChIP assays using cells cultured in the absence or presence of hydroxyurea (HU), a DNA replication inhibitor, to examine whether the CTCF binding observed here is DNA replication-dependent (Fig. 2C). The recruitment of CTCF into those regions was observed only at 12 hpi, and this was inhibited by the addition of HU, indicating that CTCF is recruited onto viral chromatin in a DNA replication-dependent manner.

Discussion

The results obtained in this study indicate that CTCF binds to Ad chromatin depending on its DNA replication and plays a pivotal role in late phases of infection. Our KD experiments clearly reveal that CTCF is required for viral DNA replication and late gene expression (Fig. 1B and C). It is demonstrated that viral late genes are activated depending on its DNA replication¹³. Thus, the mRNA levels of late genes also should be affected when only DNA replication would be directly repressed by CTCF KD. Conversely, viral factors involved in




Figure 2 | CTCF binding on viral chromatin. (A) ChIP assays with anti-CTCF antibody. HeLa cells were infected with HAdV5 at an MOI of 100, and at 12 hpi subjected to ChIP assays using anti-CTCF and anti-FLAG antibodies. The binding levels were calculated as fold enrichment against that obtained in a negative control (anti-FLAG antibody). Mean values with s.d. were obtained from three independent experiments. (B) Structure of the Ad genome. Arrows represent promoters of viral genes. Target regions for ChIP assays are indicated by gray bars. (C) Effect of viral DNA replication on CTCF binding. HeLa cells were infected with HAdV5 and cultured in the absence or presence of 2 mM HU. At 6 and 12 hpi, ChIP assays were carried out as described above. *P*-values are calculated using Student's *t*-test.

its DNA replication are encoded by the E2 gene, and its expression was suppressed by CTCF KD (Fig. 1C). Therefore, the inhibition of viral late gene expression also could be the cause for less efficient viral DNA replication. Because of this interdependency, we could not precisely discriminate whether CTCF KD primarily affects viral DNA replication or late gene expression (or both). In addition, we could not exclude the possibility that the lower level of the MLP mRNA in siCTCF-treated cells results from less amount of viral DNA templates due to the defect in viral DNA replication. Nonetheless, we speculate that CTCF may be involved in the regulation of late gene expression: First, the binding of CTCF is dependent on viral DNA replication (Fig. 2C), suggesting that this protein possibly functions in the step(s) later than viral DNA replication; second, when the level of the MLP mRNA at 12 hpi was normalized by the amounts of the virus genome, the level of the MLP mRNA per one viral DNA in CTCF KD cells was still lower than that in control cells (Fig. 1, MLP mRNA level: ~15%/virus genome: ~30% = MLP mRNA per one viral DNA: ~50%).

It is an important question how CTCF regulates the function of the virus genome/chromatin. Although in this study we found several CTCF binding sites (Fig. 2A), there could be additional regions for CTCF on the virus genome. Thus, at the moment, it is difficult to dissect a role of each CTCF binding site and the cooperative function among those regions. It is suggested that the E1B, IX, and E2 late genes/promoters are also activated depending on viral DNA replication^{15–17} as is the MLP. However, we could not observe the CTCF binding on those regions (Fig. 2A), suggesting that the recruitment of CTCF onto each late promoter seems not to be required for the genome-wide coordination between viral DNA replication and the expression of late genes. Further studies are needed to address these points. As described above, it is shown that Ad DNA replication is required for the activation of viral late genes¹³. Similarly it is reported that DNA replication is essential for the expression of certain cellular genes such as the *HoxB* gene¹⁸, although the molecular details remain to be determined. To our knowledge, this is the first report indicating the possible involvement of CTCF in the DNA replication-dependent activation of the genes. Thus, our findings may provide insight into an uncharacterized mechanism of gene regulation that involves DNA replication.

Methods

Cells and viruses. Maintenance of HeLa cells, and purification and infection of human adenovirus type 5 (HAdV5) were carried out essentially as described previously^{12,14}. Hydroxyurea (HU) was added at the final concentration of 2 mM right after infection to block DNA replication.

Antibodies. To obtain recombinant CTCF N-terminal region (amino acids (aa) 1–267) as an antigen, the expression vector for His-tagged CTCF(1–267) was constructed. cDNA fragment of full-length CTCF was amplified by PCR with a primer set, 5'-AGGGCATATGGAAGGTGATGCAGTCGAAGCCATTGTGG-3' and 5'-AGCCTCGAGAAGTCTGGCGACGCACAAGGCTCCGCC-3', and cloned into the pBluescript-FLAG vector (pBS-FLAG-CTCF). Using pBS-FLAG-CTCF as a template, cDNA fragment corresponding to aa 1–267 was amplified by PCR with a primer set, 5'-AGGGCATATGGAAGGTGATGCAGTCGAAG-CCATTGTGG-3' and 5'-GTTGAATTCAGTGAATGTCTTCTTTACAC-3', and cloned into the pET-14b vector. *E. coli* was transformed with the resultant vector, pET-14b-CTCF(1–267), and His-CTCF(1–267) was expressed and purified using the Ni-NTA resin (Novagen) according to the manufacturer's protocol.

Rabbit anti-CTCF antibody was raised against His-CTCF(1–267) according to standard protocols. Mouse anti-FLAG M2 and mouse anti-β-actin antibodies were described elsewhere^{12,14}.

RT-qPCR assays. RT-PCR and quantitative PCR (qPCR) were performed essentially as described previously^{12,14}. Total RNAs were purified by phenol extraction followed by DNase I treatment. cDNA was synthesized from total RNA (1 μg) using ReverTraAce (Toyobo) and oligo-dT primer according to the manufacturer's protocol. qPCR was carried out using FastStart SYBR Green Master (Roche) and

Table 1 | Primers used for RT-qPCR

Primer	Sequence (5'-3')
GAPDH forward	AGCCAAAAGGGTCATCATCTC
GAPDH reverse	GGACTGTGGTCATGAGTCCTTC
CTCF forward	TGACACAGTCATAGCCCGAAAA
CTCF reverse	TGCCTTGCTCAATATAGGAATGC
E1A forward	GAGACATATTATCTGCCACGGAG
E1A reverse	AGTGAGTAAGTCAATCCCTTCCTG
E4 forward	ACAGAACCCTAGTATTCAACCTGC
E4 reverse	GACAGCGACATGAACCTTAAGTGAG
MLP forward	ACTCTCTCCGCATCGCTGT
MLP reverse	GTGACTGGTTAGACGCCTTCT
E2A forward	GTGTAGACACCTAAGCTCGCCTT
E2A reverse	CTCAAACACTGCCTGACCAAGT



Table 2 | Primers used for ChIP

Primer	Sequence (5'-3')
5' inverted terminal region forward	CAATATGATAATGAGGGGGTGG
5' inverted terminal region reverse	ACTACAACATCCGCCTAAAACC
E1A enhancer region forward	CGGTGTACACAGGAAGTGACAAT
E1A enhancer region reverse	AGTCTCCACGTAACGGTCAAAGT
E1A promoter region forward	GGGTCAAAGTTGGCGTTTTA
E1A promoter region reverse	CAAAATGGCTAGGAGGTGGA
E1A coding region forward	GAGACATATTATCTGCCACGGAG
E1A coding region reverse	AGTGAGTAAGTCAATCCCTTCCTG
E1A 3' region forward	CCTTCTAACACCTCCTGAGATAC
E1A 3' region reverse	ACACACGCAATCACAGGTTTAC
E1B promoter region forward	GTGTGTGGTTAACGCCTTTGT
E1B promoter region reverse	GAGGTACTGTTAGAGCTCTGTTCCA
IX promoter region forward	GGCTCTAGCGATGAAGATACAGAT
IX promoter region reverse	CATCACATTCTGACGCACCC
ML promoter region forward	AGGTGATTGGTTTGTAGGTGTAGG
ML promoter region reverse	CTCCTCGTTTTGGAAACTGAC
VA I gene region forward	GTGCAAAAAGGAGAGCCTGTAAG
VA I gene region reverse	AGGAAGCCAAAAGGAGCACT
Hexon ORF region forward	CGCAGTGGTCTTACATGCAC
Hexon ORF region reverse	CACACGGTTATCACCCACAG
E2B ORF region forward	AGAAGAACATGCCGCAAGAC
E2B ORF region reverse	TCCAAGGCGAGCTTAAGTGT
E2 late promoter region forward	ATTATCGGTACCTTTGAGCTGC
E2 late promoter region reverse	AGAATGTGGCCTGGGTAAT
E3 promoter region forward	AAGTTCAGATGACTAACTCAGGGG
E3 promoter region reverse	AGAGTTAGGATTGCCCTGACGAG
E4 ORF3 region forward	TGGCGTGGTCAAACCTTACA
E4 ORF3 region reverse	GATTTTACAATGGCCGGACT
E4 promoter region forward	CCATAACAGTCAGCCTTACCAGT
E4 promoter region reverse	GTGACGATTGAGGAAGTTGTG

Thermal Cycler Dice Real Time System (Takara) according to the manufacturers' protocol.

Table 1 indicates primer sequences for *GAPDH*, *CTCF*, E1A, E4, MLP, and E2A genes.

ChIP assays. ChIP assays were carried out according to the protocol from Chromatin Immunoprecipitation Assay Kit (Millipore) with minor modification, essentially as described previously^{12,14}. Briefly, cells were fixed with 1% formaldehyde, followed by the addition of glycine at the final concentration of 125 mM for quenching. After centrifugation, cell pellets were lysed with SDS lysis buffer (50 mM Tris-HCl [pH 7.9], 10 mM EDTA, and 1% SDS), and lysates were subjected to sonication to shear the chromatin DNA to ~1 kbp in size. Sonicated samples were diluted 10 fold with ChIP dilution buffer (16.7 mM Tris-HCl [pH 7.9], 1.2 mM EDTA, 167 mM NaCl, 1.1% Triton X-100, and 0.01% SDS) and then pre-cleared with Protein A Sepharose 4 Fast Flow (GE Healthcare). An antibody was added to the pre-cleared sample solution and incubated overnight at 4°C. Antibody-protein-DNA complexes were incubated with Protein A Sepharose at 4°C for 1 hr, and then the beads were washed with Low Salt Wash Buffer (20 mM Tris-HCl [pH 7.9], 2 mM EDTA, 150 mM NaCl, 1% Triton X-100, and 0.1% SDS), High Salt Wash Buffer (20 mM Tris-HCl [pH 7.9], 2 mM EDTA, 500 mM NaCl, 1% Triton X-100, and 0.1% SDS), LiCl Wash Buffer (10 mM Tris-HCl [pH 7.9], 1 mM EDTA, 0.25 M LiCl, 1% NP-40, and 1% deoxycholic acid), and TE (10 mM Tris-HCl [pH 7.9] and 1 mM EDTA) successively. Protein-DNA complexes were eluted from the beads with elution buffer (1% SDS, 100 mM NaHCO₃, and 10 mM DTT), and crosslinking was reversed by incubation at 65°C for 4 hr. After treatment with Proteinase K, DNAs were recovered by phenol/chloroform extraction and ethanol precipitation. Obtained DNA samples were subjected to qPCR as described above.

Table 2 shows primer sequences for 5' inverted terminal region (5' ITR), the E1A enhancer (E1A enh), the E1A promoter (E1A pro), the E1A coding region (E1A cds), the E1A 3' region (E1A 3'), the E1B promoter (E1B pro), the IX promoter (IX pro), the ML promoter (MLP), the VA I gene (VA I), the Hexon ORF (Hexon), the E2B ORF (E2B orf), the E2 late promoter (E2late pro), the E3 promoter (E3 pro), the E4 ORF3 (E4 orf), and the E4 promoter regions (E4 pro).

siRNA-mediated knockdown, quantification of viral DNA, and western blot assays. These experiments were carried out essentially as described previously^{12,14,19}. siRNA targeted for CTCF was commercially purchased (Stealth siRNA; Invitrogen). siRNAs were introduced into cells with Lipofectamine 2000 (Invitrogen) according to the manufacturer's protocol. Total DNAs were purified by treatment with Proteinase K, followed by phenol/chloroform extraction and ethanol precipitation, as described previously¹⁹. Quantitative determination of viral DNA was carried out by qPCR as described above. For western blot analyses, cell lysates were subjected to SDS-PAGE,

and proteins were transferred to a PVDF membrane (Millipore). The membrane was incubated with primary antibodies, followed by incubation with secondary antibodies conjugated with horseradish peroxidase (GE Healthcare). The blot was developed using Chemi-Lumi One (Nacalai tesque).

1. Raab, J. R. & Kamakaka, R. T. Insulators and promoters: closer than we think. *Nat Rev Genet* **11**, 439–46 (2010).
2. Phillips, J. E. & Corces, V. G. CTCF: master weaver of the genome. *Cell* **137**, 1194–211 (2009).
3. Kim, T. H. *et al.* Analysis of the vertebrate insulator protein CTCF-binding sites in the human genome. *Cell* **128**, 1231–45 (2007).
4. Barski, A. *et al.* High-resolution profiling of histone methylations in the human genome. *Cell* **129**, 823–37 (2007).
5. Stedman, W. *et al.* Cohesins localize with CTCF at the KSHV latency control region and at cellular *c-myc* and H19/Igf2 insulators. *EMBO J* **27**, 654–66 (2008).
6. Wendt, K. S. *et al.* Cohesin mediates transcriptional insulation by CCCTC-binding factor. *Nature* **451**, 796–801 (2008).
7. Yusufzai, T. M., Tagami, H., Nakatani, Y. & Felsenfeld, G. CTCF tethers an insulator to subnuclear sites, suggesting shared insulator mechanisms across species. *Mol Cell* **13**, 291–8 (2004).
8. Kang, H., Wiedmer, A., Yuan, Y., Robertson, E. & Lieberman, P. M. Coordination of KSHV latent and lytic gene control by CTCF-cohesin mediated chromosome conformation. *PLoS Pathog* **7**, e1002140 (2011).
9. Tempera, I., Klichinsky, M. & Lieberman, P. M. EBV latency types adopt alternative chromatin conformations. *PLoS Pathog* **7**, e1002180 (2011).
10. Giberson, A. N., Davidson, A. R. & Parks, R. J. Chromatin structure of adenovirus DNA throughout infection. *Nucleic Acids Res* **40**, 2369–76 (2012).
11. Haruki, H., Okuwaki, M., Miyagishi, M., Taira, K. & Nagata, K. Involvement of template-activating factor I/SET in transcription of adenovirus early genes as a positive-acting factor. *J Virol* **80**, 794–801 (2006).
12. Komatsu, T., Haruki, H. & Nagata, K. Cellular and viral chromatin proteins are positive factors in the regulation of adenovirus gene expression. *Nucleic Acids Res* **39**, 889–901 (2011).
13. Thomas, G. P. & Mathews, M. B. DNA replication and the early to late transition in adenovirus infection. *Cell* **22**, 523–33 (1980).
14. Komatsu, T. & Nagata, K. Replication-uncoupled histone deposition during adenovirus DNA replication. *J Virol* **86**, 6701–11 (2012).
15. Swaminathan, S. & Thimmapaya, B. Regulation of adenovirus E2 transcription unit. *Curr Top Microbiol Immunol* **199**, 177–94 (1995).
16. Matsui, T., Murayama, M. & Mita, T. Adenovirus 2 peptide IX gene is expressed only on replicated DNA molecules. *Mol Cell Biol* **6**, 4149–54 (1986).



17. Maxfield, L. F. & Spector, D. J. Readthrough activation of early adenovirus E1b gene transcription. *J Virol* **71**, 8321–9 (1997).
18. Fisher, D. & Méchali, M. Vertebrate HoxB gene expression requires DNA replication. *EMBO J* **22**, 3737–48 (2003).
19. Samad, M. A., Komatsu, T., Okuwaki, M. & Nagata, K. B23/nucleophosmin is involved in regulation of adenovirus chromatin structure at late infection stages, but not in virus replication and transcription. *J Gen Virol* **93**, 1328–38 (2012).

Acknowledgments

This work was supported in part by grants-in-aid for scientific research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (to K.N.).

Author contributions

T.K. and K.N. designed the study; T.K. and T.S. performed the experiments; T.K. and K.N. prepared the manuscript; All authors reviewed the manuscript.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Komatsu, T., Sekiya, T. & Nagata, K. DNA replication-dependent binding of CTCF plays a critical role in adenovirus genome functions. *Sci. Rep.* **3**, 2187; DOI:10.1038/srep02187 (2013).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported license. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0>