



Association between Persistent Staphylococcus aureus Nasal Carriers with High Frequency of Skin Abscesses Among American Football Players

著者	SUZUKI Kotaro, SHIBA Hideki, ARAI Nobunari, GOCHI Kentaro, MATSUMOTO Tsuyoshi, TAGAMI Kazumi
journal or publication title	筑波大学体育系紀要
volume	37
page range	173-178
year	2014-03
URL	http://hdl.handle.net/2241/121331

Association between Persistent *Staphylococcus aureus* Nasal Carriers with High Frequency of Skin Abscesses Among American Football Players

SUZUKI Kotaro^{*}, SHIBA Hideki^{*}, ARAI Nobunari^{**},
GOCHI Kentaro^{*}, MATSUMOTO Tsuyoshi^{*} and TAGAMI Kazumi^{*}

Key words: Physical contact sports, American football, *Staphylococcus aureus*, skin lesion, abscess, biofilm formation

Appendix: This project report submitted for publication in a scientific journal. Primary results were reported in this study.

1. Introduction

Staphylococcus aureus (*SA*) is a common bacterium that often resides harmlessly on the skin or in the nasal cavities of healthy individuals. It is both a human commensal and a frequent source of infections that can be mild or life-threatening. The primary reservoir of *SA* is thought to be the anterior nares (nostrils). About 25% to 50% of individuals carry nasal *SA* in at any given time¹⁾, but this does not necessarily manifest as clinical infection. However, *SA* can cause deep skin infections²⁾.

Risk factors for *SA* that contribute to a high incidence of infection among athletes include close physical contact^{3,4)}, skin damage⁵⁾ and sharing contaminated equipment⁶⁾. Direct physical contact of athletes with exposed extremities during practice or competition can lead to minor abrasions and create portals of entry for *SA*. Most *SA* infections in athletes are caused by minor skin cuts and *SA* is one of the major sources of wound infections in American football players^{7,8)}. Contact sports provide players with more opportunities for skin infections due to skin trauma from turf burns, shaving and contact with secretions and body fluids⁷⁾. Additionally, *SA* typically infects cutaneous skin and spreads to form clusters, although common-source exposure can contribute to transmission⁸⁾.

Nasal carriage of *SA* is an important risk factor for infection in both community and hospital settings⁹⁾. Longitudinal studies have shown that healthy adults can be differentiated into the following types of *SA* carriers:

10% - 20% are persistent carriers (PC), 30% - 50% are intermittently colonized and are thus intermittent carriers (IC), and the remainder comprises non-carriers (NC), who apparently never carry nasal *SA*. The risk of *SA* infection among IC and occasional carriers (OC) is similar to that of NC, suggesting that only PC and OC actually exist¹⁰⁾. Carriage rates are higher among medical patients⁹⁾. Skin wounds have recently been identified as triggers of persistent nasal carriage of *SA* and PC are at higher risk of developing *SA* infections. The mechanisms involved in nasal *SA* colonization and growth at skin lesions are largely unknown.

Skin abscesses might also play a role in nasal *SA* carriage due to a difference in carriage within populations and nasal carriage as a source of *SA*. Skin abscess formation is a pathogen-driven process that usurps the default response of the host and enhances microbial replication and dissemination¹¹⁾. Also, the frequency of nasal colonization with *SA* in the general population has increased in parallel with an increase in that of *SA* skin infection¹²⁾. Thus, skin abscesses are potential sources of nasal *SA* carriage.

Biofilms are bacterial communities that are resistant to host immune responses and antimicrobial agents¹³⁾ and whether they play roles in nasal colonization remains a matter of debate¹⁴⁾. Colonization of the nasal epithelium¹⁵⁾ and the mucosa of patients with chronic rhinosinusitis¹⁶⁾ with *SA* strains has recently been associated with their ability to form biofilms *in vitro*. Thus, we investigated

* Faculty of Health and Sport Sciences, University of Tsukuba

** Faculty of Education, Hakuoh University

whether nasal biofilm formation is involved in persistent nasal *SA* colonization in a population of athletes involved in a contact sport.

We found an association between skin abscesses and persistent nasal *SA* carriage, including nasal *SA* density and biofilm formation. Furthermore, our results suggested that persistent *SA* carriers should be profiled among populations of athletes.

2. Materials and methods

Subjects

Thirty-six male (age: 20.5 ± 1.2 years; height: 172.3 ± 5.4 cm; body mass: 81.3 ± 10.8 kg, BMI: 27.3) collegiate American football players participated in the present study. The same person collected nasal samples from the players between January and November 2012. All of the players completed a questionnaire regarding basic demographic information and clinical data about abscesses and skin lesions. The research protocol was approved by the institutional ethical review board, and volunteers signed informed consent forms before entering the study.

Bacterial isolates

Samples collected from the right or left anterior nares using sterile swabs were mixed with 100 μ L of phosphate-buffered saline (PBS) and then 10 μ L portions were spread on compact Dry X-SA media (CD-XSA; Nissui Pharmaceutical Co., Ltd., Tokyo, Japan) and incubated at 37°C for 24 ± 2 h. Specific blue *SA* colonies¹⁷⁾ were counted and *SA* was extracted from the media to determine biofilm formation.

Definitions of carriers

The types of *SA* nasal carriage comprise persistent (PC), occasional (OC), intermittent (IC) and non- (NC) carriage. However, how to precisely identify these states has not reached consensus, but most studies have been based on cultures from 10 - 12 weekly nasal swabs¹⁸⁾. We defined the carrier index as the number of positive/number of total swabs for each player. We excluded those with < 4 swabs during the study period. We defined PC, IC, OC and NC as *SA* positivity in five of seven swabs, four swabs, three consecutive swabs or sporadic, and three or less and none, respectively.

Biofilm formation assay

Staphylococcus aureus cells cultured in Luria-Bertani

(LB) overnight were 100 fold diluted in Tryptic Soy Broth (Sigma-Aldrich, Tokyo, Japan), mixed with 3% NaCl and 0.5% glucose and then incubated at 96-well plates at 37°C for 24 h. The cells were washed twice with distilled water and stained with 200 μ L of 0.3% crystal violet (CV) as previously described¹⁹⁾, rinsed twice with 300 μ L of double-distilled water (DDW) and thoroughly dried. The plates were shaken for 1 h incubation at room temperature with 200 μ L of 95% ethanol per well. After incubation, the optical density (OD) at 595 nm was measured using a plate reader (BIO RAD, Hercules, CA, USA).

Statistical data analysis

The statistical significance of *SA* colony counts and biofilm formation among carriers was determined using Student's t-test. Statistical significance was set at $P < 0.05$.

3. Results

We obtained seven swabs from each player for a total of 217 swabs. According to our definitions, 5 (13.8%), 4 (11.1%), 17 (47.2%) and 10 (27.8 %) were PC, IC, OC and NC (Fig. 1). None of the isolates obtained in the present study carried the *mecA* gene encoding methicillin resistance (data not shown).

We compared the frequency of skin abscesses in each *SA* nasal carrier (Table 1) based on responses to three questionnaires. An average of 3.60 of 5 PC had skin abscesses, most of which were located in the knees and elbows (Fig. 2), and in the armpit of one. During the study, three carriers each caused severe abscesses in 32, 34 and 36 players. Conversely, four OC and 18 IC had an average of 1.25 and 0.76 skin abscesses, respectively (Table 1), most of which were located at the knees and elbows (Fig. 2). Ten NC had an average of 1.20 skin abscesses, and these were sporadically located at the knees, shanks, fingers and ankles. Skin abscesses in PC who had been nasally infected with *SA* over the long term mostly presented in the knees and elbows.

Mean colony counts were significantly higher among PC than IC, and OC (109.9 ± 84.5 vs. 72.6 ± 82.7 CFU/mL, $P = 0.038$) (Fig. 3A). The density of *SA* colonies recovered from nasal swabs of samples from PC varied from one to a few hundred (Fig. 3B) and curves of kinetics differed among PC groups.

We compared relationships between amounts of nasal *SA* colonies and biofilm formation in six PC by




Fig. 1 Longitudinal distribution of *Staphylococcus aureus* nasal carrier indices among 36 collegiate American football players. Carrier indicator represents number of nasal swabs that were positive for *Staphylococcus aureus* among total number of swabs per player. Black and white circles, positive and negative samples respectively. IC, intermittent (n = 4); NC, Non (n = 10); OC, occasional (n = 17) and PC, persistent (n =5) carriers.

Table 1 Abscesses in carriers among American football players (n = 36).

	^a PC	^b IC	^c OC	^d NC
Players (n)	5	4	17	10
Total abscess	18	5	13	12
^e Average skin abscesses	3.60	1.25	0.76	1.20

^aPC, persistent carrier, ^bIC, intermittent carrier; ^cOC, occasional carrier; ^dNC, non-carrier. ^eCalculated




Fig. 2 Anatomical location of skin abscess in front (A) and on back (B) of players. Closed circle (●) indicates a site of skin lesions.

measuring adherence to a plastic membrane. The samples were categorized as high- (≥ 200), middle- (100–199) and low- (>99) CFU. This cutoff was selected based on the data shown in Fig. 3B. Fig. 4 shows the representative optical density (OD) of adherence. The OD of the biofilm formation was 2-fold higher in those with high-, than with middle- or low-CFU (0.497 ± 0.175 vs. 0.293 ± 0.261 and 0.268 ± 0.25 ; $P = 0.045$ and $P = 0.025$, respectively).





Fig. 3 Comparison of yield of *Staphylococcus aureus* colonies (A) and their kinetics (B) from nasal swabs of infected players. *Staphylococcus aureus* colony counts were visually determined. *Significantly different ($p < 0.05$) from intermittent and occasional carriers.




Fig. 4 Nasal colony counts are associated with *Staphylococcus aureus* biofilm formation in persistent carriers. Various concentrations of colony-forming units (≥ 200 , 100 - 199, < 99) were generated from *Staphylococcus aureus* and biofilm formation was quantified using a microplate reader. Values are averages of quadruplicate readings from at least two separate experiments \pm standard deviations. * $P < 0.05$.

4. Conclusion

- 1) We detected persistent *SA* nasal carriers (Fig. 1) in whom rates of rate of skin abscesses (Fig. 2) were higher than that in other carriers (Table 1).
- 2) The density of *SA* colonies that can be generated by nasal swabs obtained from carriers varies from single digits to a few hundred (Fig. 3B), and a high load of *SA* nasal colonization is closely associated with persistent carriage (Fig. 3A).
- 3) Biofilm formation depended on nasal *SA* density (Fig. 4).

5. Acknowledgments

We thank all participants involved in the study and the laboratory staff of Exercise and Environmental Health to help with this project. This study was supported by a Faculty of Health and Sport Sciences, University of Tsukuba Research Project.

References

- 1) Kluytmans J, van Belkum A, Verbrugh H (1997): Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks. *Clin Microbiol Rev* 10: 505-20.
- 2) Lina G, Piemont Y, Godail-Gamot F, Bes M, Peter MO, Gauduchon V, Vandenesch F, Etienne J (1999): Involvement of Pantone-Valentine leukocidin-producing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clin Infect Dis* 29: 1128-32.
- 3) Stacey AR, Endersby KE, Chan PC, Marples RR (1998): An outbreak of Methicillin resistant *Staphylococcus aureus* infection in a rugby football team. *Br J Sports Med* 32: 153-4.
- 4) Lindenmayer JM, Schoenfeld S, O'Grady R, Carney JK (1998): Methicillin-resistant *Staphylococcus aureus* in a high school wrestling team and the surrounding community. *Arch Intern Med* 158: 895-9.
- 5) Elston DM (2007): Community-acquired methicillin-resistant *Staphylococcus aureus*. *J Am Acad Dermatol* 56: 1-16; quiz 7-20.
- 6) Oller AR, Province L, Curless B (2009): *Staphylococcus aureus* recovery from environmental and human locations in 2 collegiate athletic teams. *J Athl Train* 45: 222-9.
- 7) Begier EM, Frenette K, Barrett NL, Mshar P, Petit S, Boxrud DJ, Watkins-Colwell K, Wheeler S, Cebelinski EA, Glennen A, Nguyen D, Hadler JL (2004): A high-morbidity outbreak of Methicillin-resistant *Staphylococcus aureus* among players on a college football team, facilitated by cosmetic body shaving and turf burns. *Clin Infect Dis* 39: 1446-53.
- 8) Kazakova SV, Hageman JC, Matava M, Srinivasan A, Phelan L, Garfinkel B, Boo T, McAllister S, Anderson J, Jensen B, Dodson D, Lonsway D, McDougal LK, Arduino M, Fraser VJ, Killgore G, Tenover FC, Cody S, Jernigan DB (2005): A clone of Methicillin-resistant *Staphylococcus aureus* among professional football players. *N Engl J Med* 352: 468-75.
- 9) Wertheim HF, Melles DC, Vos MC, van Leeuwen W, van Belkum A, Verbrugh HA, Nouwen JL (2005): The role of nasal carriage in *Staphylococcus aureus* infections. *Lancet Infect Dis* 5: 751-62.
- 10) van Belkum A, Verkaik NJ, de Vogel CP, Boelens HA, Verveer J, Nouwen JL, Verbrugh HA, Wertheim HF (2009): Reclassification of *Staphylococcus aureus* nasal carriage types. *J Infect Dis* 199: 1820-6.
- 11) Cheng AG, DeDent AC, Schneewind O, Missiakas D (2011): A play in four acts: *Staphylococcus aureus* abscess formation. *Trends Microbiol* 19: 225-32.
- 12) Gorwitz RJ, Kruszon-Moran D, McAllister SK,

- McQuillan G, McDougal LK, Fosheim GE, Jensen BJ, Killgore G, Tenover FC, Kuehnert MJ (2008): Changes in the prevalence of nasal colonization with *Staphylococcus aureus* in the United States, 2001–2004. *J Infect Dis* 197: 1226–34.
- 13) Rohde H, Burdelski C, Bartscht K, Hussain M, Buck F, Horstkotte MA, Knobloch JK, Heilmann C, Herrmann M, Mack D (2005): Induction of *Staphylococcus epidermidis* biofilm formation via proteolytic processing of the accumulation-associated protein by staphylococcal and host proteases. *Mol Microbiol* 55: 1883–95.
- 14) Krismer B, Peschel A (2011): Does *Staphylococcus aureus* nasal colonization involve biofilm formation? *Future Microbiol* 6: 489–93.
- 15) Quinn GA, Tarwater PM, Cole AM (2009): Subversion of interleukin-1-mediated host defence by a nasal carrier strain of *Staphylococcus aureus*. *Immunology* 128: e222–9.
- 16) Reid DW, Kirov SM, Webb JS, Carroll V, Kjelleberg S, Champion A, Sanderson K (2006): Biofilm dispersal and exacerbations of cystic fibrosis lung disease. *Pediatr Pulmonol* 41: 1254; author reply 5.
- 17) Teramura H, Mizuochi S, Kodaka H (2010): Evaluation of the compact dry X-SA method for enumerating *Staphylococcus aureus* in artificially contaminated food samples. *Biocontrol Sci* 15: 149–54.
- 18) VandenBergh MF, Yzerman EP, van Belkum A, Boelens HA, Sijmons M, Verbrugh HA (1999): Follow-up of *Staphylococcus aureus* nasal carriage after 8 years: redefining the persistent carrier state. *J Clin Microbiol* 37: 3133–40.
- 19) O’Toole GA, Kolter R (1998): Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis. *Mol Microbiol* 28: 449–61.