



Monaldi Archives for Chest Disease

eISSN 2532-5264

<https://www.monaldi-archives.org/>

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The **Early Access** service lets users access peer-reviewed articles well before print / regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

The **Monaldi Archives for Chest Disease** is, therefore, e-publishing PDF files of an early version of manuscripts that have undergone a regular peer review and have been accepted for publication, but have not been through the typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear in a regular issue of the journal.

E-publishing of this PDF file has been approved by the authors.

All legal disclaimers applicable to the journal apply to this production process as well.

Monaldi Arch Chest Dis 2023 [Online ahead of print]

To cite this Article:

Sangma VSC, Jaggi S, Saini V, et al. **Prevalence of latent tuberculosis infection in the household contacts of pulmonary tuberculosis, time to treat.** *Monaldi Arch Chest Dis* doi: 10.4081/monaldi.2023.2563

 ©The Author(s), 2023
Licensee [PAGEPress](#), Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.



Prevalence of latent tuberculosis infection in the household contacts of pulmonary tuberculosis, time to treat

Vinnie Sarah Ch Sangma, Surabhi Jaggi, Varinder Saini, Deepak Aggarwal, Pankaj Kumar,
Jagdish Chander

Name of Department: Department of Pulmonary, Critical Care and Sleep Medicine,
Government Medical College and Hospital, Sector 32, Chandigarh

Corresponding author: Dr. Surabhi Jaggi, Department of Tuberculosis and Respiratory Diseases, D block, Level 5, Government Medical College and Hospital, Sector 32, Chandigarh-160030. **Email id:** surabhijaggi@gmail.com

1) Dr Vinnie Sarah Ch Sangma – MBBS, MD- Senior Resident, Department of Critical Care, Pushpawati Singhanian Hospital and Research Institute, New Delhi- 110017

2) Dr Surabhi Jaggi - MBBS, MD, DNB, IDCCM, EDARM - Assistant Professor, Department of Tuberculosis and Respiratory Diseases, Govt. Medical College and Hospital, Sector 32, Chandigarh-160030

3) Dr Varinder Saini - MBBS, MD -Professor and Head, Department of Pulmonary, Critical Care and Sleep Medicine, Government Medical College and Hospital, Sector 32, Chandigarh-160030

4) Dr Deepak Aggarwal - MBBS, MD (Tuberculosis & Respiratory Diseases), Professor, Department of Pulmonary, Critical Care and Sleep Medicine, Govt. Medical College & Hospital, Sector 32, Chandigarh -160030

5) Dr Pankaj Kumar – MBBS, MD (Paediatrics), Professor, Department of Paediatrics, Government Medical College and Hospital, Sector 32, Chandigarh -160030

6) Dr Jagdish Chander- MBBS, MD (Microbiology), Retired Professor and Head, Department of Microbiology, Government Medical College and Hospital, Sector 32, Chandigarh -160030

Keywords- tuberculosis; latent tuberculosis infection (LTBI); mantoux; chest x ray (CXR)

Contributions: VSCS, SJ, VS, DA, PK, JC helped in the treatment of the patients, helped in retrieving the data and conceived the content of the manuscript. VSCS and SJ drafted the manuscript. All authors read and revised the manuscript critically for important intellectual concepts and VS, DA, PK and JC approved the final version of the manuscript.

Conflict of interest: The authors declare that they have no competing interests, and all authors confirm accuracy.

Funding: None

Ethics Approval: The study protocol was reviewed and approved by the Institutional Ethics Committee, GMCH (Government Medical College and Hospital, Sector 32, Chandigarh, India.

Consent: Witten informed consent for the use of data was obtained from the index cases during the study.

Abstract

Treatment of latent pulmonary tuberculosis (TB) in the household contacts of TB has been included in the National Tuberculosis Elimination Program (NTEP) to achieve the target of TB elimination of Govt of India by 2025. However, there are no clear estimates of the prevalence of latent TB among the contacts that could suggest the impact of this intervention. The study was conducted to find the prevalence of and factors predicting latent TB among household contacts of pulmonary TB. All microbiological confirmed pulmonary TB patients registered between January 2020 to July 2021 and their household contacts were enrolled. All contacts underwent Mantoux testing to find the prevalence of latent TB. All symptomatic patients also underwent CXR and sputum examination to diagnose active pulmonary TB. Thereafter different demographic and clinical factors were evaluated to find predictors of latent TB using logistic regression model. A total of 118 pulmonary TB cases and their 330 household contacts were enrolled. The prevalence of latent TB and active TB among the contacts was found to be 26.36% and 3.03% respectively. The female gender of index TB cases was independently associated with a high proportion of latent TB cases in the family. (aOR-2.32; 95%CI-1.07-5.05; p=0.03). Neither the higher sputum smear positivity nor the severity of chest radiograph of index TB cases had any association with the number of contacts being diagnosed as latent TB or active TB. The results showed a significant prevalence of latent TB among household contacts with pulmonary TB. The severity of the disease in the index patient had no association with the prevalence of latent TB.

Introduction

Tuberculosis (TB), caused by the bacteria *Mycobacterium tuberculosis*, is one of the leading causes of death worldwide [1,2]. Depending on the host's immunity, the disease may present with a varying grade of severity ranging from asymptomatic infection (termed a latent TB) to progressive disease (active TB). An active case of pulmonary TB coughs out the infectious droplet nuclei, which can cause infection to others who come in contact with him unless the patient is detected early and started on treatment [3]. So, it is important to diagnose and treat such cases at the earliest.

Recently, the Government of India reframed the revised National Tuberculosis Control Program (RNTCP) and renamed it National Tuberculosis Elimination Program (NTEP) with the aim to eliminate tuberculosis from India by 2025. In line with it, the National Strategic Plan 2017-2025 was devised, integrating four strategic pillars of “DETECT-TREAT-PREVENT-BUILD, defining 4 key areas of action [4]. Under the “PREVENT” pillar, the program planned to include treatment of latent TB in the contacts of bacteriologically confirmed index cases and in individuals who are at high risk of getting TB disease [4]. This is an echo of The World Health Organization's (WHO) End TB Strategy which has incorporated the detection and treatment of latent TB as the key component to meet the targets.

Approximately one fourth of the world population has been estimated to have latent TB [5]. The prevalence of Latent TB has been highly variable across different geographical regions. This variability is partly explained by the difference in the methods used to diagnose latent TB, the population screened and partly by bacterial, host and environmental factors. There are no clear figures on the prevalence of latent TB in India which is one of the highest TB burden countries in the world. Few studies done across small geographical areas in India have yielded a prevalence

figure between 43-52% [6,7]. Hence, this study was conducted to evaluate the prevalence of latent and active TB through screening of the contacts of sputum-positive tuberculosis patients from this geographical area and to evaluate various predictors of latent TB.

Materials and Methods

This cross-sectional study was conducted in the Department of Pulmonary, Critical Care and Sleep Medicine, Government Medical College and Hospital, Chandigarh over a period of one and half years from January 2020 to July 2021. The subjects were microbiologically confirmed Pulmonary TB cases and their household contacts. Informed consent was taken from all the index cases as well as their household contacts. The study was approved by the institutional ethics committee of GMCH.

An index case was defined as a microbiologically confirmed pulmonary tuberculosis (drug-sensitive/drug-resistant) patient and “a household contact” was defined as a person living with the index case and sharing food from the same kitchen for a minimum of 3 months prior to the diagnosis of TB disease in the index case [8]. Among the index cases, extrapulmonary tuberculosis and those who did not give consent were excluded from the study. Among the household contacts, those who had past history of treated tuberculosis as well as those who did not give consent were excluded from the study.

All index cases were evaluated for demographic data, their baseline symptoms, smoking habit, chulha exposure, sputum smear grading for AFB and/or sputum CBNAAT and chest X-ray. Baseline chest x-ray of the index patients was classified into non-extensive disease and extensive

disease. The extensive disease was defined by the presence of bilateral cavitory disease or extensive parenchymal damage on chest radiography [9].

All household contacts of index cases were selected by verbal questioning with the index case. Demographic profile, symptoms and signs were noted using a structured questionnaire along with their clinical examination. Additionally, details about their known co-morbidities, immunocompromised states or therapy and smoking/alcohol use were noted. For the identification of presumptive TB four symptoms suggestive of TB i.e., productive cough of any duration, fever/night sweats, loss of weight and haemoptysis were asked as defined by the WHO [4].

After clinical assessment, all contacts underwent the Mantoux test. Latent tuberculosis was defined by Mantoux test reading of >10 mm induration measured after 48-72 hours of administration [10]. All symptomatic contacts also underwent chest X-ray PA view and sputum examination to evaluate for active TB disease. Chest radiography findings that were considered suggestive of active TB included infiltrates/consolidation, any cavitory lesion, any nodule with poorly defined margins, pleural effusion, hilar or mediastinal lymphadenopathy and military pattern.[10,11] In case of paediatric household contacts (<12 years), screening was done in the paediatric OPD by Mantoux test, chest x-ray and gastric lavage as per the advice of the paediatrician. Appropriate tests like fine needle aspiration of fluid/pus were sent for cytology and cartridge based nucleic acid amplification test (CBNAAT) in suspected extra-pulmonary tuberculosis. Active tuberculosis was confirmed in the contacts either on the basis of sputum smear microscopy/CBNAAT (microbiologically confirmed TB) or on the basis of relevant clinico-radiological findings (clinically diagnosed TB) [4]. Finally, the number of cases diagnosed as active tuberculosis and latent tuberculosis among the household contacts were analyzed in terms of their prevalence as

well as for any association with different demographic and clinical factors like sputum smear grade and disease extent on X-ray of the index patient.

Results

There was a total of 118 freshly diagnosed pulmonary TB patients and 330 household contacts that were enrolled in the study

Clinical profile of the index cases

The mean age of the index cases was 40.02±18.6 years with 39 (33%) patients presenting in the age group of 20-40 years. In contrast, 46% of contacts were in the age group of 20-40 years. Males comprised 62% of the TB patients in contrast to 54% of the household contacts. Smoking, diabetes and hypertension were more common among the index cases as compared to contacts. Among 118 TB patients, sputum high bacillary load and extensive involvement on X-ray were seen in 51 (43.2%) and 43 (36.4%) patients, respectively (Table 1). Approximately, one-fourth (23%) of the contacts were symptomatic at baseline. Cough was the most common symptom (13.0%) followed by fever/night sweat (4.2%), and significant weight loss (3.9%). Chest X-ray findings suggestive of TB were seen in 33 patients.

Comparison of contacts with and without latent TB

Out of 320 household contacts, 87 (26.4%) subjects had a positive Mantoux test and were categorized as latent TB; 10 patients (3%) were diagnosed with pulmonary TB on the basis of positive sputum AFB smear. There was no statistically significant difference in age, gender or smoking prevalence between contacts with or without latent TB (Table 2).

Association of latent TB with the TB among index cases

Out of 118 index patients, 41 had >50% of their household contacts diagnosed with latent TB and 77 (65.3%) had <50% of their contacts suffering from latent TB. Out of different demographic and clinical parameters, the female gender was independently associated with the higher (>50%) proportion of household contacts suffering from latent TB (aOR-2.32; 95%CI-1.07-5.05; p=0.034).

Discussion

The present study evaluated the prevalence of latent TB among household contacts of microbiologically proven pulmonary TB cases. The results showed that more than one-fourth of the contacts had latent TB which was independently associated with female gender of index TB cases. The global prevalence of latent TB has been estimated to be around 25% [5]. In India, there are no estimated figures on the prevalence of latent TB. However, in view of the high TB burden in India, it is likely that the figure will be higher than the global estimate. The present study showed the prevalence of latent TB to be 26.4% among household contacts with pulmonary TB which is lower than the global estimate. These lower figures need to be interpreted in light of the fact that they represent the data from a small geographical region and do not represent a country's prevalence. This prevalence was also found to be comparable to studies from California (30%) [11] and Scotland (31.2%) [12], but lower than the studies from India (52.6%) [6] and Malawi (66%) [13] that have high TB burden. The possible reasons for reduced prevalence may be the difference in the disease severity, socioeconomic status, smaller sample size and difference in the measurement tools used. Apart from the modifiable factors, Mantoux positivity significantly

depends on the host's immune response to tubercular protein. Hence, the Mantoux induration will be affected in immunocompromised states like diabetes, HIV, malnourished and patients on immunosuppressive therapy. The present study included subjects from household contacts of index patients which might also have created some amount of selection bias. Nevertheless, a relatively lower prevalence of latent TB found among household contacts in the present study undermines the contribution of seroconversion, secondary to exposure to fresh TB patients, in the prevalence of latent TB. In contrast to our results, a Chinese study from Shanghai showed a much lower prevalence of LTBI (17.2%) [14] in their study. This is possibly because Shanghai being China's one of the biggest city is inhabited by people with better socioeconomic status with greater awareness, better housing, less overcrowding and better health infrastructure.

The present study tried to find a causal association between latent TB and different demographic and clinical parameters of index patients as well as contacts. The results showed that latent TB had no association with the age and smoking status of either of the subject groups ($p > 0.05$). Also there was no gender difference in the prevalence of latent TB. However, it was seen that the female gender of index TB cases had a positive association with the higher proportion of household contacts being found to have latent TB (aOR-2.32; 95CI-1.07-5.05; $p = 0.03$). This association could be because the majority of the female patients were homemakers remaining at home throughout the day, resulting in prolonged exposure with the index patients leading to Mantoux positivity. In contrast, a study by Nair *et al.* found that male gender and being siblings with the index case (adjusted OR: 3.4 [95% CI.: 1.2, 9.7]) were important risk factors of an abnormal chest X-ray finding among the household contacts screened [8].

In the present study, the prevalence of active TB among household contacts was 3.03%. This was lower than the studies in Chattisgarh (10.1%) [15] and Chennai (5.3%) [8] in India and Tanzania

(6.4%) [16], Turkey (5.4%) [17] and Mandalay in Myanmar where a high prevalence of 13.8% active TB was found [18]. However, in other studies from India and California a much lower prevalence rate (1% and <1% respectively) was found [11,19]. The wide difference in the prevalence rates seen in different studies could be due to the difference in the sample size of the contacts screened ranging from 174 in Myanmar to 1556 in Chattisgarh [15,18]. The methods of screening also varied and even Gene X-pert was used in the studies conducted in Myanmar and Tanzania [16,18]. The studies from South Africa and Tanzania used sputum cultures for screening of the contacts [16,20].

The severity of tuberculosis is usually described on the basis of the grade of sputum positivity and the extent of involvement on X-ray. In the present study, 51 (43.2%) index cases had a higher sputum grading of >2+. However, on logistic regression analysis, higher sputum grading had no association with higher latent TB prevalence among the contacts. (OR-0.66; 95% CI-0.31-1.43); $p=0.29$). In case of active TB among contacts, 90% of the cases were the contacts of the index cases whose sputum smear was negative/scanty or 1+ ($p=0.03$). This is in contrast to a study conducted by Xu *et al.* where they found that the rate of active pulmonary tuberculosis among the contacts was positively associated with the amount of bacteria discharged in the sputum of the index cases ($r = 0.99, p<0.01$) [21].

In the present study chest radiograph showed extensive disease in 43 (36.4%) index TB patients. This was similar to the study in Chennai where the proportion of index cases with cavitory lesion on chest radiographs was 33% ($n=91$) [8]. In our study, the extensive disease on CXR in the index case was not a significant determinant of both latent TB (OR-1.18; 95%CI-0.54-2.6; $p=0.67$) and active TB among contacts. However, in a study from Malawi, there was an association between the cavitory disease/ the amount of lung field affected on CXR of the index case and the TST

positivity of the child contacts [13]. The difference in the findings could be attributed to the difference in the population groups included in the contact population.

The present study results showed a high prevalence of latent TB among household contacts of microbiologically confirmed pulmonary TB. A sample of 330 contacts provided adequate power to the results. The results suggest that the duration of exposure to the index case may be more important than the severity of the disease in determining latent TB. However, the study was conducted in a hospital setting among the contacts of those index cases who presented in the in-patient or the out-patient department and hence is not representative of the general population. Genotyping of the index cases and the contacts were not done to confirm whether the transmission had actually occurred in the household contacts leading to their Mantoux conversion or otherwise. Various challenges were also faced during the study including getting consent from the index cases to screen their contacts. This was due to the lack of awareness among the population about the need for early screening. Moreover, due to the ongoing COVID pandemic, socio-economic and household conditions of the index cases were not studied which may also have a strong influence on the prevalence of latent TB.

Conclusions

Latent TB is multi-factorial in origin. Hence, its prevalence has been highly variable across different studies both within and outside the same geographical region. Nevertheless, high prevalence, as has been seen in this study, highlights the need to actively diagnose and treat this form of TB. The same has been incorporated in the National Strategic Plan 2017-2025 by the Government of India. Active management of latent TB has been initiated in a phased manner in

India in order to achieve the TB elimination goal by 2025. Nevertheless, screening is a dynamic process and the programmes should regularly formulate decisions on when and how to screen, and which high-risk groups to be prioritized for management.

References

1. World Health Organization. Global tuberculosis report 2021. Accessed 4th January 2022. Available from: <http://www.who.int/teams/global-tuberculosis-programme/tb-reports>
2. Smith I. Mycobacterium tuberculosis pathogenesis and molecular determinants of virulence. *ClinMicrobiol Rev* 2003;16:463-96.
3. Doucette K, Cooper R. Tuberculosis. In: Grippi MA, Elias JA, Fishman JA, Kotloff RM, Pack AI, Senior RM, editors. *Fishman's Pulmonary Diseases and Disorders*, 5th ed. Mc Graw Hill; 2015. p. 2014-15.
4. World Health Organization. Systematic screening for active tuberculosis: principles and recommendations. World Health Organization; 2013. Accessed 5th November 2019. Available from: <https://www.who.int/tb/tbscreening>
5. Houben RM, Dodd PJ. The global burden of latent tuberculosis infection: a re-estimation using mathematical modelling. *PLoS Med* 2016;13:e1002152.
6. Krishnamoorthy Y, Ezhumalai K, Murali S, et al. Prevalence and risk factors associated with latent tuberculosis infection among household contacts of smear positive pulmonary tuberculosis patients in South India. *Trop Med Int Health* 2021;26:1645-51.
7. Praveen V. Prevalence of LTBI among household contacts of sputum positive TB patients receiving DOTS chemotherapy. *Indian J Tuberc* 2020;67:459-65.
8. Nair D, Rajshekhar N, Klinton JS, et al. Household contact screening and yield of tuberculosis cases—a clinic based study in Chennai, South India. *PLoS One* 2016;11:e0162090.
9. Ministry of Health and Family Welfare. Guidelines for Programmatic Management of Drug Resistant Tuberculosis in India, 2021. Accessed on 14th January 2022. Available from: <https://www.tbcindia.gov.in/>

10. Pai M, Joshi R, Kalantri S. Diagnosis of latent tuberculosis infection: Recent advances and future directions. In: Sharma SK, Mohan A, editors. Tuberculosis, 2nd ed. New Delhi: Jaypee Brothers Medical Publishers (P) Ltd; 2009. p. 186-99.
11. Sprinson JE, Flood J, Fan CS, et al. Evaluation of tuberculosis contact investigations in California. *The Int J Tuberc Lung Dis* 2003;7:363-8.
12. Mandal P, Craxton R, Chalmers JD, et al. Contact tracing in pulmonary and non-pulmonary tuberculosis. *QJM* 2012;105:741-7.
13. Hector J, Anderson ST, Banda G, et al. TST positivity in household contacts of tuberculosis patients: a case-contact study in Malawi. *BMC Infect Dis* 2017;17:259.
14. Xu J, Hu Y, Jiang W, et al. [Prevalence and risk factors of latent tuberculosis infection in close contacts of tuberculosis patients among non-resident populations in Shanghai, China]. [Article in Chinese]. *Zhonghua Jie He He Hu Xi Za Zhi* 2016;39:25-9.
15. Khaparde K, Jethani P, Dewan PK, et al. Evaluation of TB case finding through systematic contact investigation, Chhattisgarh, India. *Tuberc Res Treat* 2015;2015:670167.
16. Beyanga M, Kidenya BR, Gerwing-Adima L, et al. Investigation of household contacts of pulmonary tuberculosis patients increases case detection in Mwanza City, Tanzania. *BMC Infect Dis* 2018;18 110.
17. Kilicaslan Z, Kiyani E, Kucuk C, et al. Risk of active tuberculosis in adult household contacts of smear-positive pulmonary tuberculosis cases. *Int J Tuberc Lung Dis* 2009;13:93-8.
18. Htet KK, Liabsuetrakul T, Thein S, et al. Improving detection of tuberculosis among household contacts of index tuberculosis patients by an integrated approach in Myanmar: a cross-sectional study. *BMC Infect Dis* 2018;18:660.
19. Velayutham B, Jayabal L, Watson B, et al. Tuberculosis screening in household contacts of pulmonary tuberculosis patients in an urban setting. *PloS One* 2020;15:e0240594
20. Little KM, Msandiwa R, Martinson N, et al. Yield of household contact tracing for tuberculosis in rural South Africa. *BMC Infect Dis* 2018;18:299.
21. Xu C, Hu B. [Prevalence of active pulmonary tuberculosis among household contacts of recently diagnosed pulmonary tuberculosis patients with positive sputum-smear]. [Article in Chinese]. *Zhonghua Jie He He Hu Xi Za Zhi* 2008;29:693-5.

Table 1. Demographic and clinical characteristics of sputum smear positive index cases (n=118).

S.no	Variable		Index cases (n=118)	Household contacts (n=320)
1	Age(years)		40.02±18.6	31.90 ± 16.44
2	Males		73 (61.9)	176 (53.3%)
3	Smoking		42 (35.6)	56 (17%)
4	Chulha exposure		13 (11)	37 (11.2%)
5	Diabetes		19 (16.1)	8 (2.4%)
6	Hypertension		9 (7.6)	19 (5.8%)
7	Previous TB		33 (28)	
8	Sputum smear grading	Negative/ Scanty/ 1+	67 (56.8)	
		2+/3+/4+	51 (43.2)	
9	Chest X-ray findings	Extensive disease	43 (36.4%)	
		Non extensive disease	75 (63.3)	

Table 2. Comparison of different characteristics between contacts with and without latent tuberculosis.

	Latent TB (n=87)	Normal (excluding active TB) (n=233)	p-value
Age (yrs)	32.8±14.4	31.6±17.1	0.55
Male	45 (51.7%)	127 (54.5%)	0.70
Smoking	19	33	0.10
Chulha exposure	6 (16.7%)	30 (83.3%)	0.13
Comorbidity	10 (37.0%)	17 (63.0%)	0.23

Table 3. Association of latent TB with demographic and clinical parameters of index TB cases.

	Independent variable	Univariate analysis OR (95% CI)	p-value	Multivariate analysis aOR (95% CI)	p-value
1	Age	0.98 (0.96-1.003)	0.08		
2	Female gender	2.32 (1.07-5.05)	0.034	2.32 (1.07-5.05)	0.034
3	High AFB group	0.66 (0.31-1.43)	0.29		
4	Extensive disease on CXR	1.18 (0.54-2.6)	0.67		