

5-1-2008

# An Evaluation of Standard, Alternative, and Robust Slope Test Strategies


Tim Moses

*Educational Testing Service, Princeton, NJ, mosest@optonline.net*

Alan Klockars

*University of Washington, klockars@u.washington.edu*

Follow this and additional works at: <http://digitalcommons.wayne.edu/jmasm>

 Part of the [Applied Statistics Commons](#), [Social and Behavioral Sciences Commons](#), and the [Statistical Theory Commons](#)

## Recommended Citation

Moses, Tim and Klockars, Alan (2008) "An Evaluation of Standard, Alternative, and Robust Slope Test Strategies," *Journal of Modern Applied Statistical Methods*: Vol. 7: Iss. 1, Article 7.

Available at: <http://digitalcommons.wayne.edu/jmasm/vol7/iss1/7>

This Regular Article is brought to you for free and open access by the Open Access Journals at DigitalCommons@WayneState. It has been accepted for inclusion in Journal of Modern Applied Statistical Methods by an authorized administrator of DigitalCommons@WayneState.

## An Evaluation of Standard, Alternative, and Robust Slope Test Strategies

Tim Moses  
Educational Testing Service

Alan Klockars  
University of Washington

---

The robustness and power of nine strategies for testing the differences between two groups' regression slopes under nonnormality and residual variance heterogeneity are compared. The results showed that three most robust slope test strategies were the combination of the trimmed and Winsorized slopes with the James second order test, the combination of Theil-Sen with James, and Theil-Sen with percentile bootstrapping. The slope tests based on Theil-Sen slopes were more powerful than those based on trimmed and Winsorized slopes.

Key words: slopes, least squares, Theil-Sen, robust regression, James second order, nonnormality, residual variance heterogeneity

---

### Introduction

The question of whether group differences are constant or vary across levels of an individual difference variable (X) has been considered in many fields of social science, including clinical psychology (Dance & Neufeld, 1988), organizational research (Aguinis & Pierce, 1998; Hunter, Schmidt, & Hunter, 1979), learned helplessness (Seligman, 2002) and education's search for Aptitude-Treatment Interactions (Cronbach & Snow, 1977). Several strategies have been proposed for evaluating the consistency of group differences across X based on fitting regression lines that predict outcome Y from X in separate treatment groups and then conducting a significance test for the homogeneity of the groups' regression slopes.

---

Tim Moses is a Senior Psychometrician at Educational Testing Service where he works on several testing programs. Tim completed his PhD in Educational Psychology at the University of Washington. E-mail him at [mosest@optonline.net](mailto:mosest@optonline.net). Alan Klockars is Professor of Educational Psychology at the University of Washington. His research concerns multiple comparisons and, more recently, methods of conducting ATI research. E-mail him at [klockars@u.washington.edu](mailto:klockars@u.washington.edu).

The purpose of this study is to compare some of the recently-researched methods of slope estimation and testing under conditions of nonnormality and residual variance heterogeneity.

The slope test strategies considered in this study are approaches to estimating the following model,

$$Y_{ij} = \beta_{0j} + \beta_{1j}X_{ij} + \varepsilon_{ij}, \quad (1)$$

where outcome Y for individual i (= 1 to N) in group j (= 1 to J) is a linear function of a continuous X,  $\beta_{0j}$  and  $\beta_{1j}$  are the population intercepts and slopes of the regression line for each of J groups, and the  $\varepsilon_{ij}$  are the residuals. The strategies for assessing differences in the  $\beta_{1j}$ 's reviewed below are most easily understood in terms of alternative expressions of (1). When J = 2, (1) can be expressed as,

$$Y_{ij} = \beta_0 + \beta_1X_{ij} + \beta_2G_{ij} + \beta_3X_{ij}G_{ij} + \varepsilon_{ij}, \quad (2)$$

where  $G_{ij}$  is a dichotomously-coded group membership variable. A more general matrix version of (1) and (2) is,

$$Y = X\beta + e, \quad (3)$$

## STANDARD, ALTERNATIVE, AND ROBUST SLOPE TEST STRATEGIES

where  $\mathbf{Y}$  is an  $N$  by 1 column vector,  $\mathbf{X}$  is an  $N$  by  $K$  “design matrix” corresponding to the  $K$   $\beta$ 's (including a column of 1's for estimating  $\beta_0$ ),  $\boldsymbol{\beta}$  is a  $K$  by 1 column vector of  $\beta$ 's and  $\mathbf{e}$  is an  $N$  by 1 column vector of residuals.

Standard slope estimation and slope test

The standard slope test uses “least squares” estimates of the  $\beta$ 's (i.e.,  $\hat{\beta}$ 's) that minimize the sum of the squared residuals,  $\hat{\mathbf{e}}'\hat{\mathbf{e}} = (\mathbf{Y} - \mathbf{X}\hat{\boldsymbol{\beta}})'(\mathbf{Y} - \mathbf{X}\hat{\boldsymbol{\beta}})$ . Because  $\hat{\mathbf{e}}'\hat{\mathbf{e}}$  is a convex function of  $\hat{\boldsymbol{\beta}}$ , it can be minimized by differentiating with respect to  $\hat{\boldsymbol{\beta}}$ , setting this derivative to zero and solving for  $\hat{\boldsymbol{\beta}}$ , resulting in the closed form solution

$$\hat{\boldsymbol{\beta}} = (\mathbf{X}'\mathbf{X})^{-1} \mathbf{X}'\mathbf{Y} . \quad (4)$$

Equivalently, group  $j$ 's slope can be estimated

$$\hat{\beta}_{1j} = \frac{\sum_{i \in j} (x_{ij} - \bar{x}_j)(y_{ij} - \bar{y}_j)}{\sum_{i \in j} (x_{ij} - \bar{x}_j)^2}, \quad (5)$$

where  $\bar{x}_j$  and  $\bar{y}_j$  are the means of  $X$  and  $Y$  in group  $j$ .

The standard test for assessing the differences of  $J$  slopes is an  $F$  test,

$$F_{\text{SlopesStandard}} = \frac{\left(\frac{1}{J-1}\right) \sum_j \left( \hat{\beta}_{1j}^2 - \hat{\beta}_{1,\text{Standard}}^2 \right) \sum_{i \in j} (x_{ij} - \bar{x}_j)^2}{\left(\frac{1}{N-2J}\right) \sum_j (N_j - 2) \hat{\sigma}_{ej}^2} \quad (6),$$

where  $\hat{\sigma}_{ej}^2 = \frac{\sum_{i \in j} (\hat{e}_{ij})^2}{N_j - 2}$  and

$$\hat{\beta}_{1,\text{Standard}} = \frac{\sum_j \hat{\beta}_{1j} \sum_{i \in j} (x_{ij} - \bar{x}_j)^2}{\sum_j \sum_{i \in j} (x_{ij} - \bar{x}_j)^2}$$

is the variance-

weighted common slope (Myers & Well, 1995, p. 421-422). (6) is evaluated on an  $F$  distribution with  $J-1$  and  $N-2J$  degrees of freedom. With  $J=2$ , a  $t$ -test of  $\hat{\beta}_3$  in model (2) that is equivalent to the  $F$  test in (6) can be conducted by obtaining

the standard error of  $\hat{\beta}_3$  as the square root of one of the diagonal elements in the variance-covariance matrix of  $\hat{\boldsymbol{\beta}}$ ,  $\frac{\hat{\mathbf{e}}'\hat{\mathbf{e}}}{N-K}(\mathbf{X}'\mathbf{X})^{-1}$ , and

evaluating  $\frac{\hat{\beta}_3}{\text{SE}(\hat{\beta}_3)}$  on a  $t$  distribution with

$N-K = N-4$  degrees of freedom. The referencing of the standard test statistics to  $F$  and  $t$  distributions is justified when the data meet particular assumptions, namely that the  $\epsilon_{ij}$  are normally and independently distributed with equal variances across the  $J$  groups.

The standard methods for estimating and testing slopes are problematic when data are nonnormal and residual variances are heterogeneous (Conover & Iman, 1982; Conerly & Mansfield, 1988; Headrick & Sawilowsky, 2000; Klockars & Moses, 2002; Dretzke, Levin & Serlin, 1982; Overton, 2001; Alexander & Deshon, 1994; Deshon & Alexander, 1996). When distributions exhibit heavy-tailed nonnormality, extreme scores occur more often than when distributions are normal, increasing the variability of the estimated slopes, reducing the estimated standard errors, and making the standard test excessively liberal. When groups' residual variances and sample sizes differ, the standard test's pooling of groups' residual variances,  $\left(\frac{1}{N-2J}\right) \sum_j (N_j - 2) \hat{\sigma}_{ej}^2$ , is

problematic, making the standard slope test either liberal or conservative depending on whether the larger and smaller group has the larger or smaller residual variance. The inaccuracy of the standard test is disturbing given that nonnormality and residual variance heterogeneity appear to be common in actual data (Micceri, 1989; Aguinis, Peterson & Pierce, 1999). What follows are detailed definitions of slope test strategies that may outperform the standard test when distributions are nonnormal and residual variances are heterogeneous.

Slope tests for nonnormal data: Central tendency strategies

Two approaches to slope estimation view group  $j$ 's slope in (5),

$$\hat{\beta}_{1j} = \frac{\sum_{i \in j} (x_{ij} - \bar{x}_j)(y_{ij} - \bar{y}_j)}{\sum_{i \in j} (x_{ij} - \bar{x}_j)^2}, \text{ as a central value of}$$

the slopes that can be created from pairs of observations in the data,

$$b_{1,ij} = \frac{(y_{ij} - y_{i'j})}{(x_{ij} - x_{i'j})}, i \neq i', x_{ij} \neq x_{i'j}, \text{ and then try}$$

reduce the influence of the extreme observations on the central value. These ‘central tendency’ approaches define extreme observations in terms of both X and Y, so that the screening of extreme observations caused by nonnormality could potentially address slope estimation problems such as leverage (observations that are extreme on X), discrepancy (observations that are extreme with respect to the regression line), and outliers on Y. One popular strategy is the Theil-Sen slope estimator (Theil, 1950; Sen, 1968; Wilcox, 2004; Wilcox & Keselman, 2004; Ebrahem & Al-Nasser, 2005; Wang, 2005). The Theil-Sen estimate is the median of the slopes that can be computed from the  $N_j(N_j-1)/2$  pairs of observations in the data. Percentile bootstrapping methods can be used to test for differences between groups’ Theil-Sen slopes (i.e., draw 599 random samples with replacement from the  $J = 2$  datasets, compute the differences in Theil-Sen slopes in each of these datasets, and determine if the middle  $(1-\alpha)\%$  of the 599 slope differences contain zero, Wilcox, 2005).

A less-familiar alternative to the Theil-Sen slope estimate is the application of the trimming and Winsorizing strategies that are typically proposed in tests of mean differences to  $\frac{(y_{ij} - y_{i'j})}{(x_{ij} - x_{i'j})}$  (Guo, 1996; Luh & Guo, 2000). To obtain trimmed and Winsorized estimates of slopes and their variances, rank order the  $x$ ’s in each of the  $J$  groups,  $x_{1j} < x_{2j} \dots < x_{N_jj}$ . When the number of observations in group  $j$  is even ( $N_j=2m_j$ ) consider  $m_j$  independent slope estimates,

$$b_{1,i+m_j,j} = \frac{(y_{i+m_j,j} - y_{ij})}{(x_{i+m_j,j} - x_{ij})}. \quad (7)$$

When the number of observations in  $j$  is odd ( $N_j=2m_j+1$ ), a pooling is done so that

observations  $y_{2m_j}$  and  $y_{2m_j+1j}$  are pooled,  $x_{2m_j}$  and  $x_{2m_j+1j}$  are pooled, and  $y_{2m_j}$  and  $x_{2m_j}$  are replaced by  $y_{2m_j} = (y_{2m_j} + y_{2m_j+1j})/2$  and  $x_{2m_j} = (x_{2m_j} + x_{2m_j+1j})/2$ .

The trimming and Winsorizing is done for each of the  $j$  slopes and standard errors. Let  $g_j = \gamma m_j$  where  $\gamma$  represents the proportion of observations to be trimmed from each tail of the ordered distribution of  $b_{1,lj}, l = 1$  to  $m_j, b_{1,lj} \leq b_{1,2j} \leq \dots b_{1,m_jj}$ . Let  $h_j = m_j - 2g_j$  be the effective sample size after trimming.

The trimmed mean slope is computed as

$$\bar{b}_{1,j} = \frac{\sum_{l=g_j+1}^{m_j-g_j} b_{1,lj}}{h_j}. \quad (8)$$

Winsorized slope observations are obtained by,

$$bw_{1,lj} = \left\{ \begin{array}{ll} b_{1,(g_j+1)j} & \text{if } b_{1,lj} \leq b_{1,(g_j+1)j} \\ b_{1,lj} & \text{if } b_{1,(g_j+1)j} < b_{1,lj} < b_{1,(m_j-g_j)j} \\ b_{1,(m_j-g_j)j} & \text{if } b_{1,lj} \geq b_{1,(m_j-g_j)j} \end{array} \right\}$$

(9). The variance of the trimmed mean slope is computed as a function of the Winsorized variance,

$$\sigma_{bwj}^2 = \frac{1}{h_j(h_j-1)} \sum_{l=1}^{m_j} (bw_{1,lj} - \frac{\sum bw_{1,lj}}{m_j})^2. \quad (10)$$

To assess the differences in trimmed slopes, replace the  $\hat{\beta}_{1j}$  in (6) with  $\bar{b}_{1,j}$ , the

$$\left( \frac{1}{N-2J} \right) \sum_j (N_j - 2) \hat{\sigma}_{ej}^2 \quad \text{with}$$

$$\left( \frac{1}{\sum_j h_j - J} \right) \sum_j h_j (h_j - 1) \sigma_{bwj}^2, \quad \text{and the}$$

$\sum_{i \in j} (X_{ij} - \bar{X}_j)^2$  with  $h_j$ . These replacements to

(6) cause the standard test of slope differences to resolve into an F test for independent trimmed means with  $J-1$  and  $\sum_j h_j - J$  degrees of freedom,

$$F_{\text{SlopesTrimmed}} = \frac{\left( \frac{1}{J-1} \sum_j h_j \left( \bar{b}_{ij}^2 - \left( \frac{\sum_j h_j \bar{b}_{ij}}{\sum_j h_j} \right)^2 \right) \right)}{\left( \frac{1}{\sum_j h_j - J} \sum_j h_j (h_j - 1) \sigma_{\text{bwi}}^2 \right)}. \quad (11)$$

Slope tests for nonnormal data: MM Regression

In “minimum maximum likelihood type” (MM, Yohai, 1987) regression, extreme observations are addressed in the minimization process used to estimate the regression line. While the standard slope estimation process is based on minimizing the sum of all squared residuals, the robust regression paradigm views the least squares approach as one of several possible functions,  $\xi$ , of the scaled residuals that could be minimized,

$$\sum_j \sum_{i \text{ in } j} \xi \left( \frac{\varepsilon_{ij}}{\sigma} \right). \quad (12)$$

Some choices of  $\xi$  can produce  $\beta$  estimates that outperform the standard method’s  $\beta$ ’s in terms of their “breakdown” rates (i.e., the smallest percentage of contaminated observations needed to render  $\hat{\beta}$  useless). One popular  $\xi$  (SAS, 2003) is the Tukey weight function,

$$\xi(s) = \begin{cases} 3\left(\frac{s}{\kappa}\right)^2 - 3\left(\frac{s}{\kappa}\right)^4 + \left(\frac{s}{\kappa}\right)^6 & \text{if } |s| \leq \kappa, \\ 1 & \text{otherwise.} \end{cases} \quad (13)$$

In (13),  $\kappa$  is a constant selected to obtain desirable properties. A  $\kappa$  value of 3.44 results in parameter estimates that are 85% as efficient as least squares estimates when the data are normal (Holland & Welsh, 1977). When data contain outliers that are discrepant with respect to the regression line,  $\kappa$  defines a range around which the observations outside of the range have reduced contribution to the slope estimates.

The search for  $\beta$ ’s that minimize (12) is similar to the standard test in that  $\beta_k$ ’s are found such that the derivatives of (12) with respect to

the  $\beta_k$ ’s are zero,

$$\sum_j \sum_{i \text{ in } j} \frac{\partial \xi}{\partial s} (s_{ijk}) x_{ijk} = 0, k = 1 \text{ to } K. \quad \text{Unlike the}$$

least squares estimation methods used with the standard test, with MM regression there are no closed-form solutions to minimizing (12). The following is an outline of the three-stage MM algorithm for estimating the  $\beta_k$ ’s.

The first step of MM regression is to obtain robust starting values for the  $\beta_k$ ’s and  $\sigma$ . The current SAS procedure for MM uses Least Trimmed Squares estimates as starting values (Rousseeuw, 1984; SAS Institute, 2003). The basic idea of LTS estimation is to draw samples of  $K$  observations from the  $N$  total observations in the data set. In each sample, obtain least squares estimates of the  $\beta_k$ ’s and find the ones

$$\text{that minimize } \sum_i^h (\varepsilon_i)^2, \text{ where } h = \frac{3N + K + 1}{4}$$

and observations  $i$  through  $h$  reference the  $h$  smallest squared residuals. Additional features of the LTS algorithm involve intercept adjustments that reduce  $\sum_i^h (\varepsilon_i)^2$  and

computational search processes designed to find final  $\beta_k$  estimates quickly in extremely large datasets (Rousseeuw & Van Driessen, 2000). One preliminary estimate of  $\sigma$  is computed as,

$$s_{\text{LTS}} = d \sqrt{\frac{1}{h} \sum_i^h (\varepsilon_i)^2}, \quad (14)$$

$$\text{where } d = 1 / \sqrt{1 - \frac{2N}{hc} \phi(1/c)}, \quad c = 1 / \Phi\left(\frac{h+N}{2N}\right),$$

and  $\Phi$  and  $\phi$  are the cumulative and probability density functions of the standard normal distribution.

A more efficient estimate of  $\sigma$  than  $s_{\text{LTS}}$  can also be computed,

$$W_{\text{scale}} = \sqrt{\frac{\sum_i^N w_i (\varepsilon_i)^2}{\sum_i w_i - K}}, \quad (15)$$

$$\text{where } w_i = \begin{cases} 0 & \text{if } |\varepsilon_i| / s_{\text{LTS}} > 3 \\ 1 & \text{otherwise} \end{cases}.$$

With initial estimates of the  $\beta_k$ 's and  $\sigma$ , the second step is to conduct iterative calculations to produce a converged  $\sigma$  value,

$$(\sigma^{m+1})^2 = \frac{1}{(N-K) \left( \int \xi(s) \partial \Phi(s) \right)} \sum_i^N \xi \left( \frac{\varepsilon_i}{\sigma^m} \right) (\sigma^m)^2,$$

(16) where  $\int \xi(s) \partial \Phi(s)$  denotes an expected value of  $\xi(s)$  when the  $s$  are from a normal distribution (about .25 for the Tukey bisquare  $\xi(s)$  with  $\kappa = 2.9366$ ). In (16), setting  $\kappa = 2.9366$  results in the  $\sigma$  having a breakdown rate of 25% (SAS, 2003).

The third step is to conduct an iterative search for a final solution of the  $\beta_k$ 's with a fixed  $\sigma$  value

$$\beta^{m+1} = (\mathbf{X}^t \mathbf{\Omega} \mathbf{X})^{-1} \mathbf{X}^t \mathbf{\Omega} \mathbf{Y}, \quad (17)$$

where  $\mathbf{\Omega}$  is an  $N$  by  $N$  matrix with diagonal entries  $\frac{\partial \xi(s)}{\partial s} \frac{1}{s}$  where the  $s$  are the scaled residuals from the  $m$ th iteration step and  $\kappa = 3.44$  in default SAS routines (SAS, 2003). The entries for  $\mathbf{\Omega}$  are the "reweighted" part of MM's iteratively reweighted least squares algorithm, and for the Tukey  $\xi(s)$  given in (13) are known as the Tukey bisquare weight function.

At convergence, there are several estimates of the asymptotic variance-covariance matrix of  $\beta$  (SAS, 2003). One version is,

$$\left( 1 + \frac{K}{N \left( \frac{1}{N} \sum_i \partial^2 \xi(\varepsilon_i) / \partial^2 \varepsilon \right)^2} \right) \frac{(1/(N-K) \sum_i \partial \xi(\varepsilon_i) / \partial \varepsilon)^2}{\left( \frac{1}{N} \sum_i \partial^2 \xi(\varepsilon_i) / \partial^2 \varepsilon \right)} \mathbf{W}^1, \quad (18)$$

where  $\left( 1 + \frac{K}{N \left( \frac{1}{N} \sum_i \partial^2 \xi(\varepsilon_i) / \partial^2 \varepsilon \right)^2} \right)$  is a

correction factor,  $\partial^2 \xi(\varepsilon) / \partial^2 \varepsilon$  is the second derivative of  $\xi$  with respect to the residuals, and

$\mathbf{W}$  is a  $K$  by  $K$  matrix with entries  $\mathbf{W}_{kk'} = \sum_i \left( \partial^2 \xi(\varepsilon_i) / \partial^2 \varepsilon \right) x_{ik} x_{ik'}$ .

The preceding review provides some insight into the kinds of nonnormality problems for which MM might be especially useful, which are probably situations with outliers that do not "mask" themselves by exerting heavy influence on the regression line. Many of the steps of the MM estimation process are analogues to the standard method's estimation, including the use of least squares estimation used in the LTS starting values, the computation of the  $\beta_k$ 's (equation 17 is a weighted version of equation 4), and the computation of the MM standard errors ( $\mathbf{W}$  in equation 18 is a weighted version of  $(\mathbf{X}^t \mathbf{X})$  in  $\frac{\hat{\mathbf{e}}^t \hat{\mathbf{e}}}{N-K} (\mathbf{X}^t \mathbf{X})^{-1}$ ). The relatedness of

MM computations to the standard method's computations suggest that both procedures would do well with normal populations, while MM should outperform the standard method when there are outliers on  $Y$  (Anderson & Schumacker, 2003).

Slope tests for heterogeneous residual variances

Alternative parametric significance tests have been developed by Welch (1938), James (1951) and Deshon and Alexander (1994) to test for slope differences when residual variances are unequal. All three methods avoid the standard test's pooling of groups' residual variances in (6). Comparative research has shown that the three parametric alternative tests perform similarly in terms of robustness and power (Luh & Guo, 2000; Luh & Guo, 2002; Deshon & Alexaner, 1996), so this study focuses solely on the James second-order test, which is slightly better than the Welch and Deshon and Alexander tests in terms of power and robustness to nonnormality.

The steps of the James second order test are as follows:

- 1) Define a James weight,  $w_{j_j}$ , based on each group slope's standard error,

$$w_{j_j} = \frac{1/\sigma_{\beta_j}^2}{\sum_j 1/\sigma_{\beta_j}^2}. \quad (19)$$

## STANDARD, ALTERNATIVE, AND ROBUST SLOPE TEST STRATEGIES

- 2) Define a variance-weighted common slope as,

$$\beta^+ = \sum_j w_{j,j} \beta_{1j} \quad (20)$$

- 3) Define the James' test statistic as,

$$\text{James} = \sum_j \frac{(\beta_{1j} - \beta^+)^2}{\sigma_{\beta_{1j}}^2} \quad (21)$$

- 4) Evaluate the significance of the James' test statistic by determining if it exceeds the following critical value,

$$\begin{aligned} \text{James}_{\text{crit}} = & c + (1/2)(3\chi_4 + \chi_2) \sum_j [(1 - w_j)^2 / v_j] \\ & + (1/16)(3\chi_4 + \chi_2)^2 [1 - (J-3)/c] \sum_j [(1 - w_j)^2 / v_j]^2 \\ & + (1/2)(3\chi_4 + \chi_2) [(8R_{23} - 10R_{22} + 4R_{21} - 6R_{12}^2 + 8R_{12}R_{11} \\ & - 4R_{11}^2) + (\chi_2 - 1)(2R_{23} - 4R_{22} + 2R_{21} - 2R_{12}^2 + 4R_{12}R_{11} - 2R_{11}^2) \\ & + (1/4)(3\chi_4 - 2\chi_2 - 1)(4R_{12}R_{11} - R_{12}^2 - 2R_{12}R_{10} - 4R_{11}^2 \\ & + 4R_{11}R_{10} - R_{10}^2)] + (5\chi_6 + 2\chi_4 + \chi_2)(R_{23} - 3R_{22} + 3R_{21} - R_{20}) \\ & + (3/16)(35\chi_8 + 15\chi_6 + 9\chi_4 + 5\chi_2)(R_{12}^2 - 4R_{23} + 6R_{22} - 4R_{21} \\ & + R_{20}) + (1/16)(9\chi_8 - 3\chi_6 - 5\chi_4 - \chi_2)(4R_{21} - 2R_{22} - R_{20} + 2R_{12}R_{10} \\ & - 4R_{11}R_{10} + R_{10}^2) + (1/4)(27\chi_8 + 3\chi_6 + \chi_4 + \chi_2)(R_{11}^2 - R_{22}) \\ & + (1/4)(45\chi_8 + 9\chi_6 + 7\chi_4 + 3\chi_2)(R_{23} - R_{12}R_{11}) \end{aligned}$$

(22), where  $v_j = N_j - 2$ ,  $c$  is the  $1-\alpha$  quantile of the central chi-square distribution with  $J-1$  degrees of freedom,  $R_{\text{ut}} = \sum_j \frac{w_{j,j}^t}{v_j^u}$  and

$$\chi_{2s} = \frac{c^s}{\prod_{q=1}^s (J + 2q - 3)} \quad (\text{for } \chi_2, \chi_4, \chi_6, \text{ and } \chi_8,$$

$s$  is 1, 2, 3, and 4, respectively).

Hybrid slope tests for nonnormal data and heterogeneous residual variances

Slope test strategies are not necessarily robust to problems for which they were not directly designed. The parametric alternative strategies that were designed to address residual variance heterogeneity have documented problems with nonnormal data (Deshon & Alexander, 1996). The slope test strategies that have been proposed for nonnormal data do not directly address residual variance heterogeneity.

An important area of research assesses so-called hybrid slope test strategies that may be robust to several assumption violations by use of nonnormality-robust group slopes and standard errors with parametric alternative tests that avoid the pooling of heterogeneous residual variances.

Recent research on hybrid slope test strategies has considered using standard slope estimates and standard errors or trimmed slope estimates and Winsorized standard errors with skew-corrected versions of parametric alternative tests (Luh & Guo, 2000; 2002). The use of the trimmed slopes and Winsorized standard errors with parametric alternative tests like James is straightforward, with groups' degrees of freedom calculated as  $v_j = h_j - 1$  rather than as  $N_j - 2$ . Luh and Guo also transformed the test-statistics of the parametric alternatives to eliminate the effect of skewness (Johnson, 1978; Hall, 1992). For example, the proposed transformation for skewness for the James second order test statistic from (21) is,  $\text{James\_TT} =$

$$\sum_j \left[ \left( \frac{\sqrt{N_j} [(\beta_{1j} - \beta^+) / \sqrt{N_j} \sigma_{\beta_{1j}}] - \gamma_{x,j}^3 \gamma_{\varepsilon,j}^3 (\beta_{1j} - \beta^+)^2 / N_j \sigma_{\beta_{1j}}^2}{6 + \gamma_{x,j}^3 \gamma_{\varepsilon,j}^3 / (6N_j)} \right) \right]^2, \quad (23)$$

where  $\gamma_{x,j}^3$  and  $\gamma_{\varepsilon,j}^3$  are the sample skews of  $X$  and  $\varepsilon$  in group  $j$ . Luh and Guo's studies showed that their hybrid strategies were robust to both nonnormality and residual variance heterogeneity.

This study

This study extends prior research on the relative performance of slope testing strategies under nonnormality and residual variance heterogeneity. This study directly compares the standard, MM, and Theil-Sen tests, extending the previous comparisons based on estimating one slope that have given recommendations for MM regression over the standard method (Anderson & Schumaker, 2003) and for Theil-Sen over MM regression and the standard method (Wilcox & Keselman, 2004). The comparison of the trimmed and Winsorized slope test with the Theil-Sen and MM methods

has not been considered in previous studies, and it allows for an evaluation of some trimming (trimmed and Winsorized) with the most extreme trimming possible (Theil-Sen).

This study also extends Luh and Guo's (2000, 2002) work, first by separately evaluating the trimmed and Winsorized slope test and the skewness transformation of the James test statistic. Because the accuracy of slope estimation has more to do with the heaviness of the distribution's tails rather than its skew (Klockars & Moses, 2002), the test statistic transformation ought to have a smaller impact in correcting for nonnormality than the trimmed and Winsorized, Theil-Sen and MM methods. Finally, Luh and Guo's efforts to form hybrid slope test strategies that are robust to both nonnormality and residual variance heterogeneity are extended to consider hybrid slope tests based not only on integrating the trimmed and Winsorized methods and the skewness transformation with James second order method, but also the MM and Theil-Sen methods.

Methodology

A simulation study was conducted to investigate the relative robustness and power of the slope test strategies for comparing two groups' slopes. Empirical rejection rates of the null hypothesis were computed based on 10,000 replications for each condition. Two treatment groups were used throughout the study. The following conditions were considered.

Slope Test Strategies

Five stand-alone slope test strategies and four hybrids of the five strategies were evaluated.

- 1) The standard F test of slope differences in (6) (Standard).
- 2) The James parametric alternative test in (21) (James).
- 3) Significance testing of the  $\beta_3$  in model (2) based on MM estimation with the default settings in SAS PROC ROBUSTREG (SAS Institute Inc., 2003) (MM).
- 4) The trimmed and Winsorized slope test in (11) using 10% trimming (TW).

- 5) The Theil-Sen estimator with percentile bootstrapping for the significance testing (TS).

The following four hybrid strategies were also considered:

- 6) The James procedure with the Johnson's one-sample t-statistic transformation for skewness in (23) (James-TT).
- 7) The James procedure using MM slope estimates and standard errors (James-MM).
- 8) The James procedure using 10% trimmed slope estimates and Winsorized standard errors from Luh and Guo (2000) (James-TW).
- 9) The James procedure using the Theil-Sen slope estimates and the standard deviations of 599 bootstrapped Theil-Sen estimates from strategy 5 for the group slopes' standard errors (James-TS).

Defining groups' observations and degrees of freedom

For the James-MM and James-TS strategies, some consideration was given for defining the groups' degrees of freedom. Initial efforts were based on Luh and Guo's (2000) attempt to account for the number of observations used in the slope estimate in James-TW ( $v_j = h_j - 1$ ). Directly applying this to James-TS would mean setting  $v_j = 2 - 1$ . From initial results it was clear that using  $v_j = 1$  resulted in extremely conservative tests for James-TS, so in an effort to obtain more reasonable results, the  $v_j$  was set as  $N_j(N_j - 1)/2 - 1$ . For James-MM, degrees of freedom were set to account for the weighting of the observations used in the MM slope estimate,

$$v_j = \sum_{i \in j} \frac{\partial \xi(\frac{\epsilon_i}{\sigma})}{\partial \left(\frac{\epsilon_i}{\sigma}\right)} \frac{1}{\left(\frac{\epsilon_i}{\sigma}\right)} - 2. \text{ This } v_j \text{ produced}$$

James-MM results that were very similar to setting  $v_j = N_j - 2$ .



## STANDARD, ALTERNATIVE, AND ROBUST SLOPE TEST STRATEGIES

### Y's Distribution

Eight shapes were used for Y, including a normal shape (skew=0, kurtosis=0), and seven other shapes with various degrees of skews and kurtosis (Table 1).

### Variance heterogeneity

The two considered residual variance ratios for the groups were 1/1 and 3/1. For conditions of unequal sample size, the residual variances were directly and inversely paired with the treatment group sample sizes.

### Sample sizes

Twenty and forty subjects per treatment group were used. The conditions of unequal sample size used twenty subjects in one group and forty in the other.

### Data generation method: Robustness

The following data generation method was used to create X and Y variables of desired distributions and variances with equal slopes in the two groups.

- 1) N values of one standard normal variate, Z, were generated, where N is the total sample size in two groups.
- 2) Y was created as a transformation of Z using Fleishman's (1978) method for generating nonnormal variables:

$$Y = a + bZ + cZ^2 + dZ^3 \quad (24)$$

The constants (a, b, c, and d) and resulting distributions are listed in Table 1.

- 3) An error variable for X ( $\varepsilon$ ) was generated as a standard normal variate. X's degree of nonnormality was a compromise between Y's nonnormality and  $\varepsilon$ 's normality.
- 4) Desired numbers of Ys and  $\varepsilon$ s were randomly assigned to treatment groups 1 and 2.
- 5) X was created as a function of Y and  $\varepsilon$ :

$$X_{ij} = \rho_j Y_{ij} + \sqrt{(1 - \rho_j^2)} \varepsilon_{ij}, \quad (25)$$

where  $\rho_j$  is the desired XY correlation for treatment group j.

- 5)  $Y_{ij}$  was multiplied by a number,  $\sigma_{Yj}$ , that resulted in a desired standard

deviation for Y in the jth treatment group and, in conjunction with  $\rho_j$ , a desired residual variance. The values of  $\sigma_{Yj}$  and  $\rho_j$  for the two groups achieved a particular residual variance ratio (Table 2), while keeping the slopes equal in the two groups.

### Data generation method: Power

The data generation process used to assess strategies' power was similar to the data generation process used to assess robustness. All variables' distributions were normal. One group's XY correlation and Y standard deviation were 0.5 and 1.0, respectively, while the second group's XY correlation and Y standard deviation were 0.0 and 0.866, respectively. The XY correlations and Y standard deviations across the groups resulted in a population slope difference of 0.5 while meeting the normality and equal residual variances assumptions of the standard test.

### Analysis strategy

The assessment of strategies' robustness involved comparing their average rejection rates to the nominal 0.05 rate for conditions where no slope differences existed in the population. Deviations from the nominal 0.05 rate were determined to be excessively conservative or liberal when they were outside of two standard errors band reflective of the number of replications used in this study

$$(0.05 + /- 2 \sqrt{\frac{(0.05)(0.95)}{10,000}} = 0.046 \text{ to } 0.054).$$

The standard error band roughly corresponded to Bradley's (1978) conservative range for robust Type I error rates, 0.045 to 0.055.

The assessment of strategies' power involved comparing strategies' average rejection rates to each other for conditions where actual slope differences existed in the population.

Follow-up analyses were also conducted to gain further insight into how the slope estimation strategies were working in the conditions of this study. These follow-up analyses included assessments of averages and standard deviations of the strategies' slope estimates to indicate their bias and efficiency,

and assessments of strategies' average standard errors to provide understanding of the accuracy of strategies' significance tests.

### Results

Tables 3-9 present the considered strategies' empirical Type I error rates across the 56 combinations of nonnormality, residual variance heterogeneity and sample size. Nonnormality affected the Standard, James and MM tests similarly, creating liberal Type I error rates when the Y distributions were leptokurtic and conservative Type I error rates when the distributions were platykurtic. The TW test had Type I error rates that were close to the nominal rate across the conditions of nonnormality. The TS test had Type I error rates that were consistently conservative across the considered levels of nonnormality. In terms of the hybrid strategies, James-TT had Type I error rates that were almost indistinguishable from James, while the James-MM, James-TW and James-TS strategies had Type I error rates reflective of the nonnormality strategy used, being excessively liberal for James-MM, being near 0.05 for James-TW, and being excessively conservative for James-TS.

The effect of residual variance heterogeneity on Type I error differed for the equal and unequal sample size conditions. When sample sizes were equal (Tables 4 & 9), MM was the only strategy affected by residual variance heterogeneity, becoming excessively liberal. When sample sizes were unequal (Tables 6 & 7), the groups' sample size-residual variance pairing affected the Standard, MM and TW tests similarly, making them liberal with an inverse pairing and conservative with a direct pairing. The James hybrid strategies were largely unaffected by the combination of unequal sample sizes and residual variances. James-TS produced conservative Type I error rates for most of the considered residual variance conditions.

The effect of combining nonnormality and residual variance heterogeneity (Tables 4, 6, 7, & 9) produced somewhat unique Type I error patterns for the nine tests. For the Standard test, residual variance heterogeneity usually made the effect of nonnormality less extreme except for when sample sizes were inversely-paired with

residual variances, in which case Type I error was made more extreme. For James and James-TT, residual variance heterogeneity made the effects of nonnormality less extreme, though James did not react as much to the combination of unequal sample sizes and residual variances as the Standard test. The MM test often had the most problematic Type I error rates for combinations of nonnormality and residual variance heterogeneity. The TW and TS tests were not particularly affected by the combination of nonnormality and residual variance heterogeneity, where the TW strategy was mainly impacted by the combination of unequal sample sizes and residual variances while the TS strategy was largely uninfluenced by anything. The Type I errors of hybrid strategies were reflective of the nonnormality strategy on which they were based, being liberal for James-MM, conservative for James-TS, and close to the 0.05 level for James-TW.

### Power

Table 10 compares the power of the nine strategies across three considered sample size conditions with normal distributions, equal residual variances and a population slope difference of 0.5. The most powerful strategies were the Standard, James and James-TT strategies, of which there was no overwhelming winner. The MM test had lower power rates than the Standard, James and James-TT tests. The James-MM hybrid strategy had less power than the MM strategy. The TW and James-TW tests had the lowest power rates of the considered strategies. The James-TS and TS strategies had higher power rates than the TW and James-TW strategies and (mostly) lower power rates than the MM and James-MM strategies. The use of TS as a hybrid with James (James-TS) increased its power relative to the TS strategy.

### Slope Estimation

To gain further insight into the four slope estimation methods (Standard, MM, TW and TS), Table 11 summarizes each methods' 10,000 estimates of one slope with population value 0.5 in samples of size 20. When distributions were normal, all four methods gave average slope values close to 0.5. The methods' standard deviations show that the Standard

## STANDARD, ALTERNATIVE, AND ROBUST SLOPE TEST STRATEGIES

method's estimates were least variable, followed by the MM estimates, the TS estimates and finally the TW estimates (corresponding to TW's relatively low power). The methods' average estimated standard errors correspond to the overall liberalness/conservativeness of the methods' significance tests, and for normal distributions show that on average all methods except for TS have standard errors that closely approximate slope variability. TS's bootstrapped standard errors over-estimated TS slope variability, corresponding to the conservativeness of its Type I error rates.

The slope estimation results in Table 11 for a leptokurtic Y (kurtosis = 12) differ from those for a normal Y (kurtosis = 0). For a leptokurtic Y, all estimation methods underestimate the population slope value of 0.5, where the least biased estimator is the Standard method while the most biased is the MM estimate. The Standard method's slope estimates are the most variable while the TS estimates are the least variable. The average standard errors of the Standard test and MM underestimate slope variability, corresponding to the liberalness of the Standard's and MM's Type I error rates. The TW estimates have standard errors that slightly underestimate slope variability. The TS estimate has standard errors that overestimate slope variability, corresponding to the conservativeness of TS. The results in Table 11 support previous findings that the TS estimator is more stable than the MM and Standard estimates when distributions are nonnormal (Wilcox & Keselman, 2004). These results extend previous work by showing that with nonnormality, the Standard test and MM regression underestimate slope variability (making the Type I error rates of the Standard and MM slope tests liberal), the Winsorized standard errors provide relatively accurate estimates of the variability of the trimmed slopes, while the TS bootstrap method overestimates slope variability (making the Type I error rates of the TS slope test conservative).

### Conclusion

The purpose of this study was to compare some recently-researched strategies for testing independent groups' regression slopes. The standard test of slope differences was shown to

have its usual robustness problems with respect to nonnormality and the pairing of unequal sample sizes and residual variances. Alternative strategies proposed for addressing nonnormality and used in hybrid strategies for addressing both nonnormality and residual variance heterogeneity were also assessed. The most promising of the alternative strategies in terms of robustness and power were the Theil-Sen strategy and a hybrid of Theil-Sen and the James second-order parametric alternative test. These Theil-Sen strategies had somewhat conservative Type I error rates that were largely unaffected by nonnormality and residual variance heterogeneity, and slope estimates that were efficient even for nonnormal data. The hybrid strategy of trimming and Winsorizing slope estimates and using them with the James test had Type I error rates that were closest of all the considered strategies to the nominal 0.05 level, but trimming and Winsorizing also produced slope tests with the lowest power rates of the considered strategies. Of the other strategies considered, James, James with a test statistic transformation for skewness, MM regression and the use of MM estimates with James are not recommended due to their robustness problems with nonnormal data.

In evaluating the results of this and other studies, it is important to acknowledge that the effects of nonnormality have been considered in very different ways, all of which have implications for studies' results. When the nonnormality of  $\epsilon$  is directly manipulated, the standard and James tests have appeared to be robust to all but the most extreme shapes (e.g., skew=6.2, kurtosis=114 in Luh & Guo, 2000, 2002).

When the nonnormality of Y and/or X is manipulated, the standard and James tests become problematic for relatively small degrees of nonnormality (e.g., skew=1.95, kurtosis=7.69 in Deshon & Alexander, 1996). When nonnormality has been studied in terms of outliers in multivariate distributions, the standard test is problematic and MM regression performs well (Anderson & Schumacker, 2003). The second type of nonnormality, in Y and X, creates great problems for methods that use least squares estimation methods due to the higher likelihood of leverage points.

MOSES & KLOCKARS

Table 1. Shapes and Fleishman constants used to generate the variables

Skew	Kurtosis	a	b	c (= -a)	d
0	-1.15	0	1.34	0	-0.132
0	0	0	1	0	0
1.2	1.11	-0.340774	1.095718	0.340774	-0.080735

Table 2. Correlations and standard deviations used to create levels of residual variance heterogeneity.

Residual Variance Ratio	$\rho_1$	$\sigma_{Y1}$	$\rho_2$	$\sigma_{Y2}$
1/1	0.5	1.0	0.5	1.0
1/3	0.5	1.0	0.3162	1.5811

Table 3. Empirical Type I error rates for group sample sizes of 20, 20 and a residual variance ratio of 1/1.

Skew	Kurtosis	Standard	James	MM	TW	TS	Hybrid Strategies			
							James-TT	James-MM	James-TW	James-TS
0	-1.15	0.0256*	0.0257*	0.0260*	0.0479	0.0207*	0.0258*	0.0214*	0.0456	0.0204*
0	0	0.0486	0.0481	0.0610*	0.0497	0.0282*	0.0483	0.0486	0.0479	0.0262*
1.2	1.11	0.0668*	0.0676*	0.1099*	0.0508	0.0304*	0.0667*	0.0985*	0.0490	0.0358*
1.6	2.86	0.0941*	0.0961*	0.1383*	0.046*	0.0302*	0.0949*	0.1341*	0.0455*	0.0345*
0	3	0.0912*	0.0936*	0.0999*	0.0532	0.0307*	0.0935*	0.0874*	0.0508	0.0317*
0	6	0.1178*	0.1200*	0.1226*	0.0500	0.0341*	0.1198*	0.1091*	0.0482	0.0328*
0	9	0.1359*	0.1403*	0.1257*	0.0458	0.0308*	0.1395*	0.1193*	0.0429*	0.0288*
0	12	0.1645*	0.1727*	0.1347*	0.0542	0.0303*	0.1714*	0.1343*	0.0510	0.0281*

\* Type I error rates outside +/- 2 standard errors of the nominal 0.0500 rate (0.0456 to 0.0544).

Table 4. Empirical Type I error rates for group sample sizes of 20, 20 and a residual variance ratio of 3/1.

Skew	Kurtosis	Standard	James	MM	TW	TS	Hybrid Strategies			
							James-TT	James-MM	James-TW	James-TS
0	-1.15	0.0428*	0.0398*	0.0820*	0.0566*	0.0291*	0.0393*	0.0346*	0.0531	0.0281*
0	0	0.0514	0.0481	0.1020*	0.0511	0.0323*	0.0482	0.0438*	0.0486	0.0305*
1.2	1.11	0.0630*	0.0598*	0.1339*	0.0550*	0.0304*	0.0591*	0.0809*	0.0509	0.0292*
1.6	2.86	0.0792*	0.0774*	0.1525*	0.0517	0.0317*	0.0757*	0.1063*	0.0477	0.0278*
0	3	0.0689*	0.0682*	0.1141*	0.0523	0.0303*	0.0674*	0.0653*	0.0480	0.0302*
0	6	0.0931*	0.0946*	0.1286*	0.0517	0.0304*	0.0938*	0.0824*	0.0470	0.0251*
0	9	0.1106*	0.1096*	0.1368*	0.0481	0.0326*	0.1091*	0.0886*	0.0440*	0.0263*
0	12	0.1176*	0.1206*	0.1367*	0.0479	0.0323*	0.1202*	0.0904*	0.0437*	0.0235*

\* Type I error rates outside +/- 2 standard errors of the nominal 0.0500 rate (0.0456 to 0.0544).

STANDARD, ALTERNATIVE, AND ROBUST SLOPE TEST STRATEGIES

Table 5. Empirical Type I error rates for group sample sizes of 20, 40 and a residual variance ratio of 1/1.

Skew	Kurtosis	Standard	James	MM	TW	TS	Hybrid Strategies			
							James-TT	James-MM	James-TW	James-TS
0	-1.15	0.0300*	0.0306*	0.0247*	0.0546*	0.0263*	0.0304*	0.0252*	0.0532	0.0259*
0	0	0.0496	0.0486	0.0581	0.0507	0.0314*	0.0482	0.0506	0.0468	0.0320*
1.2	1.11	0.0701*	0.0720*	0.1174*	0.0571*	0.0327*	0.0713*	0.1090*	0.0504	0.0407*
1.6	2.86	0.1040*	0.1054*	0.1451*	0.0556*	0.0321*	0.1049*	0.1530*	0.0526	0.0412*
0	3	0.0931*	0.0929*	0.0981*	0.0531	0.0327*	0.0927*	0.0870*	0.0482	0.0349*
0	6	0.1235*	0.1286*	0.1143*	0.0525	0.0325*	0.1269*	0.1080*	0.0482	0.0341*
0	9	0.1524*	0.1599*	0.1226*	0.0522	0.0371*	0.1593*	0.1225*	0.0485	0.0356*
0	12	0.1677*	0.1804*	0.1253*	0.0528	0.0349*	0.1799*	0.1293*	0.0508	0.0338*

\* Type I error rates outside +/- 2 standard errors of the nominal 0.0500 rate (0.0456 to 0.0544).

Table 6. Empirical Type I error rates for group sample sizes of 20, 40 and a residual variance ratio of 1/3 (Direct Pairing).

Skew	Kurtosis	Standard	James	MM	TW	TS	Hybrid Strategies			
							James-TT	James-MM	James-TW	James-TS
0	-1.15	0.0134*	0.0353*	0.0251*	0.0273*	0.0312*	0.0352*	0.0299*	0.0492	0.0295*
0	0	0.0218*	0.0534	0.0406*	0.0258*	0.0313*	0.0535	0.0508	0.0505	0.0335*
1.2	1.11	0.0324*	0.0660*	0.0851*	0.0274*	0.0279*	0.0656*	0.0913*	0.0499	0.0358*
1.6	2.86	0.0443*	0.0905*	0.1171*	0.0269*	0.0345*	0.0893*	0.1203*	0.0488	0.0383*
0	3	0.0379*	0.0789*	0.0654*	0.0244*	0.0349*	0.0792*	0.0746*	0.0488	0.0352*
0	6	0.0612*	0.1087*	0.0883*	0.0283*	0.0348*	0.1088*	0.0953*	0.0493	0.0363*
0	9	0.0765*	0.1327*	0.0997*	0.0290*	0.0347*	0.1319*	0.1011*	0.0466	0.0344*
0	12	0.0900*	0.1516*	0.1172*	0.0294*	0.0342*	0.1503*	0.1151*	0.0486	0.0286*

\* Type I error rates outside +/- 2 standard errors of the nominal 0.0500 rate (0.0456 to 0.0544).

Table 7. Empirical Type I error rates for group sample sizes of 40, 20 and a residual variance ratio of 1/3 (Inverse Pairing).

Skew	Kurtosis	Standard	James	MM	TW	TS	Hybrid Strategies			
							James-TT	James-MM	James-TW	James-TS
0	-1.15	0.0827*	0.0360*	0.1722*	0.1028*	0.0305*	0.0356*	0.0342*	0.0497	0.0337*
0	0	0.0986*	0.0504	0.1860*	0.0989*	0.0341*	0.0504	0.0497	0.0477	0.0320*
1.2	1.11	0.1165*	0.0624*	0.2039*	0.0981*	0.0347*	0.0615*	0.0860*	0.0509	0.0342*
1.6	2.86	0.1392*	0.0798*	0.2017*	0.0954*	0.0386*	0.0769*	0.0998*	0.0503	0.0312*
0	3	0.1359*	0.0760*	0.1880*	0.1007*	0.0370*	0.0760*	0.0682*	0.0510	0.0343*
0	6	0.1620*	0.0997*	0.1902*	0.1032*	0.0354*	0.0995*	0.0797*	0.0531	0.0311*
0	9	0.1812*	0.1164*	0.1879*	0.0936*	0.0390*	0.1148*	0.0863*	0.0467	0.0301*
0	12	0.1862*	0.1263*	0.1810*	0.0887*	0.0391*	0.1260*	0.0903*	0.0471	0.0291*

\* Type I error rates outside +/- 2 standard errors of the nominal 0.0500 rate (0.0456 to 0.0544).

MOSES & KLOCKARS

Table 8. Empirical Type I error rates for group sample sizes of 40, 40 and a residual variance ratio of 1/1.

Skew	Kurtosis	Standard	James	MM	TW	TS	Hybrid Strategies			
							James-TT	James-MM	James-TW	James-TS
0	-1.15	0.0280*	0.0284*	0.0206*	0.0526	0.0289*	0.0283*	0.0200*	0.0526	0.0300*
0	0	0.0479	0.0477	0.0559*	0.0504	0.0321*	0.0479	0.0498	0.0501	0.0353*
1.2	1.11	0.0672*	0.0677*	0.1115*	0.0484	0.0388*	0.0678*	0.1101*	0.0478	0.0521
1.6	2.86	0.1028*	0.1042*	0.1435*	0.0495	0.0371*	0.1032*	0.1635*	0.0494	0.0483
0	3	0.1006*	0.1028*	0.0900*	0.0518	0.0391*	0.1030*	0.0878*	0.0518	0.0428*
0	6	0.1396*	0.1436*	0.1044*	0.0489	0.0352*	0.1432*	0.1057*	0.0483	0.0381*
0	9	0.1709*	0.1752*	0.1151*	0.0467	0.0389*	0.1743*	0.1235*	0.0458	0.0398*
0	12	0.1952*	0.2004*	0.1200*	0.0483	0.0396*	0.1990*	0.1347*	0.0476	0.0387*

\* Type I error rates outside +/- 2 standard errors of the nominal 0.0500 rate (0.0456 to 0.0544).

Table 9. Empirical Type I error rates for group sample sizes of 40, 40 and a residual variance ratio of 3/1.

Skew	Kurtosis	Standard	James	MM	TW	TS	Hybrid Strategies			
							James-TT	James-MM	James-TW	James-TS
0	-1.15	0.0360*	0.0354*	0.0809*	0.0506	0.0352*	0.0353*	0.0310*	0.0494	0.0388*
0	0	0.0494	0.0479	0.0884*	0.0506	0.0386*	0.0482	0.0470	0.0490	0.0423*
1.2	1.11	0.0662*	0.0636*	0.1511*	0.0499	0.0432*	0.0639*	0.0958*	0.0484	0.0503
1.6	2.86	0.0851*	0.0849*	0.1828*	0.0515	0.0390*	0.0841*	0.1236*	0.0506	0.0434*
0	3	0.0844*	0.0821*	0.1191*	0.0548	0.0386*	0.0820*	0.0733*	0.0520	0.0400*
0	6	0.1025*	0.1011*	0.1279*	0.0528	0.0386*	0.1008*	0.0820*	0.0513	0.0378*
0	9	0.1272*	0.1290*	0.1427*	0.0512	0.0408*	0.1278*	0.0945*	0.0482	0.0386*
0	12	0.1426*	0.1460*	0.1431*	0.0486	0.0367*	0.1439*	0.0998*	0.0474	0.0322*

\* Type I error rates outside +/- 2 standard errors of the nominal 0.0500 rate (0.0456 to 0.0544).

Table 10. Empirical Power rates for population slope differences of .5 and normality and residual variance assumptions met.

Sample Sizes		Standard	James	MM	TW	TS	Hybrid Strategies			
Group 1	Group 2						James-TT	James-MM	James-TW	James-TS
20	20	0.3910	0.3901	0.3735	0.2270	0.2655	0.3906	0.3372	0.2215	0.2891
20	40	0.5096	0.4989	0.4730	0.3082	0.3931	0.4994	0.4403	0.2930	0.4202
40	40	0.6909	0.6912	0.6359	0.4369	0.6179	0.6912	0.6180	0.4353	0.6483

## STANDARD, ALTERNATIVE, AND ROBUST SLOPE TEST STRATEGIES

Table 11. Descriptive analyses for the slope estimation strategies (population slope = 0.5).

Skew	Kurtosis	Sample Size		Standard	MM	TW	TS
0	0	20	Mean of Slope Estimates	0.5043	0.5043	0.5079	0.4998
			Standard Deviation of Slope Estimates	0.2080	0.2214	0.2819	0.2275
			Mean of Slope Standard Error Estimates	0.2046	0.2205	0.2721	0.2752
0	12	20	Mean of Slope Estimates	0.4598	0.2987	0.3538	0.3280
			Standard Deviation of Slope Estimates	0.2866	0.2361	0.2528	0.2159
			Mean of Slope Standard Error Estimates	0.1907	0.1701	0.2383	0.2575




Figure 1




Figure 2




Figure 3




Figure 4

This type of nonnormality was of most interest in this study. It caused serious problems for the standard test that warranted the use of alternative and robust strategies, and it created data situations that differentiated all of the studied strategies.

To gain some final insight into the for considered slope estimation methods, a representative sample of twenty observations was generated from this study's kurtosis=12 condition. Figure 1 shows these XY data and plots the Standard, least squares regression line. There is one very extreme X observation (almost 3 standard deviations from X's population mean of zero) that is also very low on Y (i.e., a bad leverage point). This observation causes the standard slope estimation method to underestimate the population slope of 0.5 in its slope estimate of 0.421. Figures 2 and 3 plot the observations in the data that are not excluded in computing the trimmed slope (Figure 2) and the Theil-Sen slope (Figure 3). The trimmed and Theil-Sen methods underestimate the population slope more than the Standard method, producing slope estimates of 0.393 and 0.231, respectively.

Figure 4 is especially useful for understanding the very complicated MM regression procedure. All twenty of the original observations are used in MM regression, but contribute in weighted form to the final MM slope estimate. The observations' weights in Figure 4 show that the high-leverage observation is weighed very heavily by the MM method, causing the MM slope estimate to be relatively small (0.148). The observations that are far from the MM regression line are assigned small weights. Figure 4 shows that with MM high-leverage points can be weighted such that they influence the final slope estimate much more than the Standard least squares estimate. The large weights that are assigned to high leverage points in MM result in MM standard errors that underestimate slope variability (the  $\mathbf{W}$  in equation 18 is large) and inflate the Type I error of the MM strategy. Figure 4 makes it clear that the problems of the MM strategy with respect to high leverage points are not likely to be fixed by altering the weighting function,  $\xi$ , or the  $\kappa$  that determines how each of the scaled residuals are weighted. It may be possible to address MM's

problems with high leverage data points through a wise choice of starting values that define the MM regression line and the residuals with respect to this line.

#### Implications

This article considered some of the recently-researched slope test strategies. Some of the strategies not considered in this paper were excluded because they have had noted problems and criticisms, including nonparametric alternative tests (Marascuilo, 1966; Dretzke, Levin & Serlin, 1982; Deshon & Alexander, 1996), residuals-based bootstrapping (Luh & Guo, 2000), ranked data (Headrick & Sawilowsky, 2000; Klockars & Moses, 2002), data transformations (Wilcox & Keselman, 2004; Aguinis & Pierce, 1998; Keselman, Carriere & Lix, 1995; Glass, Peckham & Saunders, 1972), several robust regression strategies (Anderson & Schumaker, 2003) and judgment-based elimination of outliers (Wilcox, 1996; He & Portnoy, 1992).

There are other strategies that are variations on the ones considered in this study, such as the use of Theil-Sen after trimming outliers (Wilcox & Keselman, 2004), the use of Theil-Sen based on less than the  $N(N-1)/2$  slopes that could be created out of all pairs of observations (Ebrahim & Al-Nasser, 2005), other parametric alternative tests for residual variance heterogeneity (Alexander & Deshon, 1994; Welch, 1938), and trimmed and Winsorized estimates with varied amounts of trimming.

The results of this study suggest that an especially promising slope test strategy would combine the best features of the trimming and Winsorizing methods with Theil-Sen. By using the trimming and Winsorizing strategy on the  $N(N-1)/2$  slopes that could be created out of all pairs of observations rather than only  $N/2$  pairs, the final trimmed slope estimates should have stability levels that are similar to those of Theil-Sen, ultimately improving the power of the trimmed and Winsorized slope test. This proposed test would avoid the excessively time consuming and excessively-conservative bootstrapping that accompanies the Theil-Sen method, reduce the bias of the Theil-Sen estimates for nonnormal data, provide a



## STANDARD, ALTERNATIVE, AND ROBUST SLOPE TEST STRATEGIES

reasonable answer to the awkward definition of the number of observations used by the median-based Theil-Sen, and provide the analyst some flexibility in terms of the extent of trimming used in the final slope estimates. A study that considers how the number of slopes (Ebrahim & Al-Nasser, 2005) and the extent of trimming contribute to Type I error and power across conditions of nonnormality would be especially useful for creating the best version of this proposed test of slope differences.

### References

- Aguinis, H., & Pierce, C. A. (1998). Heterogeneity of error variance and the assessment of moderating effects of categorical variables: A conceptual review. *Organizational Research Methods, 1*, 296-314.
- Aguinis, H., Petersen, S. A., & Pierce, C. A. (1999). Appraisal of the homogeneity of error variance assumption and alternatives to multiple regression for estimating moderating effects of categorical variables. *Organizational Research Methods, 2*(4), 315-339.
- Aiken, L. S. & West, S. G. (1991). *Multiple regression: Testing and interpreting interactions*. Newbury Park, CA: Sage.
- Alexander, R. A., & DeShon, R. P. (1994). Effect of error variance heterogeneity on the power of tests for regression slope differences. *Psychological Bulletin, 115*(2), 308-314.
- Anderson, C., & Schumacker, R. E. (2003). A comparison of five robust regression methods with ordinary least squares regression: Relative efficiency, bias, and test of the null hypothesis. *Understanding Statistics, 2*(2), 79-103.
- Bradley, J. C. (1978). Robustness? *British Journal of Mathematical and Statistical Psychology, 31*, 144-152.
- Conerly, M. D. & Mansfield, E. R. (1988). An approximate test for comparing heteroscedastic regression models. *Journal of the American Statistical Association, 83*(403), 811-817.
- Conover, W. J. & Iman, R. L. (1982). Analysis of covariance using the rank transformation. *Biometrics, 38*, 715-724.
- Cronbach, L. J. & Snow, R. E. (1977). *Aptitude and instructional methods*. Irvington, NY.
- Dance, K. A., & Neufeld, R. W. J. (1988). Aptitude-treatment interaction research in the clinical setting: A review of attempts to dispel the "Patient Uniformity" myth. *Psychological Bulletin, 104*(2), 192-213.
- DeShon, R. P., & Alexander, R. A. (1994). A generalization of James' second-order approximation to the test for regression slope equality. *Educational and Psychological Measurement, 54*, 328-335.
- DeShon, R. P. & Alexander, R. A. (1996). Alternative procedures for testing regression slope homogeneity when group error variances are unequal. *Psychological Methods, 1*(3), 261-277.
- Dretzke, B. J. & Levin, J. R.; Serlin, R. C. (1982). Testing for regression homogeneity under variance heterogeneity. *Psychological Bulletin, 91*, 376-383.
- Ebrahim, M. A., & Al-Nasser, A. D. (2005). Estimating the slope of simple linear regression in the presence of outliers. *Journal of Modern Applied Statistical Methods, 4*(2), 509-513.
- Fleishman, A. I. (1978). A method for simulating non-normal distributions. *Psychometrika, 43*(4), 521-532.
- Glass, G. V., Peckham, P. D., & Sanders, J. R. (1972). Consequences of failure to meet assumptions underlying the fixed effects analyses of variance and covariance. *Review of Educational Research, 42*, 237-288.
- Guo, J. H. (1996). Distribution-free test for slope in linear regression model with replications. *Communications in Statistics-Theory and Methods, 25*, 2069-2078.
- Hall, P. (1992). On the removal of skewness by transformation. *Journal of Royal Statistical Society, B54*, 221-228.
- He, X., & Portnoy, S. (1992). Reweighted LS estimators converge at the same rate as the initial estimator. *Annals of Statistics, 20*, 2161-2167.
- Headrick, T. C. & Sawilowsky, S. S. (2000). Properties of the rank transformation in factorial analysis of covariance. *Communications in Statistics- Simulation and Computation, 29*, 1059-1087.

- Holland, P. & Welsch, R. (1977). Robust regression using interactively reweighted least squares. *Communications in Statistics-Theor. Meth.*, 6, 813-827.
- Hunter, J. E., Schmidt, F. L., & Hunter, R. (1979). Differential validity of employment tests by race: A comprehensive review and analysis. *Psychological Bulletin*, 86(4), 721-735.
- James, G. S. (1951). The comparison of several groups of observations when the ratios of population variances is unknown. *Biometrika*, 38, 324-329.
- Johnson, N. J. (1978). Modified *t* tests and confidence intervals for asymmetrical populations. *Journal of the American Statistical Association*, 73, 536-544.
- Keselman, H. J., Carriere, K. C., & Lix, L. M. (1995). Robust and powerful nonorthogonal analyses. *Psychometrika*, 60(3), 395-418.
- Keselman, H. J., Wilcox, R. R., Othman, A. R. & Fradette, K. (2002). Trimming, transforming statistics, and bootstrapping: Circumventing the biasing effects of heteroscedasticity and nonnormality. *Journal of Modern Applied Statistical Methods*, 1(2), 288-309.
- Klockars, A. & Moses, T. (2002). Type I error rates for rank-based tests of homogeneity of slopes. *Journal of Modern Applied Statistical Methods*, 1(2), 452-460.
- Luh, W., & Guo, J. (2000). Approximate transformation trimmed mean methods to the test of simple linear regression slope equality. *Journal of Applied Statistics*, 27(7), 843-857.
- Luh, W., & Guo, J. (2002). Using Johnson's transformation with approximate test statistics for the simple regression slope homogeneity. *The Journal of Experimental Education*, 71(1), 69-81.
- Marascuilo, L. A. (1966). Large-sample multiple comparisons. *Psychological Bulletin*, 65, 280-290.
- Micceri, T. (1989). The unicorn, the normal curve, and other improbable creatures. *Psychological Bulletin*, 105, 156-166
- Moses, T., & Klockars, A. (2005). Testing for aptitude-treatment interactions in analysis of covariance and randomized block designs under assumption violations. *Journal of Modern Applied Statistical Methods*, 4(2), 460-472.
- Myers, J. L. & Well, A. D. (1995). *Research design and statistical analysis*. Hillsdale, NJ: Lawrence Erlbaum Associates.
- Overton, R. C. (2001). Moderated multiple regression for interactions involving categorical variables: A statistical control for heterogeneous variance across two groups. *Psychological Methods*, 6(3), 218-233.
- Pedhazur, E. J. (1997). *Multiple regression in behavioral research*. Fort Worth, Texas.
- Rousseeuw, P. J. (1984). Least median of squares regression. *Journal of the American Statistical Association*, 79, 871-880.
- Rousseeuw, P. J. & Van Driessen, K. (2000). An algorithm for positive-breakdown regression based on concentration steps. *Data Analysis: Scientific Modeling and Practical Application*, ed. W. Gaul, O. Optiz, & M. Schader. New York: Springer-Verlag, 335-346.
- SAS Institute Inc. (2003). *SAS/STAT user's guide, version 9.1: The ROBUSTREG Procedure*. Cary, NC: SAS Institute.
- Seligman, M. E. P. (2002). *Authentic happiness: Using the new positive psychology to realize your potential for lasting fulfillment*. Free Press, New York.
- Sen, P. K. (1968). Estimate of the regression coefficient based on Kendall's tau. *Journal of the American Statistical Association*, 63, 1379-1389.
- Theil, H. (1950). A rank-invariant method of linear and polynomial regression analysis. *Indagationes Mathematicae*, 12, 85-91.
- Wang, X. (2005). Asymptotics of the Theil-Sen estimator in the simple linear regression model with a random covariate. *Nonparametric Statistics*, 17(1), 107-120.
- Welch, B. L. (1938). The significance of the difference between two means when the population variances are unequal. *Biometrika*, 29, 350-362.
- Wilcox, R. R. (1996). A review of some recent developments in robust regression. *British Journal of Mathematical and Statistical Psychology*, 49, 253-274.
- Wilcox, R. R. (2004). Some results on extensions and modifications of the Theil-Sen regression estimator. *British Journal of Mathematical and Statistical Psychology*, 57, 265-280.
- Wilcox, R. R. (2005). *Introduction to robust estimation and hypothesis testing* (2<sup>nd</sup> ed.). San Diego, CA: Academic Press.
- Wilcox, R. R., & Keselman, H. J. (2004). Robust regression methods: Achieving small standard errors when there is heteroscedasticity. *Understanding Statistics*, 3(4), 349-364.
- Yohai, V. J. (1987). High breakdown point and high efficiency robust estimates for regression. *Annals of Statistics*, 15, 642-656.