

Addendum

Pubertal development mediates the association between family environment and brain structure and function in childhood – ADDENDUM

Sandra Thijssen, Paul F. Collins and Monica Luciana

doi: 10.1017/S0954579419000580. Published online by Cambridge University Press, 1 July 2019.

In November 2019, the Adolescent Brain and Cognitive Development consortium communicated that previously released functional MRI data from Philips scanners has been processed incorrectly and should not be analyzed. The resting-state fMRI analyses reported in Thijssen et al. (2019) include data from Philips scanners. We have reanalyzed our resting-state fMRI data excluding participants scanned on a Philips scanner ($n = 256$). Excluding the Philips data did not significantly affect our results. For the new results, please see below. The conclusions described in the manuscript remain unchanged.

Resting-state fMRI

In the total sample excluding those scanned with Phillips scanners, the total, direct, and indirect effects of Family Environment on cingulo-opercular network–left amygdala functional connectivity were $\beta = 0.068$, $p = .003$, $\beta = 0.059$, $p = .010$, $\beta = 0.009$, $p = .071$, respectively. For cingulo-opercular network–right amygdala functional connectivity, the total, direct, and indirect effects were $\beta = 0.044$, $p = .055$, $\beta = 0.036$, $p = .122$, $\beta = 0.008$, $p = .106$, respectively. Thus, Family Environment was positively

associated with cingulo-opercular network–amygdala functional connectivity. For the left amygdala–cingulo-opercular network functional connectivity, the indirect effect of family environment on functional connectivity via pubertal stage indicated a trend in the expected direction. For right amygdala–cingulo-opercular network functional connectivity, the indirect effect no longer indicates a trend ($p > .1$). As the effect size of the indirect effect increased from $\beta = 0.007$ to $\beta = 0.008$ when excluding the Philips data, this difference is solely explained by decreased power.

The exploratory analyses stratified by sex suggest that the total and direct effects of Family Environment on cingulo-opercular network–left amygdala functional connectivity were significant for girls, whereas a trend was found for the indirect effect ($\beta = 0.090$, $p = .005$, $\beta = 0.078$, $p = .017$, $\beta = 0.012$, $p = .093$, respectively). For boys, no significant effects were found ($\beta = 0.049$, $p = .112$, $\beta = 0.044$, $p = .157$, $\beta = 0.005$, $p = .459$, respectively). For cingulo-opercular network–right amygdala functional connectivity, no significant effects were found for girls nor boys (girls: $\beta = 0.061$, $p = .071$, $\beta = 0.053$, $p = .132$, $\beta = 0.008$, $p = .226$ for total, direct, and indirect effects, respectively; boys $\beta = 0.030$, $p = .322$, $\beta = 0.023$, $p = .459$, $\beta = 0.007$, $p = .289$, for total, direct, and indirect effects, respectively).

Cite this article: Thijssen S, Collins PF, Luciana M (2021). Pubertal development mediates the association between family environment and brain structure and function in childhood – ADDENDUM. *Development and Psychopathology* 33, 372–375. <https://doi.org/10.1017/S0954579420000322>

© Cambridge University Press 2020. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 6. Mediation model parameters--Cingulo-opercular network-amygdala connectivity

	CON-left amygdala				CON-right amygdala			
	β	S.E.	β /S.E.	p	β	S.E.	β /S.E.	p
Family Environment [∞]	0.059	0.023	2.582	.010	0.036	0.023	1.548	.122
Pubertal stage ⁺	-0.064	0.033	-1.923	.054	-0.055	0.033	-1.692	.091
Age	0.002	0.023	0.106	.915	0.021	0.024	0.878	.380
Sex	-0.031	0.027	-1.160	.246	-0.055	0.028	-1.984	.047
Race	-0.040	0.023	-1.768	.077	-0.032	0.021	-1.531	.126
	Outcome: Pubertal stage				Outcome: Pubertal stage			
Family Environment ^f	-0.136	0.022	-6.189	<.001	-0.136	0.022	-6.189	<.001
Age	0.231	0.021	11.159	<.001	0.231	0.021	11.159	<.001
Sex	-0.504	0.016	-30.950	<.001	-0.504	0.016	-30.950	<.001
Race	0.136	0.020	6.686	<.001	0.136	0.020	6.686	<.001

Note: CON = cingulo-opercular network; [∞] = direct effect; ⁺ = indirect effect Step 2; ^f = indirect effect Step 1.

Table S9. Mediation model parameters--Cingulo-opercular network-amygdala connectivity in girls

	CON-left amygdala				CON-right amygdala			
	β	S.E.	β /S.E.	p	β	S.E.	β /S.E.	p
Family Environment [∞]	0.078	0.033	2.391	.027	0.053	0.035	1.505	.132
Pubertal stage ⁺	-0.071	0.039	-1.816	.069	-0.049	0.039	-1.261	.207
Age	-0.048	0.034	-1.404	.160	0.014	0.031	0.416	.677
Race	-0.044	0.036	-1.209	.227	-0.050	0.033	-1.637	.102
	Outcome: Pubertal stage				Outcome: Pubertal stage			
Family Environment ^f	-0.171	0.034	-5.058	<.001	-0.171	0.034	-5.058	<.001
Age	0.300	0.030	9.991	.001	0.300	0.030	9.991	.001
Race	0.111	0.034	3.295	<.001	0.111	0.034	3.295	<.001

Note: [∞] = direct effect; ⁺ = indirect effect Step 2; ^f = indirect effect Step 1; CON = cingulo-opercular network.

Table S12. Mediation model parameters--Cingulo-opercular network-amygdala connectivity in boys

	CON-left amygdala				CON-right amygdala			
	β	S.E.	β /S.E.	p	β	S.E.	β /S.E.	p
Family Environment [∞]	0.044	0.031	1.415	.157	0.023	0.031	0.740	.459
Pubertal stage ⁺	-0.034	0.043	-0.794	.427	-0.048	0.042	-1.146	.252
Age	0.042	0.032	1.298	.194	0.027	0.030	0.900	.368
Race	-0.038	0.029	-1.286	.198	-0.016	0.029	-0.554	.579
	Outcome: Pubertal stage				Outcome: Pubertal stage			
Family Environment ^f	-0.142	0.036	-3.960	<.001	-0.142	0.036	-3.960	<.001
Age	0.223	0.035	6.289	<.001	0.223	0.035	6.289	<.001
Race	0.212	0.031	6.846	<.001	0.212	0.031	6.846	<.001

Note: [∞] = direct effect; ⁺ = indirect effect Step 2; ^f = indirect effect Step 1; CON = cingulo-opercular network.

Somato-motor mouth network-amygdala functional connectivity

For the resting-state model with motor processing measures, only the total and direct effects of Family Environment on

SOMM-left amygdala FC were significant ($\beta = 0.060$, $p = .005$, $\beta = 0.061$, $p = .006$, respectively), but not the indirect effect ($\beta = -0.001$, $p = .847$). No associations between Family Environment and SOMM-right amygdala were found ($\beta = -0.013$,

$p = .583$, $\beta = -0.015$, $p = .542$, $\beta = 0.002$, $p = .693$, for total, direct, and indirect effects, respectively).

For the resting-state model with motor processing measures, in girls the total and direct effects, and in boys only the total effect of Family Environment on SOMM-left amygdala FC were significant (girls: $\beta = 0.056$, $p = .048$, $\beta = 0.064$, $p = .036$, $\beta = -0.008$, $p = .302$ for total, direct, and indirect effects, respectively; boys

$\beta = 0.063$, $p = .041$, $\beta = 0.056$, $p = .074$, $\beta = 0.006$, $p = .357$. for total, direct, and indirect effects, respectively). No significant associations were found between Family Environment and SOMM-right amygdala (girls: $\beta = 0.027$, $p = .421$, $\beta = 0.026$, $p = .452$, $\beta = 0.001$, $p = .886$ for total, direct, and indirect effects, respectively; boys $\beta = -0.043$, $p = .157$, $\beta = -0.046$, $p = .141$, $\beta = 0.003$, $p = .606$ for total, direct, and indirect effects, respectively).

TableS6. Mediation model parameters--Somatomotor-mouth network-amygdala connectivity

	SOMM-left amygdala				SOMM-right amygdala			
	β	S.E.	$\beta/S.E.$	p	β	S.E.	$\beta/S.E.$	p
Family Environment [∞]	0.061	0.022	2.767	.006	-0.015	0.024	-0.610	.542
Pubertal stage ⁺	0.007	0.035	0.199	.843	-0.013	0.033	-0.404	.686
Age	-0.021	0.023	-0.925	.355	-0.011	0.025	-0.437	.662
Sex	-0.009	0.029	-0.309	.757	0.007	0.026	0.279	.780
Race	-0.015	0.022	-0.678	.498	-0.026	0.022	-1.182	.237
	Outcome: Pubertal stage				Outcome: Pubertal stage			
Family Environment ^f	-0.136	0.022	-6.189	<.001	-0.136	0.022	-6.189	<.001
Age	0.231	0.021	11.159	<.001	0.231	0.021	11.159	<.001
Sex	-0.504	0.016	-30.950	<.001	-0.504	0.016	-30.950	<.001
Race	0.136	0.020	6.686	<.001	0.136	0.020	6.686	<.001

Note: [∞] = direct effect; ⁺ = indirect effect Step 2; ^f = indirect effect Step 1; SOMM = somato-motor mouth network.

TableS17. Mediation model parameters--Somatomotor-mouth network-amygdala connectivity in girls

	SOMM-left amygdala				SOMM-right amygdala			
	β	S.E.	$\beta/S.E.$	p	β	S.E.	$\beta/S.E.$	p
Family Environment [∞]	0.064	0.031	2.099	.036	0.026	0.035	0.0753	.452
Pubertal stage ⁺	0.045	0.041	1.094	.274	-0.005	0.037	-0.146	.884
Age	-0.004	0.031	-0.113	.910	0.010	0.035	0.281	.779
Race	-0.071	0.032	-2.208	.027	-0.052	0.032	-1.592	.111
	Outcome: Pubertal stage				Outcome: Pubertal stage			
Family Environment ^f	-0.171	0.034	-5.058	<.001	-0.171	0.034	-5.058	<.001
Age	0.300	0.030	9.991	.000	0.300	0.030	9.991	.000
Race	0.111	0.034	3.295	<.001	0.111	0.034	3.295	<.001

Note: [∞] = direct effect; ⁺ = indirect effect Step 2; ^f = indirect effect Step 1; SOMM = somato-motor mouth network.

TableS20. Mediation model parameters—Somatomotor-mouth network—amygdala connectivity in boys

	SOMM-left amygdala				SOMM-right amygdala			
	β	S.E.	β /S.E.	<i>p</i>	β	S.E.	β /S.E.	<i>p</i>
Family Environment [∞]	0.056	0.032	1.787	.074	-0.046	0.031	-1.474	.141
Pubertal stage [†]	-0.043	0.045	-0.972	.331	-0.023	0.044	-0.528	.597
Age	-0.031	0.031	-1.003	.316	-0.024	0.034	-0.698	.485
Race	0.034	0.029	1.160	.246	-0.002	0.031	-0.060	.952
	Outcome: Pubertal stage				Outcome: Pubertal stage			
Family Environment ^f	-0.142	0.036	-3.960	<.001	-0.142	0.036	-3.960	<.001
Age	0.223	0.035	6.289	<.001	0.223	0.035	6.289	<.001
Race	0.212	0.031	6.846	<.001	0.212	0.031	6.846	<.001

Note: [∞] = direct effect; [†] = indirect effect Step 2; ^f = indirect effect Step 1. SOMM = somato-motor mouth network.

Table3. Correlation between MRI measures

	ACC CA	ACC FA	Amygdala SV	CON-l amygdala FC	CON-r amygdala FC
ACC CT	-.082	-.072	.067	.003	.036
ACC CA		-.257	-.017	.033	.013
ACC FA			-.049	-.012	-.038
Amygdala SV				-.058	-.033
CON-- amygdala FC					.586

Note: All measures are residualized for data collection site. Gray matter measures were further residualized for total brain volume. ACC = anterior cingulate cortex; CT = cortical thickness; CA = cortical area; FA = fractional anisotropy; SC = subcortical volume; CON = cingulo-opercular network; l = left; r = right; FC = functional connectivity.

TableS1. Correlations among brain measures of motor processing

	Precentral CA	Precentral FA	SOMM-L Amygdala FC	SOMM--R Amygdala FC
Precentral CT	-.423	.204	.070	.023
Precentral CA		-.114	-.046	-.011
Precentral FA			-.031	.033
SOMM-L Amygdala FC				-.197

Note: SOMM = somatomotor-mouth network; FC = functional connectivity; CT = cortical thickness; CA = cortical area; FA = fractional anisotropy.

Reference

Tijssen, S., Collins, P. F., & Luciana, M. Pubertal development mediates the association between family environment and brain structure and function

in childhood. *Development and Psychopathology*. Published online 1 July 2019. doi: 10.1017/S0954579419000580.