



OPEN ACCESS

Monitoring of mitochondrial oxygenation during perioperative blood loss

Floor A Harms,¹ Alexandra R M Brandt-Kerkhof,² Egbert G Mik¹

¹Laboratory for Experimental Anesthesiology, Department of Anesthesiology, Erasmus MC, Rotterdam, Zuid-Holland, The Netherlands

²Surgery, Erasmus MC, Rotterdam, Zuid-Holland, The Netherlands

Correspondence to
Dr Floor A Harms;
f.harms@erasmusmc.nl

Accepted 31 December 2020

SUMMARY

One of the challenges in the management of acute blood loss is to differentiate whether blood transfusion is required or not. The sole use of haemoglobin values might lead to unnecessary transfusion in individual cases. The suggestion is that mitochondrial oxygen tension can be used as an additional monitoring technique to determine when blood transfusion is required. In this case report, we report mitochondrial oxygen measurements in a patient with perioperative blood loss requiring blood transfusion.

BACKGROUND

Goal-directed management of perioperative blood loss remains a major challenge for clinicians. Acute anaemia resulting in inadequate oxygen supply should be avoided at all times, but currently, no specific endpoint for personalised transfusion medicine is available. Perioperative insufficient oxygen delivery to tissues is an important determinant for postoperative complications such as stroke,¹ declined cognitive function,² kidney injury³ and cardiac ischaemia.⁴ Tissue oxygenation relies on adequate oxygen delivery, which is predominantly maintained by the arterial oxygen saturation, haemoglobin concentration and cardiac output. Transfusion of allogeneic erythrocyte concentrates plays an important role in treating acute anaemia for the prevention of tissue hypoxia. However, allogeneic blood transfusion itself is not without risks and is an independent risk factor for increased mortality and morbidity.⁵ One of the challenges in the management of anaemia is to determine whether blood transfusion is required or not. Current transfusion guidelines recommend haemoglobin levels as a trigger for red blood cell transfusion.^{6,7} These guidelines are based on mean data and thus incorporate safety margins, which might lead to unnecessary transfusion in individual cases. Additionally, more physiologically based transfusion triggers may enable optimisation and personalised treatment during the management of acute blood loss and may help in preventing transfusion-related complications like haemolytic reactions, transfusion-related acute lung injury, infections and transfusion-associated circulatory overload. In a recent study in haemodiluted pigs, mitochondrial oxygen tension (mitoPO₂) was suggested as a useful parameter to distinguish whether blood transfusion is necessary or not.⁸ Given that the mitochondrion is the final destination of oxygen, it seems logical to use mitoPO₂ as a measure for transfusion need. The mitoPO₂ can be measured by the COMET

(an acronym for Cellular Oxygen METabolism) measuring system (Photonics Healthcare, Utrecht, the Netherlands).⁹ The measurement is based on the principle of oxygen-dependent quenching of delayed fluorescence of protoporphyrin IX (PpIX).^{10,11} Application of the porphyrin precursor 5-aminolevulinic acid (5-ALA) on the skin induces PpIX in the mitochondria where it acts as a mitochondrially located oxygen-sensitive dye.¹² After photoexcitation with a pulse of green light, PpIX emits delayed fluorescence of which the lifetime is inversely related to the amount of oxygen. The technique is non-invasive and can be safely used in humans.^{13,14} In this case report, we report the results of mitochondrial oxygen measurements in a patient with major intraoperative blood loss requiring blood transfusion.

CASE PRESENTATION

A 69-year-old man with a history of diabetes mellitus (type II), hypertension and dyslipidaemia and a recent diagnosis of metastatic ascending colon carcinoma (cT4N2M1) required extensive surgery and hyperthermic intraperitoneal chemotherapy (HIPEC). His medication included metformin, gliclazide, enalapril and simvastatin. Preoperative abdominal, respiratory and cardiac examination were unremarkable. Preoperative haemoglobin levels were 93.5 g/L.

Prior to induction of anaesthesia, an epidural catheter was placed, and epidural analgesia (ropivacaine/sufentanil) was given. For induction of anaesthesia, an intravenous bolus dose of propofol 110 mg followed by rocuronium 50 mg was administered, together with a continuous infusion of remifentanyl 9 µg/kg/hour. Anaesthesia was maintained using sevoflurane. Directly after induction of anaesthesia, a continuous infusion of noradrenalin (0.40–0.60 µg/kg/min) was necessary to maintain normal blood pressure values (mean arterial pressure (MAP) >65 mm Hg).

During surgery, extensive peritoneal carcinomatosis was observed with a Peritoneal Carcinomatosis Index of 16.¹⁵ The surgeons performed a low anterior resection and an omentectomy. Thereafter, warm mitomycin C (chemotherapeutic agent) was rinsed abdominally for 1.5 hours via three inflow and two outflow catheters. Finally, a previously constructing ileostomy was reversed, and a terminal colostomy was placed.

After 1 hour of coma, during the low anterior resection, an acute rapid blood loss of 2500 mL occurred. Resuscitation of blood loss was initially done using crystalloids and colloids (figure 1). Haemodynamic parameters were kept stable,



© BMJ Publishing Group Limited 2021. Re-use permitted under CC BY. Published by BMJ.

To cite: Harms FA, Brandt-Kerkhof ARM, Mik EG. *BMJ Case Rep* 2021;**14**:e237789. doi:10.1136/bcr-2020-237789

Case report




Figure 1 Haemodynamics and transfusion during surgery. HR, heart rate; MAP, mean arterial pressure; MitoPO₂, mitochondrial oxygen tension; rHb, recombinant haemoglobin; StO₂, tissue oxygen saturation.

without the need for extra vasoactive medication. Blood transfusion was started after haemoglobin levels dropped below 88.6 g/L.

INVESTIGATIONS

In addition to the standard intraoperative monitoring consisting of invasive blood pressure measurements, peripheral oxygen saturation, 5-lead electrocardiography and temperature measurements, two additional monitoring systems were used during this operation: the COMET for monitoring mitochondrial oxygenation and the oxygen to see (O2C) (LEA Medizintechnik, Giessen, Germany) for monitoring microcirculatory blood flow velocity, tissue oxygen saturation (StO₂) and relative amount of recombinant haemoglobin (rHb) in the skin.¹⁶

For the COMET measurements, a 5-ALA patch was applied the evening before surgery. Directly after the induction of anaesthesia, the 5-ALA patch was removed, and the skin sensor of the COMET was fixated to the chest. COMET measurement was performed during surgery with an interval of one measurement per minute. The O2C probe was placed on the skin of the sternum next to the COMET measurement probe providing semicontinuous readings.

OUTCOME AND FOLLOW-UP

The mitoPO₂ value started around 70 mm Hg and slowly declined in the first hour, parallel to StO₂, to reach values around 50–60 mm Hg. After approximately 1 hour of operation time, a sudden blood loss of 2.5 L occurred. Initially, adequate haemodynamic status was ensured by infusion of crystalloids and colloids resulting in haemodilution and acute anaemia. Due to anaemia, oxygen delivery to the tissues decreased, accompanied by declining microvascular and mitochondrial parameters, while other parameters such as MAP, StO₂, rHb and lactate levels did not change during the ongoing blood loss. Heart rate and capillary blood flow did show a response to the bleeding but at a later stage than mitoPO₂. Directly after resuscitation with red blood cells, a rapid increase of mitoPO₂ was observed, with

mitoPO₂ values restored from below 10 mm Hg to up to 40 mm Hg (figure 1).

After surgery, the patient was transferred to the intensive care unit (ICU), and discharge from the ICU to a surgical ward was possible after 2 days. The stay in the surgical ward was complicated by a paracolic fluid collection, which was surgically drained. After a further trouble-free recuperation, the patient was released from the hospital 15 days after surgery. Eight months later, a CT scan diagnosed extensive recurrence of disease, eliminating curative treatment options. Unfortunately, the patient died 1 year later from the consequences of his illness.

DISCUSSION

In this case report, we show mitoPO₂ use during acute perioperative blood loss and propose mitoPO₂ as an additional monitoring parameter for perioperative transfusion management.

In the current literature, there is controversy regarding the transfusion of blood components in oncological surgery. As early as in the 1980s, the effect of blood transfusion on malignancy recurrence rate was noted.¹⁷ These findings were based on simple bivariate correlation without the adjustment of confounders. The correlation between blood transfusion and malignant recurrence rate became less obvious with the introduction of new statistical techniques whereby confounding factors were included in the analysis.^{18 19} Although the negative effect of blood components does not appear to apply in all cancer types, it is not the case in patients with peritoneal colorectal carcinomatosis undergoing cytoreductive surgery and HIPEC. Two recently published articles both showed an independent relationship between perioperative blood transfusion and survival rate, especially in patients with high-grade mucinous neoplasms.^{18 19} These findings underline the importance of blood-sparing protocols during cytoreductive surgery and HIPEC. In this case report, the potential value of perioperative monitoring of the mitochondrial oxygenation in the decision as to whether or not to transfuse a patient is shown. The mitochondria are the largest oxygen consumers within the cell. Therefore, mitoPO₂ reflects the oxygen balance between oxygen supply and oxygen demand.²⁰ Oxygen supply is dependent not only on the amount of haemoglobin but also on microvascular blood flow, the haemoglobin dissociation characteristics, the level of oxygen saturation of haemoglobin and the diffusion barriers between red blood cells and the tissue cells.²¹ Because so many factors are involved in maintaining an adequate tissue oxygenation, it doesn't seem wise to use only haemoglobin levels in the decision of transfusing red blood cells. Therefore, we suggest to use mitoPO₂ and microvascular flow measurements in combination with point-of-care haemoglobin and standard operative measurements, such as blood pressure, heart rate and pulse oximetry, in the decision-making process for blood transfusion. The main goal is to reduce the number of blood transfusions required during oncological surgery, particularly during cytoreductive surgery and HIPEC and thereby improve

Learning points

- ▶ Transfusion is not without risk; a more individualised threshold for determining blood transfusion is needed.
- ▶ Monitoring of oxygenation at the mitochondrial level is clinically possible by using the delayed fluorescence of protoporphyrin IX.
- ▶ Mitochondrial oxygenation monitoring provides a new tool for research in resuscitation and transfusion medicine.

the long-term outcome of the patients. The added value of the mitochondrial oxygenation measurements during acute perioperative blood loss must be demonstrated in future studies.

Contributors FAH: Data processing measurement and writing the article. ARMB-K: Writing the article and surgeon during operation. EGM: Head of research team.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests EGM is the founder and shareholder of Photonics Healthcare, a company aimed at making the delayed fluorescence lifetime technology available to the broad public. Photonics Healthcare B.V. holds the exclusive licenses to several patents regarding this technology, filed and owned by the Academic Medical Center in Amsterdam and the Erasmus Medical Center in Rotterdam, the Netherlands.

Patient consent for publication Parental/guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

REFERENCES

- Bahrainwala ZS, Grega MA, Hogue CW, *et al*. Intraoperative hemoglobin levels and transfusion independently predict stroke after cardiac operations. *Ann Thorac Surg* 2011;91:1113–8.
- Weiskopf RB, Feiner J, Hopf HW, *et al*. Oxygen reverses deficits of cognitive function and memory and increased heart rate induced by acute severe isovolemic anemia. *Anesthesiology* 2002;96:871–7.
- Choi YJ, Kim S-O, Sim JH, *et al*. Postoperative anemia is associated with acute kidney injury in patients undergoing total hip replacement arthroplasty: a retrospective study. *Anesth Analg* 2016;122:1923–8.
- Carson JL, Duff A, Poses RM, *et al*. Effect of anaemia and cardiovascular disease on surgical mortality and morbidity. *Lancet* 1996;348:1055–60.
- Chandra S, Kulkarni H, Westphal M. The bloody mess of red blood cell transfusion. *Crit Care* 2017;21:310.
- Dietrich W, Faraoni D, von Heymann C, *et al*. Esa guidelines on the management of severe perioperative bleeding: comments on behalf of the Subcommittee on transfusion and haemostasis of the European association of cardiothoracic Anaesthesiologists. *Eur J Anaesthesiol* 2014;31:239–41.
- Carson JL, Stanworth SJ, Roubinian N, *et al*. Transfusion thresholds and other strategies for guiding allogeneic red blood cell transfusion. *Cochrane Database Syst Rev* 2016;10:CD002042.
- Römers LHL, Bakker C, Dollée N, *et al*. Cutaneous mitochondrial PO₂, but not tissue oxygen saturation, is an early indicator of the physiologic limit of hemodilution in the pig. *Anesthesiology* 2016;125:124–32.
- Ubbink R, Bettink MAW, Janse R, *et al*. A monitor for cellular oxygen metabolism (comet): monitoring tissue oxygenation at the mitochondrial level. *J Clin Monit Comput* 2017;31:1143–50.
- Mik EG, Johannes T, Ince C. Monitoring of renal venous PO₂ and kidney oxygen consumption in rats by a near-infrared phosphorescence lifetime technique. *Am J Physiol Renal Physiol* 2008;294:F676–81.
- Mik EG, Stap J, Sinaasappel M, *et al*. Mitochondrial PO₂ measured by delayed fluorescence of endogenous protoporphyrin IX. *Nat Methods* 2006;3:939–45.
- Harms FA, Bodmer SIA, Raat NJH, *et al*. Validation of the protoporphyrin IX-triplet state lifetime technique for mitochondrial oxygen measurements in the skin. *Opt Lett* 2012;37:2625–7.
- Harms FA, Stolker RJ, Mik EG. Cutaneous respirometry as novel technique to monitor mitochondrial function: a feasibility study in healthy volunteers. *PLoS One* 2016;11:e0159544.
- Mik EG, Balestra GM, Harms FA. Monitoring mitochondrial PO₂: the next step. *Curr Opin Crit Care* 2020;26:289–95.
- McMullen JRW, Selleck M, Wall NR, *et al*. Peritoneal carcinomatosis: limits of diagnosis and the case for liquid biopsy. *Oncotarget* 2017;8:43481–90.
- Forst T, Hohberg C, Tarakci E, *et al*. Reliability of lightguide spectrophotometry (O2C) for the investigation of skin tissue microvascular blood flow and tissue oxygen supply in diabetic and nondiabetic subjects. *J Diabetes Sci Technol* 2008;2:1151–6.
- Burrows L, Tartter P. Effect of blood transfusions on colonic malignancy recurrent rate. *Lancet* 1982;2:662.
- Donohue JH, Williams S, Cha S, *et al*. Perioperative blood transfusions do not affect disease recurrence of patients undergoing curative resection of colorectal carcinoma: a Mayo/North central cancer treatment group study. *J Clin Oncol* 1995;13:1671–8.
- Dent OF, Ripley JE, Chan C, *et al*. Competing risks analysis of the association between perioperative blood transfusion and long-term outcomes after resection of colorectal cancer. *Colorectal Dis* 2020;22:871–84.
- Mik EG. Special article: measuring mitochondrial oxygen tension: from basic principles to application in humans. *Anesth Analg* 2013;117:834–46.
- Wefers Bettink MA, Arbous MS, Raat NJH, *et al*. Mind the mitochondria! *J Emerg Crit Care Med* 2019;3:45–13.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow