



Wayne State University

Neurology Faculty Publications

Department of Neurology

10-23-2014

Mechanisms of T-B cell cooperation important for MOG antibody mediated demyelination

Dusanka S. Skundric
Wayne State University, skundric@wayne.edu

Recommended Citation

Skundric, DS. Mechanisms of T-B cell cooperation important for MOG antibody mediated demyelination. http://nn.neurology.org/content/1/1/e12/reply#nnn_el;44 retrieved on 5/19/15 Letter in response to: Dale RC, Tantsis EM, Merheb V, et al. Antibodies to MOG have a demyelination phenotype and affect oligodendrocyte cytoskeleton. *Neurol Neuroimmunol Neuroinflamm*. 2014 May 22; 1(1):e12. doi: 10.1212/NXI.0000000000000012

Available at: http://digitalcommons.wayne.edu/med_neurology/5

This Article is brought to you for free and open access by the Department of Neurology at DigitalCommons@WayneState. It has been accepted for inclusion in Neurology Faculty Publications by an authorized administrator of DigitalCommons@WayneState.



Article:

Antibodies to MOG have a demyelination phenotype and affect oligodendrocyte cytoskeleton

Dale RC, Tantsis EM, Merheb V, et al. Antibodies to MOG have a demyelination phenotype and affect oligodendrocyte cytoskeleton. *Neurol Neuroimmunol Neuroinflamm*. 2014 May 22; 1(1):e12. doi: 10.1212/NXI.000000000000012

Letter in Response to Above Article:

Mechanisms of T-B cell cooperation important for MOG antibody mediated demyelination

Dusanka S. Skundric, Assistant Professor

Department of Immunology and Microbiology, Wayne State University, Detroit, MI, USA

The article by Dale et al. is significant, reporting the presence of demyelinating antibodies to MOG in pediatric relapsing ON. [1] The role of complement activating antibodies to MOG is suggested in pathogenesis of adult AQP4- ON. [2] In vitro experiments have indicated that MOG antibodies isolated from pediatric ON patients perturb organization of oligodendrocyte microtubule cytoskeleton. [1]

Similarly, pronounced demyelination, damage of axonal cytoskeleton, and oligodendrocytes were observed in experimental autoimmune encephalomyelitis (EAE) induced by immunization with MOG35-55, in (B6 x SJL)F1 (H-2b/s) mice. These mice developed relapsing-remitting disease, with prominent inflammation by CD4+ T cells followed by B220+ B cells, scattered in small perivascular infiltrates throughout spinal cord. Production of a cytokine IL-16 by infiltrating CD4+ and B220+ T cells in CNS of relapsing EAE mice, has been demonstrated. [3]

IL-16 is a CD4+ T cell specific chemotactic cytokine. This cytokine regulates fundamental biological properties of CD4+ T cells, including T - B cell cooperation. [4] Cooperation between T and B cells in regulation of autoimmune responses to MOG35-55 is critical as demonstrated by double transgenic EAE mouse models. [5] Studies in appropriately selected animal models, which resemble immunopathology and histopathology of anti-MOG+, AQP4- ON are needed.

- 1. Dale RC, Tantsis EM, Merheb V, et al. Antibodies to MOG have a demyelination phenotype and affect oligodendrocyte cytoskeleton. Neurol Neuroimmunol Neuroinflam 2014;1:e12.
- 2. Mader S, Gredler V, Schanda K, et al. Complement activating antibodies to myelin oligodendrocyte glycoprotein in neuromyelitis optica and related disorders. J Neuroinflammation 2011;8:184.
- 3. Skundric DS, Dai R, Zakarian VL, et al. Anti-IL-16 therapy reduces CD4+ T-cell infiltration and improves paralysis and histopathology of relapsing EAE. J Neurosci Res 2005;79:680-693.

- 4. Kaser A, Dunzendorfer S, Offner FA, et al. B lymphocyte-derived IL-16 attracts dendritic cells and Th Cells. J Immunol 2000;165:2474 -2480.
- 5. Bettelli E, Baeten D, Jager A, et al. Myelin oligodendrocyte glycoprotein-specific T and B cells cooperate to induce a Devic-like disease in mice. J Clin Invest 2006;116:2393-2402.

For disclosures, please contact the editorial office at nnnjournal@neurology.org.

Published October 23, 2014

http://nn.neurology.org/content/1/1/e12/reply#nnn_el;44 Retrieved 5/19/15