



Wayne State University

Wayne State University Theses

1-1-2013

Defining The Mechanism Of Enhanced Cellular Invasion Induced By Mechanical Stimulation

Snehal Sunil Ozarkar
Wayne State University,

Follow this and additional works at: http://digitalcommons.wayne.edu/oa_theses

 Part of the [Biology Commons](#), and the [Cell Biology Commons](#)

Recommended Citation

Ozarkar, Snehal Sunil, "Defining The Mechanism Of Enhanced Cellular Invasion Induced By Mechanical Stimulation" (2013). *Wayne State University Theses*. Paper 241.

This Open Access Thesis is brought to you for free and open access by DigitalCommons@WayneState. It has been accepted for inclusion in Wayne State University Theses by an authorized administrator of DigitalCommons@WayneState.

**DEFINING THE MECHANISM OF ENHANCED CELLULAR INVASION INDUCED BY
MECHANICAL STIMULATION**

by

SNEHAL OZARKAR

THESIS

Submitted to the Graduate School

of Wayne State University,

Detroit, Michigan

in partial fulfillment of the requirements

for the degree of

MASTER OF SCIENCE

2013

MAJOR: BIOLOGICAL SCIENCES

Approved By:

Advisor

Date

DEDICATION

I dedicate this thesis to

My parents, Sunil and Minal Ozarkar

And my husband, Kaustubh

Who have always believed in me and motivated me to overcome all the hurdles in life.

ACKNOWLEDGMENT

I consider myself fortunate to be a part of Department of Biological Sciences, where I have always received helpful advice from many individuals. First and foremost, I would like to express my immense gratitude to my advisor, Dr. Karen A. Beningo, for all her support and guidance I received right from my first semester here at Wayne State University. She has always encouraged me and nurtured my ability to think critically and independently. This has helped me become a better scientist. I would also like to thank my committee members, Dr. Njus and Dr. Zhang, for their time and support.

I would like to thank past and current members of Beningo lab, Indra, Shalini, Shawna, Tara, Bingqing, Imjoo and Liz, for being such wonderful labmates. I am truly grateful for their constant help throughout my project and making a great lab environment to come into every day. I want to say a very special thanks to Shalini, who has not only been a colleague but also a very good friend. Shalini, thank you for training me during my rotation and your continued help later in the lab. Your suggestions have always been extremely helpful in my experiments. I would also like to thank Kiran Koya from Meller lab, for all the help and guidance he provided for the real-time PCR experiments. I would also like to thank Kang, Cunningham and Greenberg lab for access to their laboratory equipment.

I would like to acknowledge Biological Sciences Department Staff for the friendly service they have offered since the day I joined Wayne State University. I am also thankful to Linda Van Thiel and Krystyn Purvis for giving me an opportunity to teach and being very patient with me throughout the process.

Last but not the least, this journey would not be easy without the support of my friends and family. I would like to thank my friends Anura, Ashlesha, Manasi and Zahabiya for always

being there for me and making this place home away from home. All the credit for every achievement in my life goes to my parents, Sunil and Minal Ozarkar, and all the sacrifices they have made to get me this far. I would like to thank my parents and my brother, Saket, for their immense love and support. Lastly, I want to thank my husband, Kaustubh, for being a patient listener, a comforting shoulder, and most importantly a truly wonderful friend.

TABLE OF CONTENTS

| | |
|---------------------------------|-----|
| Dedication..... | ii |
| Acknowledgements..... | iii |
| List of Tables..... | vi |
| List of Figures..... | vii |
| Chapter 1 – Introduction..... | 1 |
| Chapter 2 – Methods..... | 15 |
| Chapter 3 – Results..... | 22 |
| Chapter 4 – Discussion..... | 35 |
| References..... | 42 |
| Abstract..... | 56 |
| Autobiographical Statement..... | 58 |

LIST OF TABLES

Table 1: List of up-regulated genes in mechanically stimulated cells obtained by PCR array analysis.....25

Table 2: List of down-regulated genes in mechanically stimulated cells obtained by PCR array analysis.....25

LIST OF FIGURES

| | |
|---------------------------------------------------------------------------------------------------------------------------------------------|----|
| Figure 1: Design of <i>in vitro</i> mechano-invasion assay..... | 6 |
| Figure 2: Cancer cells show enhanced invasion in response to mechanical stimulation in the presence of fibronectin..... | 7 |
| Figure 3: Cofilin is required for enhanced invasion upon mechanical stimulation..... | 13 |
| Figure 4: Differentially expressed genes identified by PCR array..... | 24 |
| Figure 5: Genes confirmed to be differentially expressed upon mechanical stimulation..... | 26 |
| Figure 6: Transcriptional down-regulation of integrin β 3 is also reflected at protein level... | 27 |
| Figure 7: Down-regulation of integrin β 3 is required for enhanced invasion..... | 29 |
| Figure 8: No change in activation of integrin β 1 upon mechanical stimulation..... | 31 |
| Figure 9: Decrease in levels of Ser3 phospho-cofilin upon mechanical stimulation is dependent on down-regulation of integrin β 3..... | 33 |
| Figure 10: Possible pathway for enhanced invasion upon mechanical stimulation..... | 40 |

CHAPTER 1 - INTRODUCTION

Cancer and Metastasis

According to the World Health Organization (2008), cancer is a leading cause of death worldwide and 90% of these deaths result from the secondary tumors. The secondary tumors are formed by a multistep process called metastasis, which involves migration and invasion of primary tumor cells into different parts of the body (Pantel and Brakenhoff, 2004). The primary tumor mass is surrounded by a basement membrane and stroma. The basement membrane is comprised of a number of extracellular matrix (ECM) proteins such as type IV collagen, laminin, elastin, and fibronectin. The tumor stroma, on the other hand, is composed of many non-tumor cells such as fibroblasts, adipocytes and various immune cells, in addition to the ECM proteins (Bissell and Radisky, 2001). The metastatic cascade begins with the acquisition of invasive properties by the primary tumor cells. By virtue of these invasive or metastatic properties, tumor cells breach the basement membrane of the tissue, invade the stroma and via blood or lymph reach other parts of the body where they can form a secondary tumor (Chambers et al., 2002; Liotta et al., 1986; Poincloux et al., 2009). A great deal of research has been done to understand the causes of origin and spread of cancer. Nevertheless, given the extreme complexity and heterogeneity of the disease etiology, it is still difficult to pinpoint exact cause of cancer development. For decades, genetic and epigenetic factors, predisposing cancer cells to become metastatic, have been the main focus of research groups. This research has resulted in the identification of many biochemical factors such as various growth factors and cytokines that promote the metastatic behavior of cancer cells (Hirakawa et al., 2005; Huang et al., 2002; Oft et al., 1998). However, with increasing knowledge about metastatic progression cancer biologists have begun to appreciate the importance of multiple mechanical factors, such as changes in the structure and

mechanics of the tissue, as well as biophysical changes in the geometry and topology of the ECM in initiating the multi-step metastatic cascade (Kumar and Weaver, 2009).

Effect of Tumor Microenvironment on Metastasis

Along with the development of the tumor mass, structure and composition of the tumor stroma undergoes constant changes, a process known as stromagenesis (Bissell and Radisky, 2001). During this process different non-tumor cells, collectively referred to as tumor associated stromal cells, are also recruited and activated. These non-tumor cells contribute to the progression of cancer through the secretion of several biochemical factors and up-regulated synthesis of ECM proteins. By virtue of its dynamic nature, the tumor stroma provides a unique microenvironment for the tumor mass. This tumor microenvironment possesses the ability to change the neoplastic properties of the tumor. The metastatic cascade begins with changes in the tumor microenvironment which triggers the dissemination of primary tumor cells to the probable sites of secondary tumor formation (Yu et al., 2011). Examination of genetic alterations in the tumor associated stromal cells showed numerous chromosomal rearrangements not found in tumor cells (Moinfar et al., 2000). The sources of mutations in stromal cells are still unknown, but lead to the abnormal cross-signaling between cancer cells and stromal cells (De Wever et al., 2008). Therefore it is extremely important to understand the tumor-stromal interactions.

Biochemical Factors in Invasion

Many biochemical factors from the stroma have been identified as promoters of tumor metastasis. For example, immune cells present in the stroma, such as macrophages, dendritic cells, natural killer cells, and mast cells, produce chemokines, cytokines and soluble cytotoxic mediators. All these biochemical components aid in the process of angiogenesis, cell proliferation, cell motility, and invasion (Tlsty and Coussens, 2006). During tumor development, stromal cells

and cancer cells together form a paracrine loop that regulates cancer progression (De Wever et al., 2008; Pollard, 2004). For example, tumor cells secrete colony stimulating factor-1 (CSF-1) for attracting macrophages and macrophages secrete epithelial growth factor (EGF) causing tumor cell proliferation (Wyckoff et al., 2004). Tumor cells also secrete various chemoattractants such as, transforming growth factor β -1 (TGF β -1), platelet-derived growth factor (PDGF), fibroblast growth factor (FGF), and EGF, in response to which stromal fibroblast cells are attracted towards the tumor (Denys et al., 2008; Dong et al., 2004; Lederle et al., 2006; Stuelten et al., 2005). Stromal fibroblasts then undergo differentiation into myofibroblasts and secrete similar growth factors, which cause proliferation of tumor cells (Ao et al., 2007; Cat et al., 2006; Shao et al., 2006), and matrix metalloproteinases, which cause degradation and remodeling of tumor stroma (Stuelten et al., 2005; Taniwaki et al., 2007).

Mechanical Factors in Invasion

In addition to the biochemical factors, physical or mechanical factors in the tumor microenvironment also contribute to cancer progression and metastasis (Desmouliere et al., 2004; Tlsty and Coussens, 2006). The tumor mass is characterized by uncontrolled cell growth and disrupted tissue structure. This leads to the loss of tissue homeostasis and mechanical equilibrium that is maintained in the physiologically normal tissue. The physical state of the tumor also evolves with the evolution of the tumor mass (Yu et al., 2011), due to changes in the cell-cell and cell-ECM interactions, as well as deposition and remodeling of the ECM components surrounding the tumor mass (Borghi et al., 2010; Ingber, 2008; Schwartz and DeSimone, 2008). On account of these changes the tumor mass and its microenvironment are acted upon by different mechanical forces such as hydrostatic pressure, shear stress, compression and tension, and changes in compliance (Butcher et al., 2009).


One of the mechanical forces within the tumor microenvironment results from the expansion of the tumor itself. The uncontrollably growing tumor mass is confined within a limited space in the tissue. These spatial constraints generate high levels of radial stress on the surrounding stroma. At the same time, compression of the stroma generates stress within the tumor itself (Shieh, 2011). Tumor cell spheroids embedded in agarose gels have been shown to generate stresses more than 10 kPa in in-vitro studies (Helmlinger et al., 1997). Interstitial fluid pressure also contributes to compressive forces within the tumor and its microenvironment. It is formed due to highly permeable microvascular networks leading to increased flow of fluids and macromolecules within the tumor and stroma (Boucher and Jain, 1992; Fukumura et al., 2010; Hagendoorn et al., 2006). Nonetheless, most studies of mechanical forces in the tumor and microenvironment have focused on compliance. After all, stiffer stromal tissue has long been a hallmark of different cancer types such as breast, liver and prostate cancer (Boyd et al., 1999; Gang et al., 2009; Hoyt et al., 2008). Mammary tumor tissue is known to have stiffness exceeding 4 kPa, which is much higher than a normal mammary gland that has an elastic modulus of 200 Pa (Paszek et al., 2005). This increase in stiffness or rigidity of the stroma is due to the abnormal production of type I collagen and fibronectin by different stromal cells (Bissell and Radisky, 2001; Tlsty and Coussens, 2006). Recent studies have shown that this rigidity of the tumor stroma positively regulates cancer cell proliferation and dissemination to other sites within the body (Kostic et al., 2009; Levental et al., 2009; Paszek et al., 2005).

Stromal cells associated with tumors also give rise to different mechanical forces, which can promote metastasis. Myofibroblasts are one of the main types of stromal cells that appear early during tumor progression. They were initially discovered to be involved in the process of wound healing (Ryan et al., 1974). In the tumor stroma, myofibroblasts differentiate from normal fibroblasts, fibrocytes, mesenchymal stem cells (MSCs) and other cells of mesenchymal origin

(De Wever et al., 2008; Desmouliere et al., 2004). This differentiation requires various paracrine signals from the cancer cells. TGF β -1 has been found to play a significant role in this process (Desmouliere et al., 1993). The hallmarks of differentiated myofibroblasts are secretion of ECM proteins like collagen and fibronectin as well as neo-expression of α -smooth muscle actin (α -SMA) (Darby et al., 1990; Serini et al., 1998). This actin isoform bundles with myosin-II and forms contractile stress fibers. A direct correlation between the level of α -SMA expression and myofibroblast contractions has been demonstrated both *in vitro* (Hinz et al., 2001a) and *in vivo* (Hinz et al., 2001b). Myofibroblasts are also known to produce large focal adhesions, through which tugging or pulling forces produced by stress fibers are transferred into the surrounding ECM (Castella et al., 2010; Hinz et al., 2003). Using a unique *in-vitro* invasion assay developed in our lab, it has been shown that mechanical stimuli in the form of pulling forces leads to enhanced invasion by cancer cells (Menon and Beningo, 2011). Thus, it is important to study the mechano-sensitivity of tumor cells to these forces generated by stromal cells and to understand how ECM remodeling, mechanical regulation and stromal cell activities contribute to cancer progression.


***In vitro* Mechano-Invasion Assay**

As mentioned earlier, contractile stromal cells such as myofibroblasts can produce pulling forces in the tumor microenvironment. We have developed an *in vitro* mechano-invasion assay, where cells can be subjected to similar pulling forces in an environment free of any cell secreted factors and the effect of these forces on the invasion response of the cells can be observed. The general set-up of the assay is illustrated in figure 1. In preparation for the invasion assay, substrates made of collagen and fibronectin were prepared. These two proteins are the most abundant ECM proteins in the tumor stroma. Paramagnetic beads, functionalized by carboxylation for protein binding, were also mixed with these collagen/fibronectin substrates. Mechanical stimulation was provided by holding these substrates over a rare earth magnet that is rotated on



an orbital shaker beneath the culture. The human fibrosarcoma cell line, HT1080, was used for this assay. HT1080 cells cultured on collagen/fibronectin matrices incubated without magnetic stimulation were used as controls (Figure 1).

Magnetic stimulation was provided for 24 hours and the number of cells that invaded into the matrix were counted using an inverted phase microscope. HT1080 cells under mechanical stimulation showed enhanced invasion (2-fold) compared to control cells. Similar enhanced invasion was also seen at 36 hours of mechanical stimulation (Figure 2A). This enhanced invasion response by HT1080 cells was stable for frequencies of bead stimulation ranging from 8 to 160



rpm. Furthermore, non-invasive mouse embryonic fibroblast (MEF) cells failed to invade in the presence, as well as the absence, of mechanical stimulation, demonstrating that in order to invade upon mechanical stimulation, cells must have a pre-defined ability to invade.

To test the importance of fibronectin in the process of enhanced invasion, the assay was performed in collagen only substrates and compared to collagen/fibronectin substrates. After 24

hours of mechanical stimulation, cells seeded on collagen only substrates failed to show enhanced invasion compared to non-stimulated controls (Figure 2B). This demonstrates the requirement of fibronectin in the extracellular environment for the cancer cell to sense the mechanical stimulation.

Mechanosensing and Mechanotransduction

In order for the cell to respond to mechanical cues in its surrounding, the cells must first be able to sense these stimuli. The process of sensing mechanical stimuli is known as mechanosensing. Secondly, cells should be able to transduce these stimuli within their cytoplasm resulting in changes in gene expression and intracellular biochemistry of the cell, by the process known as mechanotransduction (Wang et al., 2009). The signal assimilated into the cytoplasm is carried forward through a hierarchical mechanochemical signal transduction network that results in the cellular response to the mechanical stimulus. In the case of cancer cells, this response can be in the form of increased proliferation and invasion or secretion of proteases for matrix remodeling (Kumar and Weaver, 2009). Since the mechanical signal spreads once it crosses the cell membrane, this conversion of the mechanical signal into a biochemical signal likely takes place nearing close proximity to the cell membrane. Indeed, many different transmembrane proteins that connect the ECM to the inside of the cell can act as mechanotransducers (Orr et al., 2006). Integrins are one of the most studied families of transmembrane receptors known to act as mechanotransducers. As the name suggests, they integrate the outside of the cell with the inside. However, additional membrane receptors can also act as mechanotransducers. For example, abolishment of G-protein coupled receptors (GPCR) in neutrophils has been shown to inhibit the cellular response to fluid shear stress, implicating GPCRs as mechanosensors (Makino et al., 2006). Unlike all the above mentioned receptors, which bring about mechanotransduction

between ECM and cell, E-cadherin transduces the mechanical signal through cell-cell junctions with the help of other binding partners such as α -catenin and vinculin (Smutny and Yap, 2010).

Integrins are heterodimeric glycoproteins composed of one α -subunit and one β -subunit. There are eighteen α -subunits and eight β -subunits which non-covalently pair with each other to form 24 different integrin molecules (Plow et al., 2000). Both α and β -subunits have larger extracellular domains with which they bind to a specific ligand in the extracellular matrix. The cytoplasmic tail of these subunits is non-catalytic and shorter compared to the extracellular domain. The cytoplasmic tail of the β -subunits bind directly to the actin cytoskeleton of the cell, thus providing a connection between the outside and the inside of the cell (Rathinam and Alahari, 2010; Springer and Wang, 2004). The cellular response to different mechanical forces such as, fluid shear stress, stretch, hydrostatic pressure, has been shown to be dependent on the type of integrin that is engaged (Katsumi et al., 2004). Since many signaling cascades are controlled downstream of integrins, these mechanical forces can activate a variety of pathways. The outcome of this phenomenon changes from cell type to cell type and is also dependent on the type of integrin and mechanical forces involved (Schwartz and DeSimone, 2008). Considering their important role in mechanosensing and mechanotransduction, integrins have been implicated in cancer cell proliferation and migration. Even though detailed specifics vary with cancer type, in general the integrin gene expression in cancer cells is found to be in disarray. These defective expression levels of integrins, also lead to abnormal signaling downstream causing the progression of cancer (Guo and Giancotti, 2004). Higher expression levels of certain integrin subunits such as $\alpha 3$, $\alpha 5$, $\alpha 6$, αv , $\beta 1$, and $\beta 3$ have been correlated with greater metastatic potential of cancer cells, whereas, some integrins such as $\alpha 2\beta 1$ have been associated with decreased cell proliferation (Wiseman and Werb, 2002).

Integrin β 3

Integrin β 3 is one of several β -subunits of integrin receptors which non-covalently binds with the α v or α IIb subunits (Hynes, 2002). β 3 containing integrins bind to Arg-Gly-Asp (RGD) sequence containing ligands in the extracellular matrix, which include fibronectin, vitronectin, and fibrinogen (Xiong et al., 2002). Expression levels of integrin β 3 coupled with integrin α v are shown to be variable in different types of cancers. They are highly up-regulated in cases of melanoma, squamous cell carcinoma and ovarian cancer. Whereas, down-regulated in cases of kidney and colon cancer (Mizejewski, 1999). Integrin β 3 has been shown to affect different processes during cancer development and progression such as cell proliferation, apoptosis, angiogenesis and cell invasion (Jin and Varner, 2004). Integrin α v β 3 up-regulation is known induce cell death (Stupack et al., 2001). This agrees with the recent finding that down-regulation of ITGB3 in glioma and hepatocellular carcinoma renders cancer cells resistant to apoptosis and therefore causes aggressive tumor growth (Kim et al., 2011; Wu et al., 2009). Up-regulation of ITGB3 has been seen in melanoma and breast cancer and it is associated with increased tumor invasion and metastasis (Albelda et al., 1990; Gasparini et al., 1998). As integrin β 3 is known to aid in the process of tumor progression by increasing cell proliferation and invasion, it has become a potential target for cancer therapy. Different types of blocking peptides and antibodies against integrin β 3 are currently in clinical and pre-clinical trials (Eskens et al., 2003; Smith, 2003; Stoeltzing et al., 2003).

Integrin Crosstalk

As mentioned earlier, integrin heterodimers bind to specific ligands in the ECM. Integrins in the ligand bound unfolded conformation are said to be “active”, whereas those in unbound folded conformation are said to be “inactive” (Takagi et al., 2002). It has been observed that when

activity of one integrin subunit or the entire heterodimer is inhibited, it sometimes causes perturbation of ligand binding ability of another integrin family member within the same cell. This phenomenon where the activity of one integrin is controlled by other integrin is referred to as “integrin crosstalk” (Diaz-Gonzalez et al., 1996; Gonzalez et al., 2010). Evidence of integrin crosstalk has been reported between integrin $\beta 3$ and integrin $\beta 1$. Integrin $\beta 1$ is also one of the β subunits which can form heterodimers with 12 different α subunits. Integrin $\alpha 5\beta 1$ is a RGD-binding heterodimer, like integrin $\beta 3$, and it primarily binds to fibronectin (Hynes, 2002). In endothelial cells, reciprocal inhibitory activity of integrin $\alpha \beta 3$ and integrin $\alpha 5\beta 1$ has been observed (Brooks et al., 1994; Eliceiri and Cheresh, 1999). Moreover, in vivo experiments in mice have shown that down-regulation of integrin $\alpha \beta 3$ expression during tumor associated angiogenesis leads to increased activity of integrin $\beta 1$ (Reynolds et al., 2002). This integrin crosstalk is regulated by different proteins binding to the cytoplasmic tail of the integrins, for example talin. In some instances, it is also regulated by the phosphorylation state of the integrin cytoplasmic tails. There is also evidence that this integrin crosstalk is brought about, not only at the protein level, but also at the level of mRNA. It has been shown that integrin $\beta 1$ destabilizes the mRNA encoding integrin $\beta 3$, therefore decreasing the activity of integrin $\alpha \beta 3$ (Retta et al., 2001). Activity of a particular integrin matters because downstream signaling under different integrins is different (Morgan et al., 2009). Thus, the same type of cell having different integrin activity can produce totally different cellular response. For example, it has been demonstrated that the activity of Rho GTPases is regulated differently downstream of integrin $\alpha \beta 3$ and $\alpha 5\beta 1$ (Morgan et al., 2009). Cells with active $\beta 3$ integrins have low RhoA activity and show persistent migration, conversely, cells with active $\beta 1$ integrins have high RhoA activity and show random migration (Danen et al., 2005).

Invadopodia: Mechanosensory Protrusions

In 1971 Gabbiani and colleagues first discovered specialized actin-rich protrusions in Rou-

s sarcoma virus (RSV) transformed chicken embryonic fibroblasts making contacts with the ECM and having ECM degradation properties (Gabbiani et al., 1971). These protrusions were named invadopodia and were found to be a characteristic unique to invasive cancer cells and transformed cells. Matrix degradation by invadopodia enhances the process of cancer metastasis *in vivo*, by helping cancer cells breach the basement membrane and navigate through the dense stromal ECM. The process of breaching the basement membrane via invadopodia consists of three stages; 1) formation of invadopodia, 2) maturation of invadopodia by their elongation and recruitment of ECM degrading proteases and finally, 3) degradation of ECM (Schoumacher et al., 2010). The actin core formation is the principle step in invadopodia formation. Hence, different molecules required for actin nucleation and polymerization such as, Arp2/3, N-WASP, cortactin, have been shown to be required for this process. However, the molecule that stands out from these is an actin severing protein cofilin. Cofilin has been shown to be required only for invadopodia maturation and not for the formation stage. When cofilin is knocked down, cells give rise to short-lived invadopodia which are not capable of ECM degradation (Yamaguchi et al., 2005).

Cofilin is an actin binding protein that belongs to a family of “actin depolymerizing factors (ADF)” (Maciver and Hussey, 2002). Cofilin is actively involved in actin tread-milling as it binds to capped ends of the actin filaments and severs them. This function of cofilin is important in two ways; 1) the severing action keeps on refilling the actin monomer pool, 2) it creates more barbed ended actin filaments which can be further elongated (Pollard and Borisy, 2003). The actin depolymerization activity of cofilin is controlled by its phosphorylation at a serine-3 near the N-terminus by LIM-kinase. When phosphorylated, cofilin cannot interact with ADP-actin filaments of monomers and therefore, inactivated (Blanchoin et al., 2000). Rho family GTPases indirectly control the cofilin activity. Active Rho family GTPases activate PAK kinase which in turn activates

LIM-kinase leading to cofilin phosphorylation (Edwards et al., 1999). A phosphatase called slingshot causes dephosphorylation of serine residue and activates cofilin (Niwa et al., 2002). By virtue of its actin depolymerizing ability, cofilin plays an important role in cell motility. Recently it has also been observed that expression profiles of cofilin and its regulators are highly disturbed in invasive cancer cells (Wang et al., 2007). Moreover, in our *in vitro* invasion assay, we found that cofilin knock-down cells fail to show enhanced invasion in response to mechanical stimulus in their microenvironment (Figure 3; Menon and Beningo, 2012). Even though cofilin is closely related to invadopodia maturation and cancer cell invasion, its exact role in the process is still unclear.

In addition to cofilin, many other invadopodia associated proteins have been shown to be required for tumor growth and metastasis (Blouw et al., 2008; Clark et al., 2009). However, the effect of external mechanical forces on the activity of invadopodia is still unknown. Recently it was shown that increased stiffness of ECM leads to enhancement in the activity and number of invadopodia (Alexander et al., 2008). Moreover, localization of active forms of mechanosensing




Figure 3. Cofilin is required for enhanced invasion upon mechanical stimulation. HT1080 cells nucleofected with control siRNA or Cofilin siRNA and cultured for 48 hours were seeded onto collagen/fibronectin matrices containing paramagnetic beads. The cells were cultured with or without stimulation for 48hr hours and the percent of invading cells was calculated. Stimulated cells had 3-fold higher invasion as compared to non-stimulated cells ($P= 0.0065$) (Menon and Beningo; 2012).

proteins such as FAK (focal adhesion kinase) and p130Cas suggests that invadopodia are sensitive to mechanical forces in the tumor microenvironment (Parekh and Weaver, 2009). Therefore, the molecular dissection of the process of invadopodia formation and maturation in response to mechanical forces will be interesting.

In conclusion, the tumor stroma serves as a potent carcinogen by being a source of numerous mechanical factors. Recent findings suggest that many of these factors are both necessary and sufficient for the process of cancer progression. Even though influence of mechanical factors is positively linked with enhanced invasion, it is important to integrate this information into our existing knowledge of the molecular and cell biology of cancer. Identification of signaling pathways and molecules related to mechanosensing in cancer can provide new directions to chemotherapeutics, although much remains to be uncovered.

CHAPTER 2 - METHODS

Cell Culture

All the experiments in this study were performed with human fibrosarcoma cells (HT1080) purchased from ATCC. As per ATCC recommendation, these cells were cultured in Eagle's Minimum Essential Medium (EMEM; ATCC) along with 10% fetal bovine serum (FBS; Hyclone) and 1% Penicillin-Streptomycin solution (100 U/ml of penicillin and 100 µg/ml of streptomycin; Gibco by Life Technologies). The cells were maintained in a culture incubator at 37°C with 5% CO₂. Cells were allowed to grow up to 75-80% confluency before passing them into fresh culture dishes. For this purpose, cells were trypsinized with 0.25% trypsin (Sigma) and proteolysis was stopped by addition of fresh complete media. For all experiments, cells up to consecutive sixth passage were used. After the sixth passage all cells were discarded.

Invasion Matrices

Invasion assay matrices were composed of 2.5 µg/ml collagen type I (PureColl, Advanced Biomatrix), 20 µg/ml fibronectin (Sigma). 4 µl of 1 µm carboxylated paramagnetic microspheres (Polysciences Inc.) were added for each 1 ml of matrix solution to provide magnetic stimulation. 0.1N NaOH and 10X PBS were used to adjust the pH to 7.4 +/- 2. All the components were kept cold and mixed together at 4°C. 450 µl of this solution was then polymerized in a culture well, by incubating at 37°C for 30 min. The culture well, 1 mm deep and 2 cm in diameter, was created in 60 mm culture dish (Nunclon) by drilling a hole and attaching an activated coverslip (described in Beningo and Wang, 2002) at the bottom. During polymerization a 25 mm coverslip was dropped on top of the matrix to make the surface flat. For mRNA and protein extraction, 1.1 ml of matrix

mix of the same composition was polymerized in larger wells of 1 mm depth and 3 cm in diameter. The polymerization was carried out at 37°C for 60 min. After polymerization, 4 ml of warm media was added to the plates and coverslips on the top were removed. The matrices were then sterilized under ultraviolet light for 15 min within a laminar flow culture hood. “Collagen only” matrices were prepared in the same manner without adding fibronectin to the matrix mix.

***In vitro* Mechano-Invasion Assay**

For the invasion assay, 1.5×10^4 cells were counted by hemocytometer and seeded onto the sterilized matrix. The cells were allowed to adhere and spread for 1 hour in the incubator at 37°C with 5% CO₂. To provide mechanical stimulation, the culture dish with seeded matrix was held over a rare earth magnet 25 mm in diameter and 5.5 mm in thickness with magnetic field of 12,100 Gauss. The distance between matrix and the magnet was kept constant to 1.5 cm for all experiments. An orbital shaker (Barnstead Thermolyne, Roto Mix-Type 50800) was used to rotate the magnet at 160 rpm (2.6 Hz) below the seeded collagen/fibronectin matrix. Another matrix seeded with cells was incubated separately without magnetic stimulation served as a control. At 24 hours (48 hours for integrin $\beta 3$ overexpression analysis) the invasion response of cells was quantified by selecting 10 random microscopic fields of matrix under 10X phase objective on an Olympus IX81 microscope. Cells on the surface of the matrix were counted and then invaded cells were counted in eight focal planes separated by 100 μm , from 5 μm to 800 μm below the matrix surface. The percent invasion was counted as a percent of number of invaded cells compared to total number of cells counted.

Collagenase Degradation

A 2 mg/ml solution of collagenase was prepared by dissolving collagenase type 4 (Worthington Biochemical Corporation) in Hank's Balanced Salt Solution (HBSS, Gibco by Life

Technologies) warmed to 37°C. The media on top of the collagen/fibronectin matrix was aspirated and the matrix was physically removed from the culture well with a spatula. Each matrix (3 cm in diameter) was dropped into a tube containing 2 ml of collagenase solution and kept in water bath at 37°C. With intermittent shaking for approximately 10 min, the collagen matrix was degraded. Cells were separated by centrifugation at ~500 x g for 5 min in eppendorf centrifuge 5810 R at 37°C. The resulting pellet contained both whole cells and the paramagnetic beads from the collagen/fibronectin matrix. The pellet was washed with sterile 1X PBS at 37°C.

RNA Extraction

The RNA was extracted from invaded cells after the matrix had been degraded as mentioned above. The original matrices had been seeded with 7×10^4 HT1080 cells. For each independent experiment, control and stimulated matrices were made in duplicates. During collagenase degradation cells from duplicate cultures were pooled together. Qiagen RNeasy Mini kit was used for RNA extraction from pelleted cells. 700 μ l of lysis buffer from the kit was added to the pellet. To prevent plugging the column, paramagnetic beads from the pellet were pulled down with a magnet, before loading the lysate on to the column. Genomic DNA contamination was removed by Qiagen on-column DNaseI digestion. RNA was eluted in 20 μ l of DNase/RNase free water. The quality of the acquired RNA was tested spectrophotometrically using a NanoDrop spectrophotometer (Thermo Scientific). Only RNA samples having $260/280 \geq 2.0$, $260/230 \geq 1.7$ and concentration $\geq 40 \mu\text{g/ml}$ were used for further studies.

PCR Array and qPCR Analysis

RNA obtained from both control and mechanically stimulated cells from the invasion assay were used to make cDNA that was used for PCR array as well as qPCR analysis. For each experiment, 1 μg of RNA from each control and stimulated sample was converted into cDNA using

RT² First Strand Kit (SA Biosciences, for PCR array analysis) or GoScript™ Reverse Transcriptase (Promega, for qPCR). To identify differentially expressed genes between non-stimulated and stimulated conditions from the invasion assay, the following PCR arrays were purchased from SA Biosciences; Cell Motility PCR Array, Tumor Metastasis PCR Array and ECM and Adhesion Molecules PCR Array. PCR array analysis was performed using RT² qPCR SYBR Green/ROX MasterMix-12 (SA Biosciences) on a Stratagene Mx3000P instrument. The raw data obtained were analyzed using the web-based RT² Profiler PCR Array Data Analysis software (SA Biosciences). For confirming the differential expression, qPCR primers were designed for the genes showing more than 1.35 fold differential expression in PCR array analysis. For designing qPCR primers online PrimerQuest software (Integrated DNA technologies) was used. The following primers were synthesized; GAPDH (GAPDH-F: 5'-TTCGACAGTCAGCCGCATCTTCTT- 3', GAPDH-R: 5'- ACCAAATCCGTTGACTCCGACCTT- 3'), ACT3 (ACT3-F: 5'- CAATGGCCTCAAACATGCTGCT- 3', ACT3-R: 5'- TCTCTTCAGCA CCAATGGACACCA- 3'), SRC (SRC-F: 5'- ATCCTACTGTGTGCTGGAAAGCGA- 3', SRC-R: 5'- GGTGCAGATGTTCAAAACAGCCA-3'), MMP14 (MMP14-F: 5'- TGATGGATGGATACCCAA TGCCCA- 3', MMP14-R: 5'- CGCCTCATCAAACACCCAATGCTT- 3'), CTTN (CTTN-F: 5'-TCCA AAGGATTCGGCGGGAAGTAT-3', CTTN-R: 5'-ACCTGGGTGACATCCTCAAAGGTT-3'), DIAPH1 (DIAPH1-F: 5'-TGAAGGCTTGGCCTCCTTATTGGA-3', DIAPH1-R: 5'-TCTCATGCTT GTTCCGGCTATCGT-3'), MYL9 (MYL9-F: 5'-GGCCACATCCAATGTCTTCGCAAT-3', MYL9-R: 5'-AGCCATCACGGTTCTGGTCAATCA-3'), PTK2B (PTK2B-F: 5'-AGAAGTTCATGAGCGAGG CAGTGA-3', PTK2B-R:5'- ATTCCATGATGATCCAGGTGGGCT-3'), ITGB3 (ITGB3-F: 5'- TGGACAAGCCTGTGTCACCATAACA-3', ITGB3-R: 5'-TTGTAGCCAAAC ATGGGCAAGCAG-3'), IGF1 (IGF1-F: 5'-TGAAGATGCACACCATGTCCTCCT-3', IGF1-R: 5'-AACTGAAGAGCATCCA CCAGCTCA-3'), MTSS1 (MTSS1-F: 5'-ATCAAGATGGGCTTTGCCGTTTCC-3', MTSS1-R: 5'-

AGCCAAACCGCTCTGTAGGGTATT-3'), TSHR (TSHR-F: 5'-ACCCGTGTGAAGACATAATGGGCT-3', TSHR-R: 5'-AGGACAAAGACATTGCCAGGAGA-3'). The GAPDH gene was used as the normalizing gene in the analysis. The qPCR analysis of individual genes was performed using RT² qPCR SYBR Green/ROX MasterMix-12 (SA Biosciences) on Stratagene Mx3000P instrument. For every gene, at least two biological replicates were performed and for every biological replicate two technical replicates were performed. The raw data obtained were analyzed using Stratagene Mx-Pro Mx3000P software. Student's t-test was performed to determine the statistical significance of difference in gene expression.

Integrin β 3 Overexpression

A plasmid with human ITGB3 gene, pcDNA3.1-beta-3, provided by Dr. Timothy Springer (Addgene plasmid #27289), was used for integrin β 3 overexpression in HT1080 cells. HT1080 cells were grown to approximately 85% confluency and nucleofected using the Amaxa Nucleofector 2 with the Amaxa nucleofector kit T (Lonza). 4 μ g of pcDNA3.1-beta-3 plasmid were used for each nucleofection. The cells were also cotransfected with pmaxGFP[®] supplied with the kit, to determine the efficiency of nucleofection. For mock nucleofection, HT1080 cells were also transfected with pmaxGFP or with Amaxa nucleofector kit T (Lonza) alone. After nucleofection, cells were seeded into 100 mm culture dish for 24 hours and incubated at 37°C with 5% CO₂. In case of invasion assays performed for protein extractions, nucleofected cells were directly seeded on to collagen/fibronectin matrices prior to incubation.

Protein Extraction

Triple detergent lysis buffer (TDLB; 50 mM Tris HCL pH 8.0, 150 mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, and 0.1% SDS) mixed with Protease Inhibitor Cocktail (Sigma), and Halt[™] Phosphatase Inhibitor Cocktail (Thermo Scientific) was used for protein extraction for

western blot analysis. In studies checking for integrin $\beta 3$, proteins were extracted from HT1080 cells, mock nucleofected HT1080 cells and HT1080 cells nucleofected with integrin $\beta 3$ plasmid at 24, 48 and 72 hours post nucleofection. All cells were rinsed with 1X PBS and incubated with TDLB for 20 min under ice-cold conditions. Lysates were centrifuged at 4°C for 10 min at ~5500 x g to remove cell debris. Protein estimation was done using the DC assay (Bio-Rad).

For protein extraction from invaded cells, collagen/fibronectin matrices were prepared as mentioned above and seeded with HT1080 cells, mock nucleofected HT1080 cells or HT1080 cells nucleofected with integrin $\beta 3$ plasmid. For each experiment, control and stimulated matrices were made in duplicates. After 48 hours of mechanical stimulation, matrices were degraded with collagenase and cells were pelleted down by centrifugation as described above. The cell pellet was incubated with 250 μ l TDLB for 20 min under ice-cold conditions. The paramagnetic beads were separated from lysates with a magnet. Lysates were centrifuged at 4°C for 5 min at ~7000g to remove cell debris. The supernatant was again mixed with separated paramagnetic beads, in order to avoid loss of protein bound to paramagnetic beads. This solution was then mixed with 50 μ l of 6X lamelli buffer (reducing or non-reducing, based on the antibody to be used) and boiled for 10 min. Immediately following boiling, paramagnetic beads were separated from the solution using a magnet, without letting the solution cool down. Protein estimation was done by the RC-DC assay (Bio-Rad).

Western Blots

Western blots were performed with protein samples collected from HT1080 cells, mock transfected HT1080 cells, and integrin $\beta 3$ transfected HT1080 cells. Invasion assays were performed with all the above cell conditions and western blots were performed with protein samples collected from each condition following 48 hours of mechanical stimulation. 15 μ g – 25

μg of protein was run on a 4-20% Precise™ Protein Gel (Thermo Scientific) for one hour at 100 V. Proteins were then transferred onto a PVDF membrane (Bio-Rad) using a Transblot® SD Semi-Dry Transfer Cell (Bio-Rad) at 20 V for 30 min. Following transfer, the PVDF membrane was blocked for 1 hour using 5% milk in 0.1% PBS/T (for GAPDH and cofilin), 5% milk in 0.1% TBS/T (for integrin $\beta 3$ and integrin $\beta 1$) or 5% BSA in 0.1% TBS/T (for phospho-cofilin (ser3)). Primary antibody dilutions were made in the same solutions used for blocking (except for integrin $\beta 3$ for which 1% milk in 0.1% TBS/T was used) and incubated at 4°C overnight (except for cofilin; 4 hours at RT). The following day, the membrane was washed three times for 10 minutes each either with 0.1% PBS/T (GAPDH and cofilin) or 0.1% TBS/T (phospho-cofilin (ser3), integrin $\beta 3$ and active integrin $\beta 1$). Secondary antibody dilutions were made in the same solution as that of primary antibody and incubation was carried out at RT for 1 hour. The following antibodies were used for these studies; rabbit polyclonal integrin $\beta 3$ antibody (1:300, Santa Cruz Biotechnology), rat monoclonal active integrin $\beta 1$ antibody (1:5000, BD Pharmingen™), mouse monoclonal cofilin antibody (1:300, Abcam), rabbit monoclonal phospho-cofilin (ser3) antibody (1:1000; Cell Signaling Technology), mouse monoclonal GAPDH antibody (1:15000; Millipore), HRP tagged anti-mouse antibody (Fisher), HRP tagged anti-rat (Abcam), HRP tagged anti-rabbit (Amersham GE Healthcare). GAPDH served as a loading control in all experiments. Following secondary antibody incubation, membranes were washed again with 0.1% PBS/T or 0.1% TBS/T three times for 10 minutes each and for detection incubated with Amersham ECL Prime Western Blotting Detection Reagent. After development, band intensity readings were taken using ImageJ software.

CHAPTER 3 - RESULTS

Many types of mechanical forces are produced in the tumor stroma. With the growing awareness about these mechanical forces, a new door for research has opened with respect to their effect on the process of metastasis. Despite the different known mechanical forces produced in the tumor microenvironment, efforts have focused heavily on only one kind of mechanical force, increased rigidity of the stroma. Recently it has been shown that the rigidity of tumor stroma is responsible for an increase in number and activity of invadopodia as well as cell proliferation and dissemination (Kostic et al., 2009; Levental et al., 2009; Paszek et al., 2005). We have developed an *in vitro* mechano-invasion assay, where we tested the impact of mechanical stimuli in the form of pulling forces on the cells ability to invade the stroma. We have found a significant increase in the invasion efficiency of cells in this mechanically stimulated culture environment compared to that in a non-stimulated culture environment (Menon and Beningo, 2011). However, the mechanistic details about this process of mechanosensing still remain elusive. Our objective is to uncover candidate genes involved in this mechanosensing pathway, which ultimately leads to cancer metastasis. Based on our preliminary data, we hypothesized that in response to mechanical forces in the stroma, cancer cells will show an altered expression of genes involved in mechanosensing. To test our hypothesis, we used real-time PCR analysis to identify differentially expressed genes in mechanically stimulated cancer cells compared to non-stimulated cancer cells.

RNA Extraction Standardization

In order to obtain an RNA yield sufficient for real-time PCR experiments, the previously described *in vitro* invasion assay was scaled up. For scaling up, HT1080 cells were seeded on larger collagen/fibronectin substrates (3 cm in diameter) and mechanical stimulation was provided


for 48 hours. For real-time PCR it is recommended to use RNA with $260/280 \geq 2.0$ and $260/230 \geq 1.7$, in order to have RNA free of protein and chemical contamination. As the cells were invaded into the collagen/fibronectin matrix, it was necessary to degrade the matrix to isolate the cells for RNA extraction. For standardization of matrix degradation and RNA extraction two different methods were used. In the first method, the collagen/fibronectin matrix was degraded with TRIzol, which degrades the matrix and lyses the cells simultaneously. Therefore, the matrix degradation will be directly followed by RNA extraction with chloroform as per the TRIzol RNA extraction protocol. Despite numerous attempts, this method consistently produced values of 260/230 ratio significantly less than 1.7 indicating phenol contamination in the purified RNA. As an alternative approach, the degradative enzyme collagenase in Hanks's balanced salt solution at 37°C was used to degrade the matrix. However, as collagenase is known to have a low tryptic and protease activity, the concentration of collagenase solution to be used for matrix degradation, without damaging the cells, required standardization. The concentration of collagenase was standardized to 2 mg/ml which degraded the matrix within 10 min. The cells were separated by centrifugation and RNA was extracted with the Qiagen RNeasy mini kit. Any genomic DNA contamination was removed by Qiagen on-column DNaseI digestion. An RNA with sufficient concentration and of high quality was consistently obtained using this method and therefore it was used for RNA extraction.

Identification of Differentially Expressed Genes by PCR Array Analysis

For identification of differentially expressed genes under stimulated and non-stimulated conditions, three sets of PCR arrays were purchased from SA Biosciences. Each PCR array contained primers against 84 candidate genes related to cell motility, tumor metastasis and ECM and adhesion molecules. The RNA was obtained from mechanically stimulated and non-stimulated cells as described above and was then used to prepare cDNA. This cDNA was used

for PCR array analysis. The gene expression was normalized with five housekeeping genes, actin (ACTB), β -2-microglobulin (B2M), glyceraldehyde-3-phosphate dehydrogenase (GAPDH), hypoxanthine phosphoribosyltransferase 1 (HPRT1) and large ribosomal protein (RPL13A). The differentially expressed genes identified by PCR array analysis were classified into 7 groups based on their function (Figure 4). With cut-off value of 1.35, 46 genes were found to be differentially expressed, out of which 39 genes were up-regulated (Table 1) and 8 genes were down-regulated (Table 2).

From the differentially expressed genes listed in table 1 and 2, select genes were chosen for further confirmation of gene expression by qPCR. The factors for selection were based on literature survey, genes with known association with cancer cell invasion and genes associated with mechanosensing. The genes chosen were, CTTN (cortactin), DIAPH1 (diaphanous homologue 1), SRC (V-src sarcoma (Schmidt-Ruppin A-2) viral oncogene homolog), TSHR (thyroid stimulating hormone receptor), ITGB3 (integrin β 3), MTSS1 (metastatic suppressor 1),



| Up-regulated genes (> 1.35 fold) | | | | | |
|--------------------------------------------|------------------------|-----------------------------|------------------------|---------------------------|------------------------|
| Gene name | Fold regulation | Gene name | Fold regulation | Gene name | Fold regulation |
| Cell adhesion genes | | Membrane projections | | ECM proteins | |
| BCAR1 | 1.40 | CTTN [†] | 1.47 | COL14A1 | 2.35 |
| LAMA2 | 1.62 | DIAPH1 [†] | 1.49 | COL7A1 | 1.75 |
| PNN | 1.42 | RDX | 1.39 | VCAN | 1.72 |
| ROCK1 | 1.37 | SRC [†] | 1.79 | HAS1 | 1.61 |
| | | | | SPP1 | 1.48 |
| Cell growth and proliferation | | Proteases | | THBS3 | 1.64 |
| CCL7 | 3.22 | ADAMTS13 | 1.57 | VTN | 1.64 |
| CXCR4 | 1.43 | ADAMTS8 | 1.69 | | |
| DENR | 1.43 | MMP10* | 2.59 +/- 0.8 | Protease inhibitor | |
| GNRH1 | 1.86 | MMP12 | 2.20 | KAL1 | 1.47 |
| KISS1R | 1.70 | MMP13 | 1.73 | | |
| HTATIP2 | 2.31 | MMP15 | 2.27 | Rho family GTPase | |
| IL8RB | 1.59 | MMP16 | 1.51 | ARHGDIA | 1.37 |
| SET | 1.39 | MMP2 | 1.55 | RAC2 | 1.39 |
| TSHR [†] | 5.46 | MMP7 | 1.40 | | |
| VEGFA | 1.45 | SPG7 | 1.84 | | |

Table 1: Up-regulated genes in mechanically stimulated cells obtained by PCR array analysis. (*genes tested in more than one array; [†]inconsistent results)

| Down-regulated genes (> -1.35 fold) | | | |
|-----------------------------------------------|------------------------|------------------------------------|------------------------|
| Gene name | Fold regulation | Gene name | Fold regulation |
| Cell adhesion genes | | Integrin mediated signaling | |
| ACTN3** | -3.89 | ITGB3* | -1.73 +/- 0.2 |
| CLEC3B | -2.17 | Proteases | |
| MYL9** | -1.57 | MMP8 | -2.13 |
| MTSS1** | -1.37 | Rho family GTPase signaling | |
| PTK2B** | -1.67 | RHO | -1.58 |

Table 2: Down-regulated genes in mechanically stimulated cells obtained by PCR array analysis. (*genes tested in more than one array; **genes tested individually by qPCR)

PTK2B (protein tyrosine kinase 2B), MYL9 (myosin light chain 9), ACTN3 (actinin 3), MMP14 (matrix metalloproteinase 14). Similar to PCR array analysis, qPCR was performed with RNA extracted from cells incubated under stimulated and non-stimulated conditions for 48 hours. For every gene, at least two biological replicates were used and for every biological replicate two technical replicates were performed. The housekeeping gene GAPDH was used for normalization of gene expression. After repeating the qPCR, CTTN, DIAPH1, SRC and TSHR did not show a consistent change in gene expression ($P > 0.05$). However, differential expression of ITGB3, MTSS1, PTK2B, MYL9, ACTN3 and MMP14 were confirmed by qPCR. Out of these genes, ITGB3, MTSS1, PTK2B, MYL9 and ACTN3 were confirmed to be down-regulated in response to





Figure 5. Genes confirmed to be differentially expressed upon mechanical stimulation.

ITGB3 – $\beta 3$ integrin; MTSS1 – Metastasis suppressor 1; PTK2B – Protein kinase 2 β ; MYL9 – Myosin light chain 9; ACTN3 – Alpha 3 actinin; MMP14 – Matrix metalloproteinase 14. Above differentially expressed genes were selected for further confirmation by qPCR. Two biological replicates were used for qPCR of MMP14; three biological replicates were used for qPCR of MTSS1, ITGB3, MYL9 and ACTN3; four biological replicates were used for qPCR of PTK2B. For every biological replicate two technical replicates were also performed. Statistical significance was tested by performing Student's t-test ($P < 0.05$).

mechanical stimulation, whereas, MMP14 was up-regulated after mechanical stimulation (Figure 5).

Down-regulation of the Integrin $\beta 3$ Protein upon Mechanical Stimulation

From the above confirmed differentially expressed genes, ITGB3 was chosen for further analysis for a number of reasons. ITGB3 codes for integrin $\beta 3$, one of the β subunits of membrane receptors called integrins. Integrins bind to extracellular membrane proteins and are known to relay signals from outside of the cell to inside and vice-versa and they are considered to be potential mechanosensors (Rathinam and Alahari, 2010). Integrin $\beta 3$ has been found to be differentially expressed in different types of cancer and this differential expression is associated with increased cell proliferation and invasion (Jin and Varner, 2004). More importantly, integrin $\beta 3$ pairs with integrin αv or integrin $\alpha 1b$ to form heterodimeric integrin molecules. These integrin $\beta 3$ containing heterodimers are known to bind Arginine-Glycine-Aspartic acid (RGD) domains of




fibronectin (Xiong et al., 2002). As fibronectin has been shown to be required for enhanced cellular response in our invasion assay (Menon and Beningo, 2011), the down-regulation of integrin $\beta 3$ upon mechanical stimulation was intriguing.

Given that we were sure $\beta 3$ levels were down-regulated at the transcriptional level, we confirmed the down-regulation of integrin $\beta 3$ at translational level. To test for $\beta 3$ protein levels, HT1080 cells were seeded on larger collagen/fibronectin substrates and incubated with and without mechanical stimulation for 48 hours. After 48 hours, cells were removed from within the matrix by collagenase treatment and protein was extracted from them using triple detergent lysis buffer. Western blot analysis performed using these cell lysates showed down-regulation of integrin $\beta 3$ protein levels when mechanically stimulated in the invasion assay (Figure 6A). Cell lysates from mechanically stimulated cells had approximately 0.6 times integrin $\beta 3$ compared to that from non-stimulated cells. This value was comparable to gene expression levels, where stimulated samples contained 0.5 times ITGB3 mRNA compared to non-stimulated samples (Figure 5 and 6B).

Enhanced Invasion is Inhibited by Over-expression of Integrin $\beta 3$

Once the down-regulation of integrin $\beta 3$ at transcriptional and translational levels was confirmed, we addressed the functional significance of this down-regulation. Naturally we speculated that its down-regulation is directly involved in sensing the mechanical stimulus. However, it is entirely possible that down-regulation of integrin $\beta 3$ is just an outcome of enhanced invasion and not required for sensing the mechanical stimulus. To test the functional significance of the down-regulation of integrin $\beta 3$ we over-expressed integrin $\beta 3$ in HT1080 cells and tested the invasion response of these cells in the mechano-invasion assay. If the down-regulation of integrin $\beta 3$ is required for cells to sense the mechanical stimulus and show enhanced invasion re-

sponse, overexpression of integrin $\beta 3$ should inhibit the enhanced invasion observed in these cells.




Integrin $\beta 3$ was overexpressed in HT1080 cells by nucleofecting them with pcDNA3.1-beta-3 plasmid. The cells were also co-nucleofected with GFP plasmid to estimate the nucleofection efficiency. Fluorescence microscopy, indicated a 75-80% nucleofection efficiency was obtained for every experiment. HT1080 cells nucleofected with only GFP plasmid were used as control for these experiments. Western blot confirmed that ITGB3 gene was stably overexpressed up to 72 hours and approximately two-fold over-expression was obtained (Figure 7A). After nucleofection cells were incubated at 37°C for 24 hours to let them recover and allow stable over-expression of integrin $\beta 3$. After 24 hours these cells were seeded on collagen/fibronectin substrates and incubated with or without mechanical stimulation for 48 hours. After 48 hours, the number of cells invaded into the collagen/fibronectin matrix, were counted. It was observed that cells overexpressing integrin $\beta 3$ failed to respond to mechanical stimulation and did not display enhanced invasion. However, control cells nucleofected with GFP plasmid showed the typical 2-fold enhanced invasion with mechanical stimulation compared to those under non-stimulated conditions (Figure 7B). This confirmed that down-regulation of integrin $\beta 3$ is required for the process of enhanced invasion and if the integrin $\beta 3$ down-regulation is inhibited, cells fail to show enhanced invasion in response to mechanical stimulation.

Down-regulation of Integrin $\beta 3$ is Not Accompanied by Increased Activity of Integrin $\beta 1$

Numerous instances of integrin crosstalk have been reported, where binding of one integrin to its ligand causes perturbation in the expression levels or activity of other integrins. Such crosstalk is known to occur between integrin $\beta 3$ and integrin $\beta 1$ (Gonzalez et al., 2010). Integrin $\beta 1$ is another fibronectin binding β subunit of integrin. To address the potential of integrin cross-talk between these two integrins we tested if down-regulation of integrin $\beta 3$ is accompanied by an increase in the activation of integrin $\beta 1$. A re-evaluation of the PCR array analysis indicated there was no difference in the expression levels of integrin $\beta 1$ under stimulated and non-

stimulated conditions (Figure 8A). However, as a result of integrin crosstalk between integrin $\beta 1$ and $\beta 3$ the regulation may not occur at the level of expression but in the degree of receptor activation of $\beta 1$. To address this possibility, we tested if levels of active integrin $\beta 1$ are increased upon mechanical stimulation compared to non-stimulated cultures.




We used an antibody that specifically recognizes the activated conformation of integrin $\beta 1$ and probed western blots prepared from HT1080 cell lysates. HT1080 cells incubated with or without mechanical stimulation for 48 hours, were separated from collagen/fibronectin matrix and lysed to collect total protein. Western blot performed with these protein extracts determined that levels of activated integrin $\beta 1$ upon mechanical stimulation were not significantly different from those in non-stimulated cells (Figure 8B and 8C). Our results suggest that no crosstalk occurs between integrin $\beta 1$ and integrin $\beta 3$ in the invasive response observed upon mechanical stimulation.

Down-regulation of Integrin $\beta 3$ causes Decrease in Levels of Ser3 Phospho-cofilin upon Mechanical Stimulation

To identify potential down-stream pathways of integrin $\beta 3$, we focused on the relationship between integrins and cofilin. As mentioned previously, cofilin is an actin binding protein that severs actin filaments and have been shown to be required for invadopodia maturation (Yamaguchi et al., 2005). Moreover, in our *in vitro* mechano-invasion assay, cofilin siRNA treated cells failed to sense the mechanical stimulation and did not show enhanced invasion, yet retained basal levels of invasion (Menon and Beningo, 2011). Cofilin activity within the cell is controlled by its phosphorylation at serine-3 by LIM kinase. When phosphorylated at serine-3, cofilin loses its ability to bind and sever actin filaments and is considered to be inactive. The phosphorylation status of cofilin is under indirect control of Rho GTPases (Pollard and Borisy, 2003). As Rho GTPases such as Rho, Rac and Cdc42 are differentially regulated under different integrins, differences in integrin signaling might also affect down-stream cofilin activity. We hypothesized that under mechanically stimulated conditions levels of Ser3 phospho-cofilin (inactive) were lower compared to those in non-stimulated conditions, leading to enhanced invasion. In addition, if the

cofilin activity is dependent on integrin $\beta 3$ expression, overexpression of integrin $\beta 3$ should increase the levels of phospho-cofilin upon mechanical stimulation.



To test our hypothesis, cell lysates were collected from HT1080 cells cultured on collagen/fibronectin substrates and incubated with or without mechanical stimulation for 48 hours. The western blot performed with these cell lysates determined that levels of Ser3 phospho-cofilin were indeed decreased under mechanically stimulated conditions compared to non-stimulated conditions indicating that more cofilin was likely to be active (Figure 9A). Cell lysates collected from mock nucleofected cells also showed similar decrease in Ser3 phospho-cofilin upon mechanical stimulation. Quantitation of band intensities found that Ser3 phospho-cofilin upon mechanical stimulation were approximately 0.65 times lower when compared to the non-stimulated samples in both HT1080 cells and mock nucleofected HT1080 cells (Figure 9B). However, when integrin $\beta 3$ was over-expressed within the cells, the effect of mechanical stimulation on Ser3 phospho-cofilin level was reversed, such that similar levels of Ser3 phospho-cofilin were observed in integrin $\beta 3$ overexpressing HT1080 cells with or without mechanical stimulation (Figure 9A and 9B). In addition, the amount of total cofilin was unchanged in HT1080 cells, integrin $\beta 3$ overexpressing HT1080 cells and mock nucleofected HT1080 cells, between stimulated and non-stimulated conditions (Figure 9A). Our data strongly suggests that the amount of active cofilin increases in response to mechanical stimulation and this increase in active cofilin is brought about by down-regulation of integrin $\beta 3$.

CHAPTER 4 - DISCUSSION

Metastasis is a multistep process controlled by various biochemical and mechanical factors in the tumor microenvironment. Secondary tumors formed by metastatic cancer cells are the leading cause of death in cancer patients. In order to recognize potential therapeutic targets against metastasis, it is important to understand the effects of factors that instigate the process in the first place. In this study, I have focused on the effects of mechanical factors in the tumor microenvironment that can enhance cancer cell invasion. In our *in vitro* mechano-invasion assay we used magnetic beads embedded within the collagen/fibronectin matrix to mimic the pulling forces produced by cellular movements within the extracellular matrix (Menon and Beningo, 2011). Under the effect of these pulling forces, a greater number of cancer cells will invade into the ECM (Figure 2). In this study I wanted to identify molecular players in the process of mechanosensing that eventually leads to the enhanced invasion of cancer cells in our assay. My approach was to identify genes that are differentially expressed between mechanically stimulated and non-stimulated conditions by qPCR. As mRNA is the least stable biomolecule within the cell, it was challenging to standardize a method for RNA extraction from invaded cells. With the help of collagenase, invaded cells were taken out of the matrix within 15 min and RNA was successfully extracted from them.

Using PCR arrays I was able to screen over 200 genes that could be differentially regulated between mechanically stimulated and non-stimulated conditions. Candidate genes were identified and categorized based on their function. These genes belonged to the following categories; cell adhesion, cell growth and proliferation, membrane protrusions, proteases, ECM proteins, protease inhibitors, Rho family GTPase signaling (Figure 4). The pattern of overall gene regulation in some of the above categories was interesting. For instance, in the case of cell adhesion genes, a greater number of genes were down-regulated than up-regulated. This

observation is consistent with the idea that in both 3D invasion and 2D migration, optimal motility of cells occurs when cells possess an intermediate degree of adhesiveness (Palecek et al., 1997; Zaman et al., 2006). Therefore, down-regulation of cell adhesion genes is suggestive of an increase in invasive ability of the cells. Also, in my screen, many genes involved in membrane protrusions were found to be up-regulated. As these genes are known to be required for invadopodia formation in tumor cells, their up-regulation could indicate formation of more invadopodia under mechanical stimulation (Albiges-Rizo et al., 2009; Alexander et al., 2008). In addition, I also found that many genes coding for different proteases were up-regulated under mechanical stimulation. During invasion, cells need to breach the physical barriers formed by the basement membrane (BM) and stroma, and they secrete proteases to degrade the ECM proteins that form these barriers (Poincloux et al., 2009). Hence, this increase in proteases also suggests increased invasive activity by cells. In addition to these, differential regulation of Rho GTPase signaling molecules, which are the key regulators of actin dynamics, also point to modulation of cell motility upon stimulation (Jaffe and Hall, 2005).

From the above candidate genes I was able to confirm the differential expression of six genes between mechanically stimulated and non-stimulated conditions of our mechano-invasion assay. Out of these six genes, ITGB3 that codes for integrin β 3 was selected for further studies because it was the only integrin subunit found to be differentially regulated upon mechanical stimulation. Other genes confirmed to be differentially expressed were as follows. 1) MTSS1, also known as 'Missing-in-metastasis (MIM)', is expressed in low levels in cancer cells (Lee et al., 2002). It codes for an actin binding scaffold protein that regulates actin dynamics through its interactions with cortactin and protein tyrosine phosphatase δ (PTP δ) (Gonzalez-Quevedo et al., 2005; Lin et al., 2005; Quinones et al., 2010; Woodings et al., 2003). 2) MYL9 codes for myosin light chain (MLC) which is a regulatory component whose phosphorylation state dictates myosin

motor binding to actin filaments. The actomyosin contractility is required by cells to apply traction force on its substrate as well as for retracting the trailing edge of the cell as it moves forward hence it is mechanistically important in cell motility. (Jaffe and Hall, 2005; Ridley et al., 2003). 3) ACTN3 codes for a homologue of α -actinin. It is an actin crosslinking protein generally found at different cell-cell and cell-matrix adhesions. It connects transmembrane proteins such as tails of β integrins and intracellular adhesion molecule-1 (ICAM-1) to the actin cytoskeleton (Carpen et al., 1992; Otey et al., 1990). It also acts as a scaffold to bring together different signaling molecules and plays an important role in cell migration (Otey and Carpen, 2004). 4) PTK2B codes for a non-receptor protein tyrosine kinase that binds to cytoplasmic tails of integrins and is activated in response to cues from fibronectin in many developmental processes (Ma et al 1997, Schaller and Sasaki, 1997). Down-regulation of PTK2B is required for the differential cell response to rigidity and compliance in human foreskin fibroblasts (Prager-Khoutorsky et al., 2011). The kinase activity of PTK2B has been correlated with increased malignancy of cancer cells and the catalytic domain of PTK2B has become a therapeutic target (Lipinski and Loftus, 2010). 5) MMP14 encodes a zinc-dependent endopeptidase called matrix metalloproteinase 14 that causes collagenolysis (Li et al., 2008; Sabeh et al., 2004). In tumor cells, MMP14 is enriched at invadopodia. This provides invadopodial structures with an ECM degrading property, thereby increasing the rate of invasion of tumor cells through the ECM (Clark and Weaver, 2008; Linder, 2007).

As described earlier, being transmembrane proteins, β subunits of integrin are able to bind extracellular ligands outside the cell and actin cytoskeleton and a variety of signaling proteins inside the cell (Rathinam and Alahari, 2010; Springer and Wang, 2004). By virtue of this property, β subunits serve as potential mechanosensors and mechanotransducers. Using qPCR and western blot analysis I confirmed that integrin β 3 is down-regulated upon mechanical stimulation

at both the transcriptional and translational levels. As it is a candidate mechanosensing molecule, one would expect to see up-regulation of integrin $\beta 3$ expression upon mechanical stimulation, hence this decrease in integrin $\beta 3$ expression was rather surprising. To confirm that the down-regulation of integrin $\beta 3$ upon mechanical stimulation is required for enhanced invasion of cells, I overexpressed integrin $\beta 3$ and inhibited the enhanced invasion normally seen upon mechanical stimulation. This indicated that integrin $\beta 3$ must be down-regulated upon mechanical stimulation in order for the cells to respond to the stimulus. The upstream mechanism of how the mechanical stimulation causes the down-regulation of integrin $\beta 3$ is unclear. Since fibronectin in the ECM is required for sensing this mechanical stimulus, it is entirely possible that down-regulation of integrin $\beta 3$ is caused by a feedback mechanism that is initiated by mechanosensing through integrin $\beta 3$ itself. Alternatively, a second integrin could be responsible for the down-regulation I have observed.

Different signaling pathways are controlled down-stream of different integrins and can result in very different cellular responses. To control the display and engagement of the correct integrins at the proper time, cells use a mechanism of integrin crosstalk, where one type of integrin regulates expression and activity of another type of integrin (Gonzalez et al., 2010). Such integrin crosstalk has been observed between integrin $\beta 3$ and another fibronectin binding integrin, integrin $\beta 1$. As integrin $\beta 3$ is down-regulated upon mechanical stimulation, I checked if it was accompanied by an increase in activity of integrin $\beta 1$. Unfortunately, we did not observe any significant changes in the levels of activation of integrin $\beta 1$ upon mechanical stimulation. However, another possibility is that down-regulation of integrin $\beta 3$ might change localization of integrin $\beta 1$ within the cell. For example, down-regulation of integrin $\beta 3$ might enrich integrin $\alpha 5\beta 1$ in invadopodia and thus increasing their interaction with fibronectin resulting in enhanced mechanosensing.

The downstream pathway resulting from the down-regulation of integrin $\beta 3$, and eventual increase in cellular invasion upon mechanical stimulation, may not be as obscure as its upstream pathway. To begin, our previous studies have shown the requirement of cofilin for sensing the mechanical stimulation in our mechano-invasion assay (Menon and Beningo, 2011). As mentioned previously cofilin is an actin binding protein, also required for maturation of invadopodia and its activity is dependent on phosphorylation by LIM kinase (Pollard and Borisy, 2003; Yamaguchi et al., 2005). (Blanchoin et al., 2000). My results suggest cofilin is in its active state (low phosphorylation at Ser-9) upon mechanical stimulation and that this activation can be inhibited upon overexpression of integrin $\beta 3$. This evidence provides a clue to a known pathway leading from integrin beta 3 to cofilin and potentially to the maturation of invadopodia.

As cofilin is a key regulator of the actin cytoskeleton, the balance between its active and inactive forms is very important with respect to cell motility and invasion. The levels of Ser3 phosphorylated cofilin are low in highly metastatic cell lines derived from T-lymphoma and carcinoma (Nebl et al., 1996). In *in vitro* as well as *in vivo* experiments, invasiveness of cancer cells has been correlated with overall output of the cofilin pathway (Wang et al., 2006; Zebda et al., 2000). The mammary tumors formed by LIMK1 over-expressing cancer cells have been shown to have low metastatic potential. The tumors with dominant negative LIMK1 expressing cells, on the other hand, show increased invasive abilities (Wang et al., 2006). The phosphorylation of cofilin is linked to the ligand binding ability of integrin $\alpha v \beta 3$. Murine melanoma cells expressing integrin $\alpha v \beta 3$ have 10-fold higher amounts of Ser3 phospho-cofilin than integrin $\alpha v \beta 3$ negative cells. Moreover, expression of pseudophosphorylated and non-phosphorylatable mutant cofilin in these cells showed that cofilin activity is required for expression of MMP2 and MMP14 (Dang et al., 2006).

Based on our observations and current literature we have developed a model that could explain the pathway connecting integrin $\beta 3$ and cofilin. It is known that integrin engagement and activation leads to activation Rho GTPases. Activity of Rho GTPase Rac1 can be regulated by integrin $\beta 3$ in focal complexes (Morgan et al., 2009). We hypothesize that in conditions without mechanical stimulation, signaling down-stream of integrin $\beta 3$ leads to activation of Rac1. The active GTP bound Rac1 phosphorylates and activates its down-stream effector p21 activated kinase-1 (PAK1) (del Pozo et al., 2000). Active PAK1 activates LIM kinase-1 (LIMK1), which in turn phosphorylates its substrate cofilin at Ser-3 thereby inactivating it. Therefore, under non-




Figure 10. Possible pathway for enhanced invasion upon mechanical stimulation. Signaling down-stream of integrin $\beta 3$, through sequential activation of Rac1, PAK1 and LIMK1, results in inhibition of cofilin activity by phosphorylation at Ser3. Upon mechanical stimulation integrin $\beta 3$ gets down-regulated thereby decreasing the levels of down-stream kinases. Decrease in activation of LIMK1, also decreases phosphorylation of cofilin. This leads to increase in levels of active cofilin. Increased cofilin activity probably through formation of more mature invadopodia causes enhanced invasion.

stimulated conditions, the amount of active cofilin is lower than inactive cofilin, yet it is enough to produce basal levels of invasion. When mechanical stimulation is provided, integrin $\beta 3$ is down-regulated through an unknown mechanism. As the amount of integrin $\beta 3$ displayed on the surface decreases, the overall signaling down-stream of it also decreases. This leads to a decrease in the amounts of Rac1-GTP, which further causes less activation of PAK1 and LIMK1. Thus, the collective effect of integrin $\beta 3$ down-regulation is a decrease in the amount of inactive Ser3 phospho-cofilin and an increase in non-phosphorylated active form. This increase in active cofilin upon stimulation leads to the formation of more mature invadopodia which eventually cause enhanced invasion.

In conclusion, this study has uncovered a part of the pathway that leads to enhanced invasion of cancer cells in response to tugging forces in the tumor microenvironment. For future studies, we first need to confirm the involvement of Rac1, PAK1 and LIMK1 down-stream of integrin $\beta 3$ for controlling activation status of cofilin, as predicted by the model. As cofilin is known to be involved in invadopodia maturation, it will be interesting to test by microscopic studies if stimulated cultures show increased invadopodial activity. Moreover, how cofilin controls the maturation of invadopodia is still unknown. Solving these still unanswered questions will take us one step closer to understanding the mechanosensing pathway causing enhanced cancer cell invasion.

REFERENCES

- Albelda, S.M., S.A. Mette, D.E. Elder, R. Stewart, L. Damjanovich, M. Herlyn, and C.A. Buck. 1990. Integrin distribution in malignant melanoma: association of the beta 3 subunit with tumor progression. *Cancer research*. 50:6757-6764.
- Albiges-Rizo, C., O. Destaing, B. Fourcade, E. Planus, and M.R. Block. 2009. Actin machinery and mechanosensitivity in invadopodia, podosomes and focal adhesions. *Journal of cell science*. 122:3037-3049.
- Alexander, N.R., K.M. Branch, A. Parekh, E.S. Clark, I.C. Iwueke, S.A. Guelcher, and A.M. Weaver. 2008. Extracellular matrix rigidity promotes invadopodia activity. *Current biology : CB*. 18:1295-1299.
- Ao, M., O.E. Franco, D. Park, D. Raman, K. Williams, and S.W. Hayward. 2007. Cross-talk between paracrine-acting cytokine and chemokine pathways promotes malignancy in benign human prostatic epithelium. *Cancer research*. 67:4244-4253.
- Beningo, K.A., and Y.L. Wang. 2002. Flexible substrata for the detection of cellular traction forces. *Trends in cell biology*. 12:79-84.
- Bissell, M.J., and D. Radisky. 2001. Putting tumours in context. *Nature reviews. Cancer*. 1:46-54.
- Blanchoin, L., T.D. Pollard, and R.D. Mullins. 2000. Interactions of ADF/cofilin, Arp2/3 complex, capping protein and profilin in remodeling of branched actin filament networks. *Current biology : CB*. 10:1273-1282.
- Blouw, B., D.F. Seals, I. Pass, B. Diaz, and S.A. Courtneidge. 2008. A role for the podosome/invadopodia scaffold protein Tks5 in tumor growth in vivo. *European journal of cell biology*. 87:555-567.
- Borghi, N., M. Lowndes, V. Maruthamuthu, M.L. Gardel, and W.J. Nelson. 2010. Regulation of cell motile behavior by crosstalk between cadherin- and integrin-mediated adhesions.

Proceedings of the National Academy of Sciences of the United States of America.
107:13324-13329.

Boucher, Y., and R.K. Jain. 1992. Microvascular pressure is the principal driving force for interstitial hypertension in solid tumors: implications for vascular collapse. *Cancer research.* 52:5110-5114.

Boyd, N.F., G.A. Lockwood, L.J. Martin, J.A. Knight, R.A. Jong, E. Fishell, J.W. Byng, M.J. Yaffe, and D.L. Tritchler. 1999. Mammographic densities and risk of breast cancer among subjects with a family history of this disease. *Journal of the National Cancer Institute.* 91:1404-1408.

Brooks, P.C., A.M. Montgomery, M. Rosenfeld, R.A. Reisfeld, T. Hu, G. Klier, and D.A. Cheresh. 1994. Integrin alpha v beta 3 antagonists promote tumor regression by inducing apoptosis of angiogenic blood vessels. *Cell.* 79:1157-1164.

Butcher, D.T., T. Alliston, and V.M. Weaver. 2009. A tense situation: forcing tumour progression. *Nature reviews. Cancer.* 9:108-122.

Carpen, O., P. Pallai, D.E. Staunton, and T.A. Springer. 1992. Association of intercellular adhesion molecule-1 (ICAM-1) with actin-containing cytoskeleton and alpha-actinin. *The Journal of cell biology.* 118:1223-1234.

Castella, L.F., L. Buscemi, C. Godbout, J.J. Meister, and B. Hinz. 2010. A new lock-step mechanism of matrix remodelling based on subcellular contractile events. *Journal of cell science.* 123:1751-1760.

Cat, B., D. Stuhlmann, H. Steinbrenner, L. Alili, O. Holtkotter, H. Sies, and P. Brenneisen. 2006. Enhancement of tumor invasion depends on transdifferentiation of skin fibroblasts mediated by reactive oxygen species. *Journal of cell science.* 119:2727-2738.

- Chambers, A.F., A.C. Groom, and I.C. MacDonald. 2002. Dissemination and growth of cancer cells in metastatic sites. *Nature reviews. Cancer.* 2:563-572.
- Clark, E.S., B. Brown, A.S. Whigham, A. Kochaishvili, W.G. Yarbrough, and A.M. Weaver. 2009. Aggressiveness of HNSCC tumors depends on expression levels of cortactin, a gene in the 11q13 amplicon. *Oncogene.* 28:431-444.
- Clark, E.S., and A.M. Weaver. 2008. A new role for cortactin in invadopodia: regulation of protease secretion. *European journal of cell biology.* 87:581-590.
- Danen, E.H., J. van Rheenen, W. Franken, S. Huveneers, P. Sonneveld, K. Jalink, and A. Sonnenberg. 2005. Integrins control motile strategy through a Rho-cofilin pathway. *The Journal of cell biology.* 169:515-526.
- Dang, D., J.R. Bamburg, and D.M. Ramos. 2006. Alphavbeta3 integrin and cofilin modulate K1735 melanoma cell invasion. *Experimental cell research.* 312:468-477.
- Darby, I., O. Skalli, and G. Gabbiani. 1990. Alpha-smooth muscle actin is transiently expressed by myofibroblasts during experimental wound healing. *Laboratory investigation; a journal of technical methods and pathology.* 63:21-29.
- De Wever, O., P. Demetter, M. Mareel, and M. Bracke. 2008. Stromal myofibroblasts are drivers of invasive cancer growth. *International journal of cancer. Journal international du cancer.* 123:2229-2238.
- del Pozo, M.A., L.S. Price, N.B. Alderson, X.D. Ren, and M.A. Schwartz. 2000. Adhesion to the extracellular matrix regulates the coupling of the small GTPase Rac to its effector PAK. *The EMBO journal.* 19:2008-2014.
- Denys, H., L. Derycke, A. Hendrix, W. Westbroek, A. Gheldof, K. Narine, P. Pauwels, C. Gespach, M. Bracke, and O. De Wever. 2008. Differential impact of TGF-beta and EGF on fibroblast

- differentiation and invasion reciprocally promotes colon cancer cell invasion. *Cancer letters*. 266:263-274.
- Desmouliere, A., A. Geinoz, F. Gabbiani, and G. Gabbiani. 1993. Transforming growth factor-beta 1 induces alpha-smooth muscle actin expression in granulation tissue myofibroblasts and in quiescent and growing cultured fibroblasts. *The Journal of cell biology*. 122:103-111.
- Desmouliere, A., C. Guyot, and G. Gabbiani. 2004. The stroma reaction myofibroblast: a key player in the control of tumor cell behavior. *The International journal of developmental biology*. 48:509-517.
- Diaz-Gonzalez, F., J. Forsyth, B. Steiner, and M.H. Ginsberg. 1996. Trans-dominant inhibition of integrin function. *Molecular biology of the cell*. 7:1939-1951.
- Dong, J., J. Grunstein, M. Tejada, F. Peale, G. Frantz, W.C. Liang, W. Bai, L. Yu, J. Kowalski, X. Liang, G. Fuh, H.P. Gerber, and N. Ferrara. 2004. VEGF-null cells require PDGFR alpha signaling-mediated stromal fibroblast recruitment for tumorigenesis. *The EMBO journal*. 23:2800-2810.
- Edwards, D.C., L.C. Sanders, G.M. Bokoch, and G.N. Gill. 1999. Activation of LIM-kinase by Pak1 couples Rac/Cdc42 GTPase signalling to actin cytoskeletal dynamics. *Nature cell biology*. 1:253-259.
- Eliceiri, B.P., and D.A. Cheresh. 1999. The role of alphav integrins during angiogenesis: insights into potential mechanisms of action and clinical development. *The Journal of clinical investigation*. 103:1227-1230.
- Eskens, F.A., H. Dumez, R. Hoekstra, A. Perschl, C. Brindley, S. Bottcher, W. Wynendaele, J. Drevs, J. Verweij, and A.T. van Oosterom. 2003. Phase I and pharmacokinetic study of continuous twice weekly intravenous administration of Cilengitide (EMD 121974), a novel

- inhibitor of the integrins $\alpha v \beta 3$ and $\alpha v \beta 5$ in patients with advanced solid tumours. *European journal of cancer*. 39:917-926.
- Fukumura, D., D.G. Duda, L.L. Munn, and R.K. Jain. 2010. Tumor microvasculature and microenvironment: novel insights through intravital imaging in pre-clinical models. *Microcirculation*. 17:206-225.
- Gabbiani, G., G.B. Ryan, and G. Majne. 1971. Presence of modified fibroblasts in granulation tissue and their possible role in wound contraction. *Experientia*. 27:549-550.
- Gang, Z., Q. Qi, C. Jing, and C. Wang. 2009. Measuring microenvironment mechanical stress of rat liver during diethylnitrosamine induced hepatocarcinogenesis by atomic force microscope. *Microscopy research and technique*. 72:672-678.
- Gasparini, G., P.C. Brooks, E. Biganzoli, P.B. Vermeulen, E. Bonoldi, L.Y. Dirix, G. Ranieri, R. Miceli, and D.A. Cheresh. 1998. Vascular integrin $\alpha(v)\beta 3$: a new prognostic indicator in breast cancer. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 4:2625-2634.
- Gonzalez-Quevedo, R., M. Shoffer, L. Horng, and A.E. Oro. 2005. Receptor tyrosine phosphatase-dependent cytoskeletal remodeling by the hedgehog-responsive gene MIM/BEG4. *The Journal of cell biology*. 168:453-463.
- Gonzalez, A.M., R. Bhattacharya, G.W. deHart, and J.C. Jones. 2010. Transdominant regulation of integrin function: mechanisms of crosstalk. *Cellular signalling*. 22:578-583.
- Guo, W., and F.G. Giancotti. 2004. Integrin signalling during tumour progression. *Nature reviews. Molecular cell biology*. 5:816-826.
- Hagendoorn, J., R. Tong, D. Fukumura, Q. Lin, J. Lobo, T.P. Padera, L. Xu, R. Kucherlapati, and R.K. Jain. 2006. Onset of abnormal blood and lymphatic vessel function and interstitial hypertension in early stages of carcinogenesis. *Cancer research*. 66:3360-3364.

- Helmlinger, G., P.A. Netti, H.C. Lichtenbeld, R.J. Melder, and R.K. Jain. 1997. Solid stress inhibits the growth of multicellular tumor spheroids. *Nature biotechnology*. 15:778-783.
- Hinz, B., G. Celetta, J.J. Tomasek, G. Gabbiani, and C. Chaponnier. 2001a. Alpha-smooth muscle actin expression upregulates fibroblast contractile activity. *Molecular biology of the cell*. 12:2730-2741.
- Hinz, B., V. Dugina, C. Ballestrem, B. Wehrle-Haller, and C. Chaponnier. 2003. Alpha-smooth muscle actin is crucial for focal adhesion maturation in myofibroblasts. *Molecular biology of the cell*. 14:2508-2519.
- Hinz, B., D. Mastrangelo, C.E. Iselin, C. Chaponnier, and G. Gabbiani. 2001b. Mechanical tension controls granulation tissue contractile activity and myofibroblast differentiation. *The American journal of pathology*. 159:1009-1020.
- Hirakawa, S., S. Kodama, R. Kunstfeld, K. Kajiya, L.F. Brown, and M. Detmar. 2005. VEGF-A induces tumor and sentinel lymph node lymphangiogenesis and promotes lymphatic metastasis. *The Journal of experimental medicine*. 201:1089-1099.
- Hoyt, K., B. Castaneda, M. Zhang, P. Nigwekar, P.A. di Sant'agnese, J.V. Joseph, J. Strang, D.J. Rubens, and K.J. Parker. 2008. Tissue elasticity properties as biomarkers for prostate cancer. *Cancer biomarkers : section A of Disease markers*. 4:213-225.
- Huang, S., L. Mills, B. Mian, C. Tellez, M. McCarty, X.D. Yang, J.M. Gudas, and M. Bar-Eli. 2002. Fully humanized neutralizing antibodies to interleukin-8 (ABX-IL8) inhibit angiogenesis, tumor growth, and metastasis of human melanoma. *The American journal of pathology*. 161:125-134.
- Hynes, R.O. 2002. Integrins: bidirectional, allosteric signaling machines. *Cell*. 110:673-687.
- Ingber, D.E. 2008. Tensegrity-based mechanosensing from macro to micro. *Progress in biophysics and molecular biology*. 97:163-179.

- Jaffe, A.B., and A. Hall. 2005. Rho GTPases: biochemistry and biology. *Annual review of cell and developmental biology*. 21:247-269.
- Jin, H., and J. Varnier. 2004. Integrins: roles in cancer development and as treatment targets. *British journal of cancer*. 90:561-565.
- Katsumi, A., A.W. Orr, E. Tzima, and M.A. Schwartz. 2004. Integrins in mechanotransduction. *The Journal of biological chemistry*. 279:12001-12004.
- Kim, J.H., L.T. Zheng, W.H. Lee, and K. Suk. 2011. Pro-apoptotic role of integrin beta3 in glioma cells. *Journal of neurochemistry*. 117:494-503.
- Kostic, A., C.D. Lynch, and M.P. Sheetz. 2009. Differential matrix rigidity response in breast cancer cell lines correlates with the tissue tropism. *PLoS one*. 4:e6361.
- Kumar, S., and V.M. Weaver. 2009. Mechanics, malignancy, and metastasis: the force journey of a tumor cell. *Cancer metastasis reviews*. 28:113-127.
- Lederle, W., H.J. Stark, M. Skobe, N.E. Fusenig, and M.M. Mueller. 2006. Platelet-derived growth factor-BB controls epithelial tumor phenotype by differential growth factor regulation in stromal cells. *The American journal of pathology*. 169:1767-1783.
- Lee, Y.G., J.A. Macoska, S. Korenchuk, and K.J. Pienta. 2002. MIM, a potential metastasis suppressor gene in bladder cancer. *Neoplasia*. 4:291-294.
- Levental, K.R., H. Yu, L. Kass, J.N. Lakins, M. Egeblad, J.T. Ertler, S.F. Fong, K. Csiszar, A. Giaccia, W. Weninger, M. Yamauchi, D.L. Gasser, and V.M. Weaver. 2009. Matrix crosslinking forces tumor progression by enhancing integrin signaling. *Cell*. 139:891-906.
- Li, X.Y., I. Ota, I. Yana, F. Sabeh, and S.J. Weiss. 2008. Molecular dissection of the structural machinery underlying the tissue-invasive activity of membrane type-1 matrix metalloproteinase. *Molecular biology of the cell*. 19:3221-3233.

- Lin, J., J. Liu, Y. Wang, J. Zhu, K. Zhou, N. Smith, and X. Zhan. 2005. Differential regulation of cortactin and N-WASP-mediated actin polymerization by missing in metastasis (MIM) protein. *Oncogene*. 24:2059-2066.
- Linder, S. 2007. The matrix corroded: podosomes and invadopodia in extracellular matrix degradation. *Trends in cell biology*. 17:107-117.
- Liotta, L.A., C.N. Rao, and U.M. Wewer. 1986. Biochemical interactions of tumor cells with the basement membrane. *Annual review of biochemistry*. 55:1037-1057.
- Lipinski, C.A., and J.C. Loftus. 2010. Targeting Pyk2 for therapeutic intervention. *Expert opinion on therapeutic targets*. 14:95-108.
- Maciver, S.K., and P.J. Hussey. 2002. The ADF/cofilin family: actin-remodeling proteins. *Genome biology*. 3:reviews3007.
- Makino, A., E.R. Prossnitz, M. Bunemann, J.M. Wang, W. Yao, and G.W. Schmid-Schonbein. 2006. G protein-coupled receptors serve as mechanosensors for fluid shear stress in neutrophils. *American journal of physiology. Cell physiology*. 290:C1633-1639.
- Menon, S., and K.A. Beningo. 2011. Cancer cell invasion is enhanced by applied mechanical stimulation. *PloS one*. 6:e17277.
- Mizejewski, G.J. 1999. Role of integrins in cancer: survey of expression patterns. *Proceedings of the Society for Experimental Biology and Medicine. Society for Experimental Biology and Medicine*. 222:124-138.
- Moinfar, F., Y.G. Man, L. Arnould, G.L. Bratthauer, M. Ratschek, and F.A. Tavassoli. 2000. Concurrent and independent genetic alterations in the stromal and epithelial cells of mammary carcinoma: implications for tumorigenesis. *Cancer research*. 60:2562-2566.

- Morgan, M.R., A. Byron, M.J. Humphries, and M.D. Bass. 2009. Giving off mixed signals--distinct functions of alpha5beta1 and alphavbeta3 integrins in regulating cell behaviour. *IUBMB life*. 61:731-738.
- Nebi, G., S.C. Meuer, and Y. Samstag. 1996. Dephosphorylation of serine 3 regulates nuclear translocation of cofilin. *The Journal of biological chemistry*. 271:26276-26280.
- Niwa, R., K. Nagata-Ohashi, M. Takeichi, K. Mizuno, and T. Uemura. 2002. Control of actin reorganization by Slingshot, a family of phosphatases that dephosphorylate ADF/cofilin. *Cell*. 108:233-246.
- Oft, M., K.H. Heider, and H. Beug. 1998. TGFbeta signaling is necessary for carcinoma cell invasiveness and metastasis. *Current biology : CB*. 8:1243-1252.
- Orr, A.W., B.P. Helmke, B.R. Blackman, and M.A. Schwartz. 2006. Mechanisms of mechanotransduction. *Developmental cell*. 10:11-20.
- Otey, C.A., and O. Carpen. 2004. Alpha-actinin revisited: a fresh look at an old player. *Cell motility and the cytoskeleton*. 58:104-111.
- Otey, C.A., F.M. Pavalko, and K. Burridge. 1990. An interaction between alpha-actinin and the beta 1 integrin subunit in vitro. *The Journal of cell biology*. 111:721-729.
- Palecek, S.P., J.C. Loftus, M.H. Ginsberg, D.A. Lauffenburger, and A.F. Horwitz. 1997. Integrin-ligand binding properties govern cell migration speed through cell-substratum adhesiveness. *Nature*. 385:537-540.
- Pantel, K., and R.H. Brakenhoff. 2004. Dissecting the metastatic cascade. *Nature reviews. Cancer*. 4:448-456.
- Parekh, A., and A.M. Weaver. 2009. Regulation of cancer invasiveness by the physical extracellular matrix environment. *Cell adhesion & migration*. 3:288-292.

- Paszek, M.J., N. Zahir, K.R. Johnson, J.N. Lakins, G.I. Rozenberg, A. Gefen, C.A. Reinhart-King, S.S. Margulies, M. Dembo, D. Boettiger, D.A. Hammer, and V.M. Weaver. 2005. Tensional homeostasis and the malignant phenotype. *Cancer cell*. 8:241-254.
- Plow, E.F., T.A. Haas, L. Zhang, J. Loftus, and J.W. Smith. 2000. Ligand binding to integrins. *The Journal of biological chemistry*. 275:21785-21788.
- Poincloux, R., F. Lizarraga, and P. Chavrier. 2009. Matrix invasion by tumour cells: a focus on MT1-MMP trafficking to invadopodia. *Journal of cell science*. 122:3015-3024.
- Pollard, J.W. 2004. Tumour-educated macrophages promote tumour progression and metastasis. *Nature reviews. Cancer*. 4:71-78.
- Pollard, T.D., and G.G. Borisy. 2003. Cellular motility driven by assembly and disassembly of actin filaments. *Cell*. 112:453-465.
- Prager-Khoutorsky, M., A. Lichtenstein, R. Krishnan, K. Rajendran, A. Mayo, Z. Kam, B. Geiger, and A.D. Bershadsky. 2011. Fibroblast polarization is a matrix-rigidity-dependent process controlled by focal adhesion mechanosensing. *Nature cell biology*. 13:1457-1465.
- Quinones, G.A., J. Jin, and A.E. Oro. 2010. I-BAR protein antagonism of endocytosis mediates directional sensing during guided cell migration. *The Journal of cell biology*. 189:353-367.
- Rathinam, R., and S.K. Alahari. 2010. Important role of integrins in the cancer biology. *Cancer metastasis reviews*. 29:223-237.
- Retta, S.F., G. Cassara, M. D'Amato, R. Alessandro, M. Pellegrino, S. Degani, G. De Leo, L. Silengo, and G. Tarone. 2001. Cross talk between beta(1) and alpha(V) integrins: beta(1) affects beta(3) mRNA stability. *Molecular biology of the cell*. 12:3126-3138.
- Reynolds, L.E., L. Wyder, J.C. Lively, D. Taverna, S.D. Robinson, X. Huang, D. Sheppard, R.O. Hynes, and K.M. Hodivala-Dilke. 2002. Enhanced pathological angiogenesis in mice lacking beta3 integrin or beta3 and beta5 integrins. *Nature medicine*. 8:27-34.

- Ridley, A.J., M.A. Schwartz, K. Burridge, R.A. Firtel, M.H. Ginsberg, G. Borisy, J.T. Parsons, and A.R. Horwitz. 2003. Cell migration: integrating signals from front to back. *Science*. 302:1704-1709.
- Ryan, G.B., W.J. Cliff, G. Gabbiani, C. Irle, D. Montandon, P.R. Statkov, and G. Majno. 1974. Myofibroblasts in human granulation tissue. *Human pathology*. 5:55-67.
- Sabeh, F., I. Ota, K. Holmbeck, H. Birkedal-Hansen, P. Soloway, M. Balbin, C. Lopez-Otin, S. Shapiro, M. Inada, S. Krane, E. Allen, D. Chung, and S.J. Weiss. 2004. Tumor cell traffic through the extracellular matrix is controlled by the membrane-anchored collagenase MT1-MMP. *The Journal of cell biology*. 167:769-781.
- Schoumacher, M., R.D. Goldman, D. Louvard, and D.M. Vignjevic. 2010. Actin, microtubules, and vimentin intermediate filaments cooperate for elongation of invadopodia. *The Journal of cell biology*. 189:541-556.
- Schwartz, M.A., and D.W. DeSimone. 2008. Cell adhesion receptors in mechanotransduction. *Current opinion in cell biology*. 20:551-556.
- Serini, G., M.L. Bochaton-Piallat, P. Ropraz, A. Geinoz, L. Borsi, L. Zardi, and G. Gabbiani. 1998. The fibronectin domain ED-A is crucial for myofibroblastic phenotype induction by transforming growth factor-beta1. *The Journal of cell biology*. 142:873-881.
- Shao, J., G.G. Sheng, R.C. Mifflin, D.W. Powell, and H. Sheng. 2006. Roles of myofibroblasts in prostaglandin E2-stimulated intestinal epithelial proliferation and angiogenesis. *Cancer research*. 66:846-855.
- Shieh, A.C. 2011. Biomechanical forces shape the tumor microenvironment. *Annals of biomedical engineering*. 39:1379-1389.
- Smith, J.W. 2003. Cilengitide Merck. *Current opinion in investigational drugs*. 4:741-745.

- Smutny, M., and A.S. Yap. 2010. Neighborly relations: cadherins and mechanotransduction. *The Journal of cell biology*. 189:1075-1077.
- Springer, T.A., and J.H. Wang. 2004. The three-dimensional structure of integrins and their ligands, and conformational regulation of cell adhesion. *Advances in protein chemistry*. 68:29-63.
- Stoeltzing, O., W. Liu, N. Reinmuth, F. Fan, G.C. Parry, A.A. Parikh, M.F. McCarty, C.D. Bucana, A.P. Mazar, and L.M. Ellis. 2003. Inhibition of integrin alpha5beta1 function with a small peptide (ATN-161) plus continuous 5-FU infusion reduces colorectal liver metastases and improves survival in mice. *International journal of cancer. Journal international du cancer*. 104:496-503.
- Stuelten, C.H., S. DaCosta Byfield, P.R. Arany, T.S. Karpova, W.G. Stetler-Stevenson, and A.B. Roberts. 2005. Breast cancer cells induce stromal fibroblasts to express MMP-9 via secretion of TNF-alpha and TGF-beta. *Journal of cell science*. 118:2143-2153.
- Stupack, D.G., X.S. Puente, S. Boutsaboualoy, C.M. Storgard, and D.A. Cheresh. 2001. Apoptosis of adherent cells by recruitment of caspase-8 to unligated integrins. *The Journal of cell biology*. 155:459-470.
- Takagi, J., B.M. Petre, T. Walz, and T.A. Springer. 2002. Global conformational rearrangements in integrin extracellular domains in outside-in and inside-out signaling. *Cell*. 110:599-511.
- Taniwaki, K., H. Fukamachi, K. Komori, Y. Ohtake, T. Nonaka, T. Sakamoto, T. Shiomi, Y. Okada, T. Itoh, S. Itohara, M. Seiki, and I. Yana. 2007. Stroma-derived matrix metalloproteinase (MMP)-2 promotes membrane type 1-MMP-dependent tumor growth in mice. *Cancer research*. 67:4311-4319.
- Tlsty, T.D., and L.M. Coussens. 2006. Tumor stroma and regulation of cancer development. *Annual review of pathology*. 1:119-150.

- Wang, N., J.D. Tytell, and D.E. Ingber. 2009. Mechanotransduction at a distance: mechanically coupling the extracellular matrix with the nucleus. *Nature reviews. Molecular cell biology*. 10:75-82.
- Wang, W., R. Eddy, and J. Condeelis. 2007. The cofilin pathway in breast cancer invasion and metastasis. *Nature reviews. Cancer*. 7:429-440.
- Wang, W., G. Mouneimne, M. Sidani, J. Wyckoff, X. Chen, A. Makris, S. Goswami, A.R. Bresnick, and J.S. Condeelis. 2006. The activity status of cofilin is directly related to invasion, intravasation, and metastasis of mammary tumors. *The Journal of cell biology*. 173:395-404.
- Wiseman, B.S., and Z. Werb. 2002. Stromal effects on mammary gland development and breast cancer. *Science*. 296:1046-1049.
- Woodings, J.A., S.J. Sharp, and L.M. Machesky. 2003. MIM-B, a putative metastasis suppressor protein, binds to actin and to protein tyrosine phosphatase delta. *The Biochemical journal*. 371:463-471.
- Wu, Y., J. Zuo, G. Ji, H. Saiyin, X. Liu, F. Yin, N. Cao, Y. Wen, J.J. Li, and L. Yu. 2009. Proapoptotic function of integrin beta(3) in human hepatocellular carcinoma cells. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 15:60-69.
- Wyckoff, J., W. Wang, E.Y. Lin, Y. Wang, F. Pixley, E.R. Stanley, T. Graf, J.W. Pollard, J. Segall, and J. Condeelis. 2004. A paracrine loop between tumor cells and macrophages is required for tumor cell migration in mammary tumors. *Cancer research*. 64:7022-7029.
- Xiong, J.P., T. Stehle, R. Zhang, A. Joachimiak, M. Frech, S.L. Goodman, and M.A. Arnaout. 2002. Crystal structure of the extracellular segment of integrin alpha Vbeta3 in complex with an Arg-Gly-Asp ligand. *Science*. 296:151-155.

- Yamaguchi, H., M. Lorenz, S. Kempiak, C. Sarmiento, S. Coniglio, M. Symons, J. Segall, R. Eddy, H. Miki, T. Takenawa, and J. Condeelis. 2005. Molecular mechanisms of invadopodium formation: the role of the N-WASP-Arp2/3 complex pathway and cofilin. *The Journal of cell biology*. 168:441-452.
- Yu, H., J.K. Mouw, and V.M. Weaver. 2011. Forcing form and function: biomechanical regulation of tumor evolution. *Trends in cell biology*. 21:47-56.
- Zaman, M.H., L.M. Trapani, A.L. Sieminski, D. Mackellar, H. Gong, R.D. Kamm, A. Wells, D.A. Lauffenburger, and P. Matsudaira. 2006. Migration of tumor cells in 3D matrices is governed by matrix stiffness along with cell-matrix adhesion and proteolysis. *Proceedings of the National Academy of Sciences of the United States of America*. 103:10889-10894.
- Zebda, N., O. Bernard, M. Bailly, S. Welte, D.S. Lawrence, and J.S. Condeelis. 2000. Phosphorylation of ADF/cofilin abolishes EGF-induced actin nucleation at the leading edge and subsequent lamellipod extension. *The Journal of cell biology*. 151:1119-1128.

ABSTRACT**DEFINING THE MECHANISM OF ENHANCED CELLULAR INVASION INDUCED BY
MECHANICAL STIMULATION**

by

SNEHAL OZARKAR**May 2013****Advisor:** Dr. Karen Beningo**Major:** Biological Sciences**Degree:** Master of Science

Metastasis is a multistep process driven by various biochemical and mechanical factors, which eventually leads to formation of secondary tumors. The tumor mass is surrounded by basement membrane (BM) and stroma made of various extracellular matrix (ECM) proteins. During metastasis tumor cells disseminate from the primary tumor, breach the BM, invade the stroma, travel through blood and lymph and colonize tissues distant from the primary tumor. Formation of secondary tumors by metastasis is a leading cause of death in cancer patients. Even though plenty of research has been focused on biochemical factors affecting metastasis, information on role of mechanical factors in this process is very limited. Using our previously developed *in vitro* mechano-invasion assay, we had observed enhanced cellular invasion in response to tugging forces in the stroma during cancer cell invasion. In vivo, such tugging forces would be produced by contractile cells within the stroma as they migrate and remodel the matrix fibers. In addition, we found this mechanically enhanced invasion by cancer cells to be dependent on the presence of fibronectin in the extracellular matrix. The objective of our study is to understand the mechanotransduction pathway leading to enhanced invasion. We hypothesized that in response to mechanical forces in the stroma, tumor cells will show an altered expression

of genes involved in mechanosensing. We performed expression profiling of several genes related to cell migration, adhesion and tumor metastasis by real-time PCR analysis. Six genes were confirmed to be differentially expressed between mechanically stimulated and non-stimulated conditions. Surprisingly, one of the genes found to be significantly down-regulated in the mechanically stimulated invasion culture is a fibronectin specific integrin subunit, integrin $\beta 3$. Over-expression of this gene resulted in a significant decrease in enhanced invasion, supporting its role in sensing the mechanical stimulus. Furthermore, down-regulation of integrin $\beta 3$ resulted in decrease in amounts of inactive form of cofilin (Ser3 phospho-cofilin).

AUTOBIOGRAPHICAL STATEMENT**SNEHAL OZARKAR****Educational Qualification:**

2009-2013: Master of Science (M.S.) in Biological Sciences, Wayne State University, Detroit, MI

2007-2009: Master of Science (M.Sc.) in Bioanalytical Sciences, G. N. Khalsa College, University of Mumbai, India

2004-2007: Bachelor of Science (B.Sc.) in Biotechnology, University of Mumbai, India

Achievements and Awards:

2nd rank in University of Mumbai in Master of Science, April 2008

Thomas C. Rumble Graduate Fellowship, Wayne State University (August 2009 – July 2010)

Poster presentations:

- The 52nd annual meeting of American Society of Cell Biology, December 2012
- Frontiers of Tumor and Microenvironment Symposium, December 2012
- Department of Biological Sciences Research Retreat, October 2012 (3rd Position)
- Department of Biological Sciences Research Retreat, November 2010