



Correction to: Earlier chronotype in patients with rheumatoid arthritis

G. Esther A. Habers^{1,2} · Annette H. M. van der Helm-van Mil³ · Dieuwke S. Veldhuijzen^{1,2} · Cornelia F. Allaart³ · Erno Vreugdenhil⁴ · Daniëlle E. J. Starreveld⁵ · Tom W. J. Huizinga³ · Andrea W. M. Evers^{1,2,6,7,8}

Published online: 28 January 2021

© International League of Associations for Rheumatology (ILAR) 2021

Correction to: Clinical Rheumatology

<https://doi.org/10.1007/s10067-020-05546-x>

The original version of this article contained error.

In the last part of the formula under section “Norm group data for timing sleep-wake behaviour in general population (study aim 1)” was described “(=**33:52 h:mm**)”, it should have been “(=**3:52 h : mm**)”. The original article has been corrected.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s10067-020-05546-x>

✉ G. Esther A. Habers
g.e.a.habers@fsw.leidenuniv.nl

¹ Health, Medical, and Neuropsychology Unit, Institute of Psychology, Leiden University, Wassenaarseweg 52, 2333 AK Leiden, the Netherlands

² Leiden Institute for Brain and Cognition, Leiden, the Netherlands

³ Department of Rheumatology, Leiden University Medical Center, Leiden, the Netherlands

⁴ Department of Cell and Chemical Biology/Department of Dermatology, Leiden University Medical Center, Leiden, the Netherlands

⁵ Department of Psychosocial Research and Epidemiology, Netherlands Cancer Institute, Amsterdam, the Netherlands

⁶ Department of Psychiatry, Leiden University Medical Center, Leiden, the Netherlands

⁷ Industrial Design Engineering, Delft University of Technology, Delft, the Netherlands

⁸ Erasmus School of Health Policy and Management, Erasmus University, Rotterdam, the Netherlands