



Smartphone Use Associated with Refractive Error in Teenagers

The Myopia App Study

Clair A. Enthoven, MSc,^{1,2} Jan Roelof Polling, BSc,^{1,3} Timo Verzijden, MSc,¹ J. Willem L. Tideman, MD, PhD,¹ Nora Al-Jaffar, BSc,¹ Pauline W. Jansen, PhD,^{4,5} Hein Raat, MD, PhD,⁶ Lauwerens Metz, MSc,⁷ Virginia J.M. Verhoeven, MD, PhD,^{1,8} Caroline C.W. Klaver, MD, PhD^{1,9,10}

Purpose: To investigate the association between smartphone use and refractive error in teenagers using the Myopia app.

Design: Cross-sectional population-based study.

Participants: A total of 525 teenagers 12 to 16 years of age from 6 secondary schools and from the birth cohort study Generation R participated.

Methods: A smartphone application (Myopia app; Innovatic) was designed to measure smartphone use and face-to-screen distance objectively and to pose questions about outdoor exposure. Participants underwent cycloplegic refractive error and ocular biometry measurements. Mean daily smartphone use was calculated in hours per day and continuous use as the number of episodes of 20 minutes on screen without breaks. Linear mixed models were conducted with smartphone use, continuous use, and face-to-screen distance as determinants and spherical equivalent of refraction (SER) and axial length-to-corneal radius (AL:CR) ratio as outcome measures stratified by median outdoor exposure.

Main Outcome Measures: Spherical equivalent of refraction in diopters and AL:CR ratio.

Results: The teenagers on average were 13.7 ± 0.85 years of age, and myopia prevalence was 18.9%. During school days, total smartphone use on average was 3.71 ± 1.70 hours/day and was associated only borderline significantly with AL:CR ratio ($\beta = 0.008$; 95% confidence interval [CI], -0.001 to 0.017) and not with SER. Continuous use on average was 6.42 ± 4.36 episodes of 20-minute use without breaks per day and was associated significantly with SER and AL:CR ratio ($\beta = -0.07$ [95% CI, -0.13 to -0.01] and $\beta = 0.004$ [95% CI, 0.001 – 0.008], respectively). When stratifying for outdoor exposure, continuous use remained significant only for teenagers with low exposure ($\beta = -0.10$ [95% CI, -0.20 to -0.01] and $\beta = 0.007$ [95% CI, 0.001 – 0.013] for SER and AL:CR ratio, respectively). Smartphone use during weekends was not associated significantly with SER and AL:CR ratio, nor was face-to-screen distance.

Conclusions: Dutch teenagers spent almost 4 hours per day on their smartphones. Episodes of 20 minutes of continuous use were associated with more myopic refractive errors, particularly in those with low outdoor exposure. This study suggested that frequent breaks should become a recommendation for smartphone use in teenagers. Future large longitudinal studies will allow more detailed information on safe screen use in youth. *Ophthalmology* 2021;128:1681-1688 © 2021 by the American Academy of Ophthalmology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Supplemental material available at www.aaojournal.org.

Myopia is a refractive error caused by disproportionate eye growth during childhood and adolescence.¹ The prevalence of myopia is rising all over the world.^{2,3} Currently, almost 50% of the young adults in Europe and 80% to 90% of the young adults in urban areas of East Asia are myopic.^{2,4,5} Early onset of myopia results in higher degrees of myopia in adulthood.^{6,7} This can lead to visual impairment and even blindness resulting from retinal complications later in life.^{8,9} The rise in myopia prevalence in the last decade is caused by many lifestyle and behavioral changes.¹⁰ For instance, spending less time outdoors is an established risk

factor; the role of prolonged near work is still debated, but many reports conclude an association.^{11–13} These environmental factors also may explain why children growing up in urban areas more often are myopic than those growing up in rural areas.^{14–16}

In recent years, researchers have speculated that smartphone use is an additional risk factor for myopia. Time spent on smartphones adds considerably to the total hours spent on near work among teenagers.¹⁷ However, the so-called myopia boom started in 1950,¹⁸ when smartphones did not yet exist. Smartphones are relatively new, and children

growing up with smartphones are yet to become adults. Long-term effects, including the influence on the myopia prevalence, are yet to be determined. Smartphone use is prone to underreporting and therefore is difficult to determine by questionnaire.¹⁹ For the current study, we developed a smartphone application (the Myopia app; Innovatic) that registers smartphone use and face-to-screen distance electronically to allow for objective measurements. We assessed the associations among smartphone use, outdoor exposure, and refractive error as measured by the Myopia app and self-reported outdoor exposure. We hypothesized that increased smartphone use is associated with a more myopic refractive error and that this association may be modified by outdoor exposure.

Methods

Study Populations: Myopia App Study and Generation R

Teenagers 12 to 16 years of age from 2 cohorts were eligible to enrol in the study: participants in the Myopia App Study (MAS) and the Generation R study. The MAS participants were recruited from 6 secondary schools in semiurban areas in The Netherlands. Schools were asked to disseminate information on MAS among their pupils, and 300 teenagers from the first, second, and third grades (ages, 12–16 years) consented to participate (Fig S1, available at www.aaojournal.org). Generation R is a large, prospective, population-based birth cohort in which 9778 pregnant mothers were enrolled between 2002 and 2006. Details of the methodology of this study have been described elsewhere.^{20,21} Of the initial cohort, 4929 children (50%) visited the research center at 13 years of age. The app measurements were introduced during the final part of the study phase in April 2019, and 225 teenagers signed informed consent (Fig S1).

The app and ophthalmic measurements were performed between November 2018 and December 2019 in both cohorts. Two participants did not undergo eye measurements; 361 participants installed the app. Valid smartphone and eye measurements were available for 272 participants, because 25% of participants did not allow the app to run in the background of the operating system or technical issues hampered registration (Fig S1). Written informed consent from both parents and the teenagers was obtained before eye examination and app measurements. The study protocol was approved by the Medical Ethical Committee of the Erasmus Medical Center, Rotterdam (identifiers, MEC-2018-005, NL63977.078.17 [MAS] and MEC-217.595/2002/20 [Generation R study]). The study project was conducted according to the tenets of the Declaration of Helsinki.

Mobile Application

The Myopia app was developed by the company Innovatic (www.innovatic.com) and was made available for the smartphone operation systems iOS and Android. This smartphone logging app registered smartphone use and face-to-screen distance (see next section). The teenagers received questions about outdoor exposure twice weekly through pop-up notifications in the app. To encourage the teenagers to answer all questions, gamification techniques were implemented in the app, that is, different levels were used to perform the measurements. Participants were rewarded with extra points after a questionnaire was completed, and an avatar received new gadgets (i.e., hat or sunglasses) with an increasing number of points. After 5 weeks, the teenagers were

rewarded with an online shopping voucher with a value corresponding to the amount of questions answered (up to €7.50).

Smartphone Use

Smartphone use was measured over 5 weeks. The time of locking and unlocking the smartphone was registered using Unix time stamps, and participants were advised not to close the app. In that way, the app continued running in the background, which was needed because the closed operating systems of iOS and Android hampered continuous registration. We took particular care to identify measurement errors that occurred when participants (unintentionally) closed the app. Depending on whether the last measurement was registered as screen off or screen on before the app stopped running in the background, this resulted in days with very low smartphone use or extremely long continuous smartphone use. Days with fewer than 5 minutes of smartphone use in total or days with more than 5 hours of continuous use without locking the screen were excluded (on average, 7.9 days per participant [33.9%]), resulting in an average of 19.7 measurement days (standard deviation [SD], 14.5 measurement days; median, 17.0 measurement days; interquartile range [IQR] 7.0 to 30.0, 23 measurement days) per participant. To check for bias because of measurement error, we also excluded days with less than 1 minute of smartphone use in total or days with more than 4 hours of continuous use (on average, 8.7 days per participant [35.7%]), resulting in an average of 19.0 measurement days (SD, 14.0 measurement days; median, 17.0 measurement days; IQR 7.0 to 28.0, 21.0 measurement days) per participant, and days with less than 10 minutes of smartphone use in total. Excluding days with more than 6 hours of continuous use (on average, 7.4 days per participant [32.1%]) resulted in an average of 20.3 measurement days (SD, 14.7 measurement days; median, 18.0 measurement days; IQR 8.0 to 30.0, 22.0 measurement days) per participant. The main analyses were performed using the first data processing manner (excluding days with < 5 minutes in total and > 5 hours of continuous use). Sensitivity analyses were performed using the second (more strict) and third (less strict) data processing manner (excluding days with < 1 minute in total and > 4 hours of continuous use and excluding days with < 10 minutes in total and > 6 hours of continuous use) to ensure that the association between smartphone use and refractive error was not driven by our choice of excluding measurement days.

Smartphone use (hours per day) was calculated by summing the total time of smartphone use divided by the number of days the app was running. Continuous smartphone use was calculated by the sum of screen times of 20 minutes or longer divided by 20. For example, if a participant had 5, 53, 22, 19, and 68 minutes of smartphone use on one day, then continuous use was calculated by summing 53, 22, and 68 (143 minutes) divided by 20, that is, 7.15 episodes of 20 minutes of continuous smartphone use. Continuous use was determined by the sum of these episodes divided by the number of days the app was running. Smartphone use and continuous use were calculated for school days and non-school days separately. Non-school days consisted for 75.5% of weekend days and 24.5% of holidays. The density plots of smartphone use and continuous smartphone use during school days defined by the 3 different data processing manners are shown in Figure S2 (available at www.aaojournal.org).

Validation Study

We performed a validation study that included 5 Android users and 5 iOS users. They installed the Myopia app on their smartphone for 2 weeks. Smartphone use measured by the Myopia app was compared with smartphone use measured by the inbuilt screen time

tracker of the smartphone. The Spearman correlation coefficient between the smartphone use measured by the Myopia app, and the smartphone use measured by the inbuilt app was calculated.

Face-to-Screen Distance

Face-to-screen distance was measured using the front camera of the smartphone. Android device users calibrated the app by holding their smartphone exactly 29.7 cm in front of their eyes (the length of the long side of an A4 piece of paper); iOS device users did not need to calibrate face-to-screen measurement because of the technical similarities among iPhones. Face-to-screen distance was measured when the app was active and open (i.e., when participants were filling out questions). The number of face-to-screen measurements on average was 592 measurements (SD, 1246 measurements; median, 272.0 measurements; IQR 152.0 to 555.3, 403.3 measurements) per person. Mean face-to-screen distance was calculated. Sensitivity analyses were performed excluding participants with fewer than 100 measurements to ensure that measurement reflected most commonly used smartphone distance.

Outdoor Exposure

Outdoor exposure was asked repeatedly in the app for 5 weeks. On Monday afternoon and Friday evening, the participants received the question: “How much time did you spend outdoors last Saturday/Sunday/Monday or Tuesday/Wednesday/Thursday/Friday? For example, cycling, sports, walking, playing outdoors, or being outdoors with friends or family.” Mean outdoor exposure per day (in hours per day) was calculated for school days and non-school days separately.

Other Covariates

Sex, age at examination, season of app measurement, ethnic background, and operating system (iOS or Android) were considered as covariates. Ethnic background was defined according to the definitions by Health Statistics Netherlands, that is, based on the country of birth of the (grand) parents. It was assessed through a questionnaire in the app for the MAS participants and by questionnaires filled out by the parents for the Generation R participants and was stratified into European and non-European backgrounds. Operating system was assessed through the app.

Eye Measurements

The eye examination consisted of presenting monocular visual acuity with logarithm of the minimum angle of resolution-based Early Treatment Diabetic Retinopathy Study charts at 3 m by means of the fast Early Treatment Diabetic Retinopathy Study method. Ocular biometry was measured by Zeiss IOLMaster 500 or 700 (Carl Zeiss Meditec). Five axial length measurements per eye were averaged to calculate mean axial length; 3 measurements of corneal radius (K1 and K2) were averaged to calculate the mean corneal radius, and axial length-to-corneal radius (AL:CR) ratio was calculated. Cycloplegic refractive error of the nondominant eye was measured with handheld Retinomax 3 (Righton) in the MAS participants, of both eyes in the Generation R participants, both 30 minutes after 2 doses of cyclopentolate 1%. Spherical equivalent of refraction (SER) was calculated by the sum of the full spherical value plus half of the negative cylindrical value. Mean SER for Generation R participants was assessed by averaging SER of the right and left eyes. Myopia was defined as SER of -0.50 diopter (D) or less.

Data Analyses

Differences between participants who were included in the analyses and who were excluded because of missing data, as well as differences between the school-based cohort and Generation R cohort, were analyzed with independent *t* tests for continuous variables and chi-square tests for dichotomous variables. Spearman correlation coefficients were calculated for smartphone use, continuous use, face-to-screen distance, and outdoor exposure during school days and weekend days. To take into account the similarities between teenagers from the same study site, linear mixed models with restricted likelihood estimation from the nlmer package in R software (R Foundation for Statistical Computing) were used to perform the analyses (Table S1, available at www.aojournal.org).²² The associations between smartphone use, continuous use (20 minutes), outdoor exposure, and face-to-screen distance as exposures and SER and AL:CR ratio as outcomes variables were investigated, with random intercept for study sites (schools), and adjusted for age, sex, season of app measurement, and operating system (iOS or Android). The following sensitivity analyses were performed. First, outliers in smartphone use and continuous use were excluded, that is, $> 4/6$ hours continuous use, and days with $< 1/10$ minutes smartphone use (see above). Second, we additionally adjusted for outdoor exposure to ensure an independent association among smartphone use, continuous use, SER, and AL:CR ratio. Third, participants with fewer than 100 measurements for face-to-screen distance were excluded (see previous). Fourth, because of the large number of missing data for ethnicity and because the MAS participants were 97% European, we did not adjust for ethnicity in the main analyses but instead performed sensitivity analyses with European participants only. Finally, interaction analysis was performed with smartphone use, outdoor exposure, and an interaction term as exposures and SER and AL:CR ratio as outcomes variables, with random intercept for study sites (schools), and adjusted for age, sex, and operating system. Stratified analyses were performed for teenagers with high and low outdoor exposure based on the median. Analyses were performed in IBM SPSS version 25 and R statistical software version 3.6.1.

Results

The teenagers on average were 13.7 ± 0.85 years of age; 54% were girls. Myopia prevalence was 18.9%, SER was $+0.40 \pm 1.90$ D, AL:CR ratio was 2.99 ± 0.11 , and axial length was 23.4 ± 0.88 mm. The teenagers spent on average 3.71 ± 1.70 hours/day on their smartphone on school days and 3.82 ± 2.09 hours/day on non-school days, with an average face-to-screen distance of 29.1 ± 6.25 cm. Participants had 6.42 ± 4.36 episodes of 20 minutes of continuous use per day during school days and 7.10 ± 5.28 episodes during non-school days. Outdoor exposure was 2.37 ± 0.94 hours/day on school days and 2.77 ± 1.13 hours/day on non-school days. Participants with myopia demonstrated a more negative SER and larger AL:CR ratio and axial length compared with participants without myopia. Differences between participants with ($n = 45$) and without ($n = 193$) myopia regarding sex, ethnicity, smartphone use, continuous use, face-to-screen distance, outdoor exposure, season of app measurement, operating system, and study site did not reach statistical significance (Table 1).

Variables that differed between the MAS cohort and Generation R were age ($P = 0.02$), ethnic background ($P < 0.001$), and outdoor exposure during school days ($P = 0.01$). Participants who were included in the analyses were younger (13.7 years vs. 13.9 years; $P = 0.01$) and more often from a European ethnic background (86.5% vs. 67.9%; $P \leq 0.001$) than those who were not included because of missing data on smartphone use and eye

Table 1. General Characteristics

	Total (n = 272)	Missing (%)	Myopia (n = 45)	No Myopia (n = 193)	P Value
Age (yrs)	13.7 ± 0.85	0.0	13.5 ± 0.96	13.7 ± 0.87	0.36
Sex (female)	53.7	0.0	60.0	52.3	0.41
Ethnicity (European)	86.5	15.4	81.8	87.7	0.39
Spherical equivalent (diopters)	0.40 ± 1.90	12.5	-2.36 ± 2.10	1.04 ± 1.11	< 0.001
Myopia	18.9	12.5	NA	NA	NA
Axial length corneal radius ratio	2.99 ± 0.11	2.6	3.14 ± 0.13	2.96 ± 0.08	< 0.001
Axial length (mm)	23.4 ± 0.88	0.4	24.2 ± 0.91	23.2 ± 0.73	< 0.001
Smartphone use (hr/day)					
During school days	3.71 ± 1.70	7.7	3.75 ± 1.55	3.67 ± 1.73	0.78
During non-school days	3.82 ± 2.09	5.9	3.54 ± 2.11	3.77 ± 2.09	0.52
Continuous use (episodes of ≥ 20 min)					
During school days	6.42 ± 4.36	7.7	6.62 ± 4.32	6.13 ± 4.17	0.50
During non-school days	7.10 ± 5.28	5.9	6.51 ± 5.95	6.91 ± 5.11	0.66
Face-to-screen distance (cm)	29.1 ± 6.3	14.7	29.1 ± 7.47	29.4 ± 5.72	0.76
Outdoor exposure (hr/day)					
During school days	2.37 ± 0.94	11.8	2.10 ± 0.90	2.41 ± 0.96	0.06
During non-school days	2.77 ± 1.13	1.5	2.48 ± 1.21	2.83 ± 1.07	0.05
Season of app measurement		0.0			0.65
Spring	71.3		66.6	72.0	
Summer	20.2		20.0	19.2	
Autumn	8.5		13.3	8.8	
Operating system (Android)	60.7	0.0	68.9	59.1	0.24
Study site		0.0			0.16
Generation R	25.7		22.2	15.5	
School 1	36.4		28.9	44.6	
School 2	13.6		20.0	14.0	
School 3	8.8		11.1	8.3	
School 4	4.0		0.0	5.7	
School 5	4.0		8.8	3.6	
School 6	7.4		8.8	8.3	

Data are presented as mean ± standard deviation or percentage.

measurements. Differences between children included in the analysis and those excluded regarding sex, SER, myopia, axial length, and AL:CR ratio were not observed. The Spearman

correlation coefficient between the Myopia app and the inbuilt app in our validation study was 0.97 (Fig S3, available at www.aaojournal.org).




Figure 1. Diagram showing correlations between smartphone use, continuous use, face-to-screen distance, and outdoor exposure during school days and holidays. Dark blue represents a positive correlation of 1, whereas dark red represents a negative correlation of -1.




Figure 2. Scatterplots showing the association between continuous smartphone use (episodes of ≥ 20 minutes) and (A) spherical equivalent and (B) axial length-to-corneal radius ratio. Blue lines represent the unadjusted regression lines. D = diopter.

Correlations between smartphone use, face-to-screen distance, and outdoor exposure are depicted in Figure 1. Smartphone use, face-to-screen distance, and outdoor exposure were distributed normally; continuous use was slightly right skewed (Fig S2). Smartphone use was correlated strongly with continuous use ($r = 0.86$ and $P < 0.001$ during school days; $r = 0.90$ and $P < 0.001$ during weekend days), and outdoor exposure was correlated inversely with smartphone use and continuous use (smartphone use: $r = -0.19$ and $P = 0.006$ during school days; $r = -0.21$ and $P = 0.003$ during weekend days; continuous use: $r = -0.24$ and $P < 0.001$ during school days; $r = -0.26$ and $P < 0.001$ during weekend days). Face-to-screen distance was not correlated with smartphone use, continuous use, or outdoor exposure.

Continuous use during school days was associated with SER (per each extra episode of 20 minutes continuous use: $\beta = -0.07$ [95% CI, -0.13 to -0.01]) and AL:CR ratio ($\beta = 0.004$ [95% CI, 0.001 – 0.008]; Fig 2). Smartphone use during school days showed a similar trend and was borderline associated significantly with AL:CR ratio ($\beta = 0.008$ [95% CI, -0.001 to 0.017]) but not

with SER ($\beta = -0.09$ [95% CI, -0.25 to 0.07]). Outdoor exposure was associated with SER ($\beta = 0.33$ [95% CI, 0.07 – 0.60]) and $\beta = 0.32$ [95% CI, 0.10 – 0.55] both during school days) and with AL:CR ratio during non-school days ($\beta = -0.016$ [95% CI, -0.029 to -0.003]). Face-to-screen distance, continuous use during non-school days, and smartphone use during non-school days were not associated with SER or AL:CR ratio (Table 2). Sensitivity analyses with different definitions of smartphone use or adjustment for outdoor exposure yielded similar results; excluding non-Europeans and those with missing data on ethnicity resulted in similar, albeit not significant, β coefficients. Face-to-screen distance excluding participants with fewer than 100 measurements was not associated significantly with SER or AL:CR ratio (Table S1 available at www.aojournal.org).

Stratified analyses showed that the association between continuous use and SER and AL:CR ratio was observed for teenagers with low outdoor exposure ($\beta = -0.10$ [95% CI, -0.20 to -0.01] for SER and $\beta = 0.007$ [95% CI, 0.001 – 0.013] for AL:CR ratio) but not for teenagers with high outdoor exposure (Table 3).

Table 2. Linear Regression Analyses of Smartphone Use, Continuous Use during School Days and Non-School Days, and Face-to-Screen Distance on Spherical Equivalent and Axial Length-to-Corneal Radius Ratio

	Spherical Equivalent Refraction					Axial Length-to-Corneal Radius Ratio				
	No.	Estimate	Standard Error	95% Confidence Interval	P Value	No.	Estimate	Standard Error	95% Confidence Interval	P Value
Smartphone use (hr/day) during school days	207	-0.09	0.08	-0.25 to -0.07	0.30	227	0.008	0.005	-0.001 to -0.017	0.10
Continuous use (≥ 20 min) during school days	207	-0.07	0.03	-0.13 to -0.01	0.03	227	0.004	0.002	0.001–0.008	0.02
Smartphone use (hr/day) during non-school days	204	-0.02	0.10	-0.21 to -0.18	0.88	226	0.002	0.006	-0.010 to -0.013	0.75
Continuous use (≥ 20 min) during non-school days	204	-0.03	0.03	-0.11 to -0.04	0.34	226	0.002	0.002	-0.002 to -0.006	0.29
Outdoor exposure (hr/day) during school days	213	0.33	0.13	0.07–0.60	0.01	235	-0.011	0.008	-0.027 to -0.005	0.17
Outdoor exposure (hr/day) during non-school days	235	0.32	0.11	0.10–0.55	0.004	261	-0.016	0.006	-0.029 to -0.003	0.02
Face-to-screen distance	201	0.00	0.02	-0.04 to -0.04	0.98	226	0.000	0.001	-0.003 to -0.002	0.84

Adjusted for age, sex, season of app measurement, and operating system.

Table 3. Linear Regression Analyses of Smartphone Use and Continuous Use during School Days and Holidays on Spherical Equivalent Refraction and Axial Length-to-Corneal Radius Ratio Stratified by High versus Low Outdoor Exposure

Outdoor Exposure	Smartphone Use during School Days	Spherical Equivalent Refraction					Axial Length-to-Corneal Radius Ratio				
		No.	Estimate	Standard Error	95% Confidence Interval	P Value	No.	Estimate	Standard Error	95% Confidence Interval	P Value
Low	Hours/day	99	-0.12	0.13	-0.36 to -0.12	0.35	112	0.010	0.007	-0.004 to -0.024	0.17
	Continuous use (≥20 min)	99	-0.10	0.05	-0.20 to -0.01	0.03	112	0.007	0.003	0.001-0.013	0.02
High	Hours/day	99	-0.04	0.11	-0.25 to -0.17	0.72	105	0.003	0.006	-0.009 to -0.014	0.65
	Continuous use (≥20 min)	99	-0.02	0.05	-0.12 to -0.07	0.61	105	0.001	0.002	-0.003 to -0.006	0.59

Adjusted for age, sex, season of app measurement, and operating system.

However, the interaction term between continuous use and outdoor exposure was not significant ($P = 1.00$ for SER; $P = 0.40$ for AL:CR ratio).

Discussion

In this study, we used a mobile application to determine smartphone use in relationship to refractive error. We showed that those with more episodes of continuous use demonstrated a more myopic refractive error. This association disappeared in teenagers with high outdoor exposure, suggesting that outdoor exposure may moderate this effect.

Smartphone use is a relatively new behavior among youth. It became increasingly popular after the introduction of the first iPhone in 2008. Worldwide, 139 million smartphones were sold in 2008, which increased to 1496 million smartphones sold in 2016. Most smartphone owners are from the United States and Western Europe, but the Chinese market is also on the rise.²³ Research reports addressing the effect of smartphone use on myopia in teenagers are scarce. In our study, smartphone use was 3.71 hours/day during school days according to the Myopia app, which is comparable with the 4 hours/day among 19-year-old university students from the United States measured with the Moment app.¹⁷ A Chinese study showed that 1 hour/day increase in smartphone use was associated with -0.28 D SER after adjustment for age, sex, reading behavior, outdoor exposure, and sleep in 566 children 6 to 14 years of age.²⁴ We observed a particular association with continuous use: SER was -0.07 D more myopic and AL:CR ratio was 0.005 larger for each extra episode of 20 minutes of continuous use. The SER was -0.10 D more myopic and the AL:CR ratio 0.008 larger for each hour of daily smartphone use, but this association was not significant ($P = 0.22$ for SER and $P = 0.07$ for AL:CR ratio). Studies focussing on reading behavior also reported that continuous reading was associated more prominently with myopia than total reading time,^{12,25} despite their high correlation. Continuous near work may be a more important risk factor than time spent on near work, suggesting that regular breaks during near work (including smartphone use) will help to prevent myopia from developing in teenagers.

Although the association between screen time and myopia was debatable for a long time,^{26,27} recently, the results of many studies support the presence of such an association.²⁸⁻³² Exposure to screen time before the age of 1 year was associated with myopia (prevalence ratio, 4.02) among 26,433 preschool children in China.²⁸ Irish school children who spent more than 3 hours/day on a screen more often were myopic (odds ratio, 3.70), and a 1-hour increase in computer use was associated with myopia (odds ratio, 1.005) in a former study among 9-year-old children.^{29,30} Adolescents using a screen for more than 6 hours/day more often were myopic than those with fewer than 2 hours/day of screen use (odds ratio, 1.95) in Copenhagen.³¹ A longitudinal study among 5- to 15-year-old children from India showed that more than 7 hours/day of screen time also was associated with myopia progression compared with fewer than 4 hours/day of screen time (odds ratio, 3.53).³² Together with our current findings, this suggests that screen use may become an established risk factor for myopia.

Reading distance has been identified as a risk factor for myopia in many cross-sectional and longitudinal studies.^{12,25,30,33} Reading distance often was measured using a questionnaire for parents, and these studies reported positive associations for 30 cm,^{25,30} 20 cm,¹² and 33 cm.³³ The sensitivity analysis in our study showed that a 1-cm-shorter face-to-screen distance was associated with -0.03 D (95% CI, 0.02 to -0.08 D) more myopia, but this association failed to reach statistical significance. Face-to-screen distance was not correlated with smartphone use in our study. Ip et al²⁵ and Li et al¹² did not identify a correlation between reading distance and reading time either, adding to the discrepancies in the associations with refractive error for continuous smartphone use and face-to-screen distance.

Strengths of this study are the objective measurement of smartphone use and face-to-screen distance using the Myopia app. The Myopia app was made available for both iOS and Android devices and thus was accessible to almost any smartphone user. Our validation study showed a high correlation between smartphone use measured by the Myopia app and smartphone use measured by the inbuilt screen time tracker of the smartphone, supporting an accurate registration. Sensitivity analyses with different

definitions of smartphone use yielded similar results, indicating that the association was robust. Nevertheless, some limitations should be borne in mind. First, the cross-sectional design of this study hindered causal interpretation of the data. Current smartphone use most likely reflects previous smartphone use; however, cumulative smartphone use depends on the age of smartphone acquisition. In the Netherlands, most children own a smartphone from the age of 10 years onward,³⁴ and we expect that most teenagers in our study already had 2 to 3 years of smartphone exposure time. Second, the relatively large number of days with unrealistic measurements and the limited sample size may have led to inconclusive results. Future studies should incorporate a longitudinal study design in a large sample. Third, some activities on the smartphone, like calling someone, were registered as smartphone use, while not involving near work. Yet because time spent on calling is usually very short in teenagers of this age, we do not expect that this had a major influence on our results.³⁴ Finally, only the nondominant eye was measured with cycloplegia in the MAS participants. Nondominant eyes may be more hyperopic than dominant eyes in children with anisometropia.^{35,36} This may have resulted in an underrepresentation of myopia in the MAS participants

but did not distort AL:CR ratio because this was measured in both eyes.

In conclusion, our study showed that Dutch teenagers use their smartphone almost 4 hours/day. A higher number of episodes of more than 20 minutes of continuous use was associated with more myopic SER and a larger AL:CR ratio. This association was not present in teenagers with high outdoor exposure, suggesting that outdoor exposure moderates the association. Because smartphone use is becoming increasingly popular, awareness of the potential negative consequences of prolonged smartphone use is warranted. The 20-20-2 rule as recommended earlier remains good advice.³⁷

Acknowledgments

The authors thank the participating schools, principals, teachers, and children in the Myopia App Study, and the Generation R children and parents, general practitioners, hospitals, midwives, and pharmacies in Rotterdam. The Generation R Study is conducted by the Erasmus Medical Center in close collaboration with the School of Law and Faculty of Social Sciences of the Erasmus University Rotterdam, the Municipal Health Service Rotterdam area, the Rotterdam Homecare Foundation, and the Stichting Trombosedienst & Artsenlaboratorium Rijnmond, Rotterdam.

Footnotes and Disclosures

Originally received: March 2, 2021.

Final revision: June 18, 2021.

Accepted: June 29, 2021.

Available online: July 8, 2021.

Manuscript no. D-21-00436.

¹ Department of Ophthalmology and Epidemiology, Erasmus Medical Center, Rotterdam, Zuid-Holland, The Netherlands.

² The Generation R Study Group, Erasmus Medical Center, Rotterdam, Zuid-Holland, The Netherlands.

³ Orthoptics and Optometry, University of Applied Sciences, Utrecht, Utrecht, The Netherlands.

⁴ Department of Child and Adolescent Psychiatry/Psychology, Erasmus Medical Center, Rotterdam, Zuid-Holland, The Netherlands.

⁵ Department of Psychology, Education and Child Studies, Erasmus University Rotterdam, Rotterdam, Zuid-Holland, The Netherlands.

⁶ Department of Public Health, Erasmus MC, Rotterdam, Zuid-Holland, The Netherlands.

⁷ Innovatic, Delft, Zuid Holland, The Netherlands.

⁸ Clinical Genetics, Erasmus Medical Center, Rotterdam, Zuid-Holland, The Netherlands.

⁹ Department of Ophthalmology, Radboud University Medical Center, Nijmegen, Gelderland, The Netherlands.

¹⁰ Institute of Molecular and Clinical Ophthalmology, Basel, Switzerland.

Disclosure(s):

All authors have completed and submitted the ICMJE disclosures form.

The author(s) have made the following disclosure(s): P.W.J.: Financial support – ZonMW

H.R.: Board member – Kenter Jeugdhulp

L.M.: Owner – Innovatic

C.C.W.K.: Board member – LSBS; Consultant – Bayer

Supported by Oogfonds, ODAS, Uitzicht 2017-28 (LSBS, MaculaFonds, Oogfonds); Netherlands Organization for Scientific Research (grant nos.:

91815655 [C.C.W.K.], 91617076 [V.J.M.V.], and 636320005 [P.W.J.]); and the European Research Council under the European Union's Horizon 2020 research and innovation programme (grant no.: 648268 [C.C.W.K.]). The funding organizations had no role in the design or conduct of this research. They provided unrestricted grants.

HUMAN SUBJECTS: Human subjects were included in this study. The human ethics committees at Erasmus Medical Center approved the study. All research adhered to the tenets of the Declaration of Helsinki. All participants and their parents provided informed consent.

No animal subjects were included in this study.

Author Contributions:

Conception and design: Enthoven, Polling, Tideman, Verhoeven, Klaver
Analysis and interpretation: Enthoven, Polling, Verzijden, Tideman, Jansen, Verhoeven, Klaver

Data collection: Enthoven, Polling, Verzijden, Al-Jaffar, Metz

Obtained funding: Verhoeven, Klaver

Overall responsibility: Enthoven, Polling, Tideman, Jansen, Raat, Metz, Verhoeven, Klaver

Abbreviations and Acronyms:

AL:CR = axial length-to-corneal radius; **CI** = confidence interval; **D** = diopter; **iOS** = iPhone operating system; **IQR** = interquartile range; **MAS** = Myopia App Study; **SD** = standard deviation; **SER** = spherical equivalent of refraction.

Keywords:

Myopia, Outdoor exposure, Refractive error, Smartphone use, Teenagers.

Correspondence:

Caroline C.W. Klaver, MD, PhD, Erasmus Medical Center, Room Na-2808, P. O. Box 2040, 3000 CA, Rotterdam, The Netherlands. E-mail: c.w.klaver@erasmusmc.nl.

References

- Tideman JW, Polling JR, Vingerling JR, et al. Axial length growth and the risk of developing myopia in European children. *Acta Ophthalmol.* 2018;96(3):301–309.
- Chen M, Wu A, Zhang L, et al. The increasing prevalence of myopia and high myopia among high school students in Fenghua city, eastern China: a 15-year population-based survey. *BMC Ophthalmol.* 2018;18(1):159.
- Holden BA, Fricke TR, Wilson DA, et al. Global prevalence of myopia and high myopia and temporal trends from 2000 through 2050. *Ophthalmology.* 2016;123(5):1036–1042.
- Williams KM, Verhoeven VJM, Cumberland P, et al. Prevalence of refractive error in Europe: the European Eye Epidemiology (E3) Consortium. *Eur J Epidemiol.* 2015;30(4):305–315.
- Pan C-W, Dirani M, Cheng C-Y, et al. The age-specific prevalence of myopia in Asia: a meta-analysis. *Optom Vis Sci.* 2015;92(3):258–266.
- Pärssinen O, Kauppinen M, Viljanen A. The progression of myopia from its onset at age 8–12 to adulthood and the influence of heredity and external factors on myopic progression. A 23-year follow-up study. *Acta Ophthalmol.* 2014;92(8):730–739.
- Fledelius HC. Myopia profile in Copenhagen medical students 1996–98. Refractive stability over a century is suggested. *Acta Ophthalmol Scand.* 2000;78(5):501–505.
- Verhoeven VJ, Wong KT, Buitendijk GH, et al. Visual consequences of refractive errors in the general population. *Ophthalmology.* 2015;122(1):101–109.
- Tideman JL, Snabel MC, Tedja MS, et al. Association of axial length with risk of uncorrectable visual impairment for Europeans with myopia. *JAMA Ophthalmol.* 2016;134(12):1355–1363.
- Morgan IG, French AN, Rose KA. Risk factors for myopia: putting causal pathways into a social context. In: Ang M, Wong TY, eds. *Updates on Myopia: A Clinical Perspective.* Singapore: Springer Singapore; 2020:133–170.
- Huang HM, Chang DST, Wu PC. The association between near work activities and myopia in children—a systematic review and meta-analysis. *Plos One.* 2015;10(10):e0140419.
- Li SM, Li SY, Kang MT, et al. Near work related parameters and myopia in Chinese children: the Anyang Childhood Eye Study. *PLoS One.* 2015;10(8):e0134514.
- Deng L, Pang Y. Effect of outdoor activities in myopia control: meta-analysis of clinical studies. *Optom Vis Sci.* 2019;96(4):276–282.
- Morgan IG, Ohno-Matsui K, Saw S-M. Myopia. *Lancet.* 2012;379(9827):1739–1748.
- Lin LL-K, Shih Y-F, Hsiao CK, Chen CJ. Prevalence of myopia in Taiwanese schoolchildren: 1983 to 2000. *Ann Acad Med Singap.* 2004;33(1):27–33.
- Rudnicka AR, Kapetanakis VV, Wathern AK, et al. Global variations and time trends in the prevalence of childhood myopia, a systematic review and quantitative meta-analysis: implications for aetiology and early prevention. *Br J Ophthalmol.* 2016;100(7):882.
- Elhai JD, Tiarniyu MF, Weeks JW, et al. Depression and emotion regulation predict objective smartphone use measured over one week. *Pers Individ Dif.* 2018;133:21–28.
- Dolgin E. The myopia boom. *Nature.* 2015;519(7543):276–278.
- Andrews S, Ellis DA, Shaw H, Piwek L. Beyond self-report: tools to compare estimated and real-world smartphone use. *PLoS One.* 2015;10(10). e0139004–e0139004.
- Kooijman MN, Kruithof CJ, van Duijn CM, et al. The Generation R Study: design and cohort update 2017. *Eur J Epidemiol.* 2016;31(12):1243–1264.
- Kruithof CJ, Kooijman MN, van Duijn CM, et al. The Generation R Study: biobank update 2015. *Eur J Epidemiol.* 2014;29(12):911–927.
- Pinheiro J, Bates D, DebRoy S, et al. *Package “nlme.” Linear and nonlinear mixed effects models, version. 3.* CRAN; 2017, 2017;3.
- O’Dea S. Number of smartphones sold to end users worldwide from 2007 to 2020. 2020; Available at: <https://www.statista.com/statistics/263437/global-smartphone-sales-to-end-users-since-2007>; Accessed 25.03.20.
- Liu S, Ye S, Xi W, Zhang X. Electronic devices and myopic refraction among children aged 6–14 years in urban areas of Tianjin, China. *Ophthalmic Physiol Opt.* 2019;39(4):282–293.
- Ip JM, Saw S-M, Rose KA, et al. Role of near work in myopia: findings in a sample of Australian school children. *Invest Ophthalmol Vis Sci.* 2008;49(7):2903–2910.
- La Torre G, Pacella E. Use of tablet, smartphone and myopia: where is the evidence? *Senses and Sciences.* 2014;1(3):82–83.
- Lanca C, Saw SM. The association between digital screen time and myopia: a systematic review. *Ophthalmic Physiol Opt.* 2020 Mar;40(2):216–229.
- Yang G-Y, Huang L-H, Schmid KL, et al. Associations between screen exposure in early life and myopia amongst Chinese preschoolers. *Int J Environ Res Public Health.* 2020;17(3):1056.
- Harrington SC, Stack J, O’Dwyer V. Risk factors associated with myopia in schoolchildren in Ireland. *Br J Ophthalmol.* 2019;103(12):1803–1809.
- Enthoven CA, Tideman JW, Polling JR, et al. The impact of computer use on myopia development in childhood: the Generation R study. *Prev Med.* 2020;132:105988.
- Hansen MH, Laigaard PP, Olsen EM, et al. Low physical activity and higher use of screen devices are associated with myopia at the age of 16-17 years in the CCC2000 Eye Study. *Acta Ophthalmol.* 2020;98(3):315–321.
- Saxena R, Vashist P, Tandon R, et al. Incidence and progression of myopia and associated factors in urban school children in Delhi: the North India Myopia Study (NIM Study). *PLOS One.* 2017;12(12):e0189774.
- Wu L-J, Wang Y-X, You Q-S, et al. Risk factors of myopic shift among primary school children in Beijing, China: a prospective study. *Int J Med Sci.* 2015;12(8):633–638.
- Duursma E, Meijer A, De Bot K. The impact of home literacy and family factors on screen media use among Dutch preteens. *J Child Fam Stud.* 2017;26(2):612–622.
- Cheng C-Y, Yen M-Y, Lin H-Y, et al. Association of ocular dominance and anisometropic myopia. *Invest Ophthalmol Vis Sci.* 2004;45(8):2856–2860.
- Jiang F, Chen Z, Bi H, et al. Association between ocular sensory dominance and refractive error asymmetry. *PLoS One.* 2015;10(8):e0136222.
- Klaver C, Polling JR. Myopia management in the Netherlands. *Ophthalmic Physiol Opt.* 2020;40(2):230–240.