

## University of Rhode Island DigitalCommons@URI

---

Rhode Island Agricultural Experiment Station  
Bulletin

College of the Environment and Life Sciences

---

2013

# Leek Variety Trials, 2012


Rebecca Brown

*University of Rhode Island*, [brownreb@uri.edu](mailto:brownreb@uri.edu)

Noah Leclaire-Conway

*University of Rhode Island*, [noah.conway@gmail.com](mailto:noah.conway@gmail.com)

Follow this and additional works at: [http://digitalcommons.uri.edu/riaes\\_bulletin](http://digitalcommons.uri.edu/riaes_bulletin)

 Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Horticulture Commons](#), and the [Plant Breeding and Genetics Commons](#)

---

### Recommended Citation

Brown, Rebecca and Leclaire-Conway, Noah, "Leek Variety Trials, 2012" (2013). *Rhode Island Agricultural Experiment Station Bulletin*. Paper 4.

[http://digitalcommons.uri.edu/riaes\\_bulletin/4](http://digitalcommons.uri.edu/riaes_bulletin/4)

This Article is brought to you for free and open access by the College of the Environment and Life Sciences at DigitalCommons@URI. It has been accepted for inclusion in Rhode Island Agricultural Experiment Station Bulletin by an authorized administrator of DigitalCommons@URI. For more information, please contact [digitalcommons@etal.uri.edu](mailto:digitalcommons@etal.uri.edu).

LEEK VARIETY TRIALS 2012

---

---

**RESEARCH BULLETIN No. 3, 2013**

---

**Agricultural Experiment Station**

of the

**University of Rhode Island**

**Kingston, RI, 02881, U.S.A., February, 2013**

---



---

---

Publications of the Agricultural Experiment Station of the University of Rhode Island are made available free to the public through the Repository of the Digital Commons of the University of Rhode Island libraries at <http://digitalcommons.uri.edu/>

## Leek Variety Trials, 2012

Rebecca Brown and Noah Leclaire-Conway, Department of Plant Sciences and Entomology; funded by a Farm Viability Grant from RIDEM  
University of Rhode Island, Kingston, Rhode Island

The 2012 leek trial included 28 varieties, ranging from new F1 hybrids to traditional heirloom varieties. The leeks were seeded in the greenhouse March 17-30, and transplanted to the field April 17. In the field spacing was 15" between rows, and 5" between plants in the row. Leeks were hilled every few weeks with the hiller/furrower attachment on the walk-behind rototiller. Weeds were controlled by hand cultivation; fertilizer was incorporated at transplanting and irrigation was from overhead sprinklers. The trial developed Purple Blotch, caused by *Alternaria porri*, in early August. The disease continued to be problematic into September, despite repeated fungicide applications. Leeks were harvested beginning September 20<sup>th</sup>. Harvested plants were trimmed, counted, and weighed to determine average size. One plot of each variety was left in the field to evaluate the overwintering potential. The overwintered leeks were not mulched. Survival was rated on January 14<sup>th</sup> after repeated exposure to temperatures below freezing, and scattered overnight lows below 10°F.

The top-performing varieties in the trial were DP-12-02, Megaton, and Tadorna. DP-12-02 was an outstanding leek for fall harvest, but did not overwinter. Megaton and Tadorna were similar, except that Megaton was larger and more uniform, while Tadorna had better overwintering quality. Belton also did well, except for poor establishment.

Establishment ratings measured the survival of transplants in the field; it does not reflect percent seed germination. The variety Carentan had the best establishment, 100%, but extremely poor seed germination such that it had to be replanted in the greenhouse, and was not put into the field until mid-May. Tadorna, DP-12-02, and Megaton had the best establishment of the varieties transplanted in April.

Average size varied widely, from 5.6 oz. for Belton down to 0.7 oz. for Matisse. Data were not available for Matejko, as the harvested leeks were misplaced prior to trimming and weighing. Belton produced very large leeks, but it had poor seed germination and poor transplant survival, so the actual yield was low. Megaton and DP-12-02 had the highest yields due to the combination of large leeks and good establishment. DP-12-02 and Striker had the most uniformity among the leeks, but any variety with a score above 3.0 was statistically similar.

Vertical leaves are desirable in leeks as they decrease the amount of soil that accumulates in the leaf axils and thus give cleaner leeks. However, tight leaf axils can also provide a haven for thrips, and exacerbate damage. Runner, Surfer, and Megaton had the most upright leaves. Leaf color can also be important, as waxy, blue leaves are more attractive to thrips, but may be less susceptible to fungal diseases and more tolerant of freezing. Tadorna, 0050 TZ, Electra, and Lexton had very blue leaves, while DP-12-02 had glossy green leaves similar to many of the heirloom varieties. Tadorna had the best quality after overwintering in the field – the healthy blue plants were visible from a significant distance. Autumn Giant and Bandit also held well in the field. In general the varieties that survived overwintering had blue foliage, while those that were killed by hard freeze had glossy green foliage. Overwintering

data was not available for Belton, as the plot in the row that was left unharvested failed to establish. There were no differences among the varieties in the amount of damage from purple blotch.

### Leek variety performance data

Variety	Source	Establish. <sup>a</sup>	Size (oz.) <sup>b</sup>	Uniformity <sup>c</sup>	Leaf Angle <sup>d</sup>	Color <sup>e</sup>	Purple Blotch <sup>f</sup>	Over-wintering <sup>g</sup>
0050 TZ	Siegers	88%	2.4	3.0	2.3	4.0	1.3	3
98118 TZ	Siegers	83%	1.9	3.0	3.0	3.3	2.7	2
Autumn Giant	Baker Creek	94%	2.1	2.0	2.3	3.7	1.7	4
Bandit	High Mowing	65%	2.3	2.0	2.7	3.3	1.7	4
Belton	Siegers	42%	5.6	2.5	3.0	3.5	2.0	
Bleu de Solaize	Fedco	52%	1.1	1.7	2.0	3.3	2.0	2
Bulgarian Giant	Baker Creek	63%	2.0	1.7	1.7	2.0	2.0	1
Carentan	Baker Creek	100%	1.1	2.7	2.7	2.7	1.7	1
Conway	Siegers	83%	2.5	3.3	3.0	3.0	2.0	2
DP-12-02	dp seeds	97%	4.0	3.7	2.3	1.7	1.7	1
Electra	Harris	81%	2.9	3.0	2.0	4.0	2.3	2
Gevaria	Siegers	37%	2.9	1.7	2.0	3.0	1.7	2
Giant Musselburg	Baker Creek	92%	2.0	2.3	2.3	2.3	1.7	1
King Richard	Fedco	73%	1.2	2.0	2.7	3.0	2.7	3
King Sieg	Fedco	66%	3.7	2.7	2.0	2.3	1.3	1
Lancelot	JSS <sup>h</sup>	82%	2.7	2.3	2.7	3.7	1.7	2
Lexton	JSS	42%	2.3	2.3	2.3	4.0	1.0	3
Lincoln	Fedco	53%	3.7	3.0	2.7	2.0	1.3	2
Matejko	Siegers	64%		2.0	3.0	3.7	1.7	2
Matisse	Siegers	49%	0.7	1.7	2.3	3.3	1.3	3
Megaton	Siegers, JSS	97%	4.7	3.0	3.3	3.3	1.7	3
Oarsman	Siegers	61%	2.9	1.7	2.7	3.0	1.3	2
Pandora	JSS	40%	2.9	2.7	3.0	3.0	2.0	1
Runner	High Mowing, Bejo	75%	1.3	2.7	4.0	2.3	2.3	2
Striker	Bejo	92%	2.9	3.7	3.0	3.3	1.7	2
Surfer	Bejo	58%	1.5	2.7	3.7	3.7	2.0	2
Tadorna	Siegers, JSS	97%	3.3	3.3	2.7	4.0	2.0	5
Tornado	Siegers	44%	3.9	1.3	2.0	3.0	1.7	2

<sup>a</sup> Establishment refers to the % of transplants that established in the field.

<sup>b</sup> Size is the average weight after trimming and cleaning.

<sup>c</sup> Uniformity was rated on a scale of 1-4 with 4 indicating that the plants were highly uniform for size and color. Average of 3 reps; LSD = 1.7

<sup>d</sup> Leaf angle was rated on a scale of 1-4 where 1 indicates a leaf angle close to horizontal at the stem, and 4 indicates vertical leaves. Average of 3 reps; LSD = 0.9

<sup>e</sup> Foliage color ranged from glossy green (1) to blue (4). Average of 3 reps; LSD = 1.0

<sup>f</sup> Purple blotch was rated on a scale of 1-4 with 4 indicating heavy disease. There were no significant differences among varieties.

<sup>g</sup> Overwintering was evaluated in January 2013 after exposure to temperatures below 10°F. A score of 1 indicates that plants were completely dead, while a score of 5 indicates minimal damage.

<sup>h</sup> JSS is Johnny's Selected Seeds

THE BACTERIAL INFECTION OF FRESH EGGS

---

---

BULLETIN 164

---

Agricultural Experiment Station

OF THE

Rhode Island State College

KINGSTON, R. I., U. S. A., JANUARY, 1916

---



---

The publications of the Station are sent free upon request. Visitors are always welcome. Railway Station, Telegraph, Express and Post-Office—Kingston, R. I. Telephone Exchange, Narragansett Pier, No. 232-J.

---

PRESS OF E. L. FREEMAN COMPANY, PRINTERS TO THE STATE

*Scan of an original frontispiece from 1916. Courtesy of Special Collections, Robert L. Carothers Library, University of Rhode Island, Kingston, RI*