

University of Rhode Island DigitalCommons@URI

Graduate School of Oceanography Faculty **Publications** 

Graduate School of Oceanography

2006

# The Sedimentary Environment Below Earth's Polar Ice Cap as a Microbial Habitat

David C. Smith

Unviersity of Rhode Island. Graduate School of Oceanography, dcsmith@uri.edu

Stephanie Forschner University of Rhode Island

See next page for additional authors

Follow this and additional works at: https://digitalcommons.uri.edu/gsofacpubs Part of the Marine Biology Commons, and the Oceanography Commons



Terms of Use

All rights reserved under copyright.

### Citation/Publisher Attribution

Smith, David C.; Forschner, Stephanie; Rowley, David C.; and D'Hondt, Steven, "The Sedimentary Environment Below Earth's Polar Ice Cap as a Microbial Habitat" (2006). Graduate School of Oceanography Faculty Publications. Paper 1. https://digitalcommons.uri.edu/gsofacpubs/1https://digitalcommons.uri.edu/gsofacpubs/1

This Article is brought to you for free and open access by the Graduate School of Oceanography at DigitalCommons@URI. It has been accepted for inclusion in Graduate School of Oceanography Faculty Publications by an authorized administrator of DigitalCommons@URI. For more information, please contact digitalcommons@etal.uri.edu.

Authors David C. Smith, Stephanie Forschner, David C. Rowley, and Steven D'Hondt

# The Sedimentary Environment Below Earth's Polar Ice Cap as a Microbial Habitat

#### David C. Smith

Graduate School of Oceanography
University of Rhode Island
Narragansett, RI 02882
USA
dcsmith@gso.uri.edu

#### Stephanie Forschner

Department of Biomedical and Pharmaceutical Sciences
University of Rhode Island
USA

## David C. Rowley

Department of Biomedical and Pharmaceutical Sciences
University of Rhode Island
USA

#### Steven D'Hondt

Graduate School of Oceanography University of Rhode Island USA

Due to low energy and anaerobic conditions, deeply buried marine sediments provide an opportunity to study microbial diversity and adaptation to subsurface life under conditions that may mimic extraterrestrial subsurface conditions. The URI team of the NASA Astrobiology Institute recently participated in the first scientific drilling expedition to the ice-covered central Arctic Ocean (Integrated Ocean Drilling Program expedition 302). With an armada that consisted of three icebreakers, this expedition successfully cored the entire 428 m sediment stack on the Lomonosov Ridge during August and September 2004. The drill sites were located ~250 km from the North Pole under ~1300 m of water. The sediments accumulated over the past ~56 Ma. The recovered cores vary from siliciclastic sediment low in organic carbon (< 0.2 %) to organic-rich black sediments that rapidly accumulated in the early middle Eocene. Three geochemical environments were characterized based on chemical analyses of porewater: an upper ammonium oxidation zone, a carbonate dissolution zone and a deep (> 200 meters below sea floor) sulfate reduction zone. We are using phylogenetic markers to assess the diversity of microbes within each zone. We will analyze these data in the context of sediment lithology and porewater geochemistry, to document the environmental properties that control occurrences of these microorganisms and to guide sampling efforts for future subsurface exploration.